

SUMS OF STATIONARY RANDOM VARIABLES¹

ENDERS A. ROBINSON

A sequence $x(t)$ ($-\infty < t < \infty$, t an integer) of elements in Hilbert space is called stationary if the inner product $(x(t+s), x(t))$ does not depend upon t . If the Hilbert space is L^2 space with probability measure, then $x(t)$ is a random variable and the sequence $x(t)$ ($-\infty < t < \infty$) is called a second-order stationary random process. Let \mathbf{X} be the closed linear manifold spanned by all the elements of the stationary process. Then Kolmogorov [1] has shown that the equation $x(t)U = x(t+1)$, $-\infty < t < \infty$, uniquely determines the unitary operator U with domain and range \mathbf{X} . Using the von Neumann [2] spectral representation of U , we obtain the spectral representation of the random process

$$x(t) = \int_{-.5}^{.5} e^{2\pi i u t} dx(0)E(u), \quad -\infty < t < \infty.$$

The von Neumann [3] ergodic theorem, in the framework of Khintchine [4], is applicable, and shows that the average $\sum_1^n x(t)/n$ converges in the mean to the random variable $x(0)[E(0+) - E(0-)]$ as $n \rightarrow \infty$. In this paper we consider sums instead of averages; that is, we consider $\sum_1^n x(t)$, and establish the following theorem.

THEOREM. *Let the random variables $x(t)$ ($-\infty < t < \infty$, t an integer) be a second-order stationary random process with spectral distribution function $F(u)$. For variance $\{\sum_1^n x(t)\}$ to be bounded for all positive integers n , each of the following two conditions is necessary and sufficient:*

$$(1) \quad \int_{-.5}^{.5} \sin^{-2} \pi u dF(u) < \infty.$$

(2) *There is a second-order stationary random process*

$$y(t) \quad (-\infty < t < \infty) \text{ satisfying } y(t) - y(t+1) = x(t).$$

PROOF. (NECESSARY CONDITIONS). We are given that variance $\{\sum_1^n x(t)\} < B$ for all positive integers n . Without loss of generality we assume that the $x(t)$ are centered so that their mean values are zero. Then

Presented to the Society, February 28, 1959; received by the editors April 6, 1959.

¹ This paper was sponsored by the United States Army under Contract No. DA-11-022-ORD-2059 at the Mathematics Research Center, United States Army, Madison, Wisconsin.

$$\text{variance} \left\{ \sum_1^n x(t) \right\} = \left(\sum_1^n x(t), \sum_1^n x(t) \right).$$

From the spectral representation we have

$$\sum_1^n x(t) = \int_{-.5}^{.5} e^{2\pi i u} \frac{1 - e^{2\pi i u n}}{1 - e^{2\pi i u}} dx(0) E(u),$$

so

$$\text{variance} \left\{ \sum_1^n x(t) \right\} = \int_{-.5}^{.5} \frac{|1 - e^{2\pi i u n}|^2}{|1 - e^{2\pi i u}|^2} dF(u)$$

where $F(u) = \|x(0)E(u)\|^2$, $-.5 \leq u \leq .5$, is the spectral distribution function. Hence we have

$$B > \text{variance} \left\{ \sum_1^n x(t) \right\} = \left\{ \int_{-.5}^{0-} + \int_{0+}^{.5} \right\} \frac{\sin^2 \pi u n}{\sin^2 \pi u} dF(u) \\ + n^2 [F(0+) - F(0-)]$$

which shows that $F(0+) - F(0-)$ must vanish. Moreover, we have

$$B > \frac{1}{N} \sum_{n=1}^N \left\{ \int_{-.5}^{0-} + \int_{0+}^{.5} \right\} \frac{\sin^2 \pi u n}{\sin^2 \pi u} dF(u) \\ = \left\{ \int_{-.5}^{0-} + \int_{0+}^{.5} \right\} \left[\frac{1}{N} \sum_{n=1}^N \left(\frac{1}{2} - \frac{1}{2} \cos 2\pi u n \right) \right] \sin^{-2} \pi u dF(u).$$

Clearly the limit of the expression in brackets, as $N \rightarrow \infty$, is $1/2$, so $\int_{-.5}^{.5} \sin^{-2} \pi u dF(u)$ is finite. Q.E.D. (1).

The distribution function $F(u)$ defines a Lebesgue-Stieltjes measure on the real line segment $-.5 \leq u \leq .5$. Let \mathcal{W} denote the L^2 space of complex-valued measurable functions $\Phi(u)$ defined on $-.5 \leq u \leq .5$ for this measure. Define a correspondence between an element x of \mathcal{X} and an element $\Phi(u)$ of \mathcal{W} by

$$x = \int_{-.5}^{.5} \Phi(u) dx(0) E(u) \leftrightarrow \Phi(u).$$

Then Stone [5] and Kolmogorov [1] have shown that this correspondence establishes an isomorphism between \mathcal{X} and \mathcal{W} that preserves inner products. The function $e^{2\pi i u t} / (1 - e^{2\pi i u})$ belongs to \mathcal{W} since

$$\int_{-.5}^{.5} \left| \frac{e^{2\pi i u t}}{1 - e^{2\pi i u}} \right|^2 dF(u) = \frac{1}{4} \int_{-.5}^{.5} \sin^{-2} \pi u du < \infty.$$

If we define the element $y(t)$ of \mathbf{X} by the correspondence $y(t) \leftrightarrow e^{2\pi i u t} / (1 - e^{2\pi i u})$ we see that

$$y(t) - y(t+1) \leftrightarrow \frac{e^{2\pi i u t} - e^{2\pi i u (t+1)}}{1 - e^{2\pi i u}} = e^{2\pi i u t}.$$

But by the spectral representation, we know that $x(t) \leftrightarrow e^{2\pi i u t}$, and hence we have $y(t) - y(t+1) = x(t)$ for all integers t . Since

$$\begin{aligned} (y(t+s), y(t)) &= \int_{-.5}^{.5} \frac{e^{2\pi i u (t+s)} e^{-2\pi i u t}}{|1 - e^{2\pi i u}|^2} dF(u) \\ &= \frac{1}{4} \int_{-.5}^{.5} e^{2\pi i u s} \sin^{-2} \pi u dF(u) \end{aligned}$$

depends only on s , we see that $y(t)$ is a stationary random process. Q.E.D. (2).

PROOF. (SUFFICIENT CONDITIONS). Let condition (1) of the theorem be given. Since

$$\text{variance} \left\{ \sum_1^n x(t) \right\} = \int_{-.5}^{.5} \frac{\sin^2 \pi u n}{\sin^2 \pi u} dF(u) \leq \int_{-.5}^{.5} \sin^{-2} \pi u dF(u)$$

we see that the variance is bounded. Q.E.D. (1).

Let condition (2) of the theorem be given. Then $\|y(t)\|$ is a finite constant. Because $\sum_1^n x(t) = y(1) - y(n+1)$ we have $\left\| \sum_1^n x(t) \right\| \leq \|y(1)\| + \|y(n+1)\|$, and so $\text{variance} \left\{ \sum_1^n x(t) \right\} = \left\| \sum_1^n x(t) \right\|^2$ is bounded. Q.E.D. (2).

REFERENCES

1. A. N. Kolmogorov, *Stationary sequences in Hilbert space*, Bull. Moscow State Univ. Ser. Math. vol. 2 no. 6 (1941) 40 pp.
2. J. von Neumann, *Eigenwerttheorie Hermitescher Functionaloperatoren*, Math. Ann. vol. 102 (1929) pp. 49-131.
3. ———, *Proof of the quasiergodic hypothesis*, Proc. Nat. Acad. Sci. U.S.A. vol. 18 (1932) pp. 70-82.
4. A. Ja. Khintchine, *Korrelationstheorie der stationären stochastischen Prozesse*, Math. Ann. vol. 109 (1934) pp. 604-615.
5. M. H. Stone, *Linear transformations in Hilbert space*, Amer. Math. Soc. Colloquium Publications, vol. 15, 1932.

MATHEMATICS RESEARCH CENTER, UNITED STATES ARMY,
MADISON, WISCONSIN