

## A THEOREM OF LEVITZKI

I. N. HERSTEIN

Levitzki [2] proved that in a ring satisfying the ascending chain condition on left-ideals every left-ideal consisting of nilpotent elements is nilpotent. More recently, Goldie [1] has given a simpler proof. We present here a proof which is, in our opinion, even simpler and more elementary than that of Goldie. Moreover, it is completely self-contained.

**THEOREM (LEVITZKI).** *In a ring with ascending chain condition on left-ideals a nil left-ideal must be nilpotent.*

**PROOF.** We claim that we may assume that the ring  $R$  has no nonzero nilpotent left-ideals, for if it did, it would have a nonzero nilpotent two-sided ideal, hence a maximal such,  $N$ . If the result were false, in  $R/N$ , which has no nilpotent left-ideals we would have a nonzero nil left-ideal.

So suppose that  $R$  is a ring with ascending chain condition on left-ideals which has no nonzero nilpotent left-ideals and that  $A \neq (0)$  is a nil left-ideal of  $R$ . Let  $a \neq 0 \in A$ , and let  $\mathfrak{M} = \{ax \neq 0 \mid x \in R\}$ . For  $ax \in \mathfrak{M}$  let  $L(ax) = \{y \in R \mid yax = 0\}$ ; these give us a set of left-ideals of  $R$ , which, by the ascending chain condition has maximal elements. We denote these maximal elements by  $L(ax_i)$ .

If  $t \in R$  and  $ax_it \neq 0$  then  $L(ax_it) \supset L(ax_i)$  which, together with  $ax_it \in \mathfrak{M}$  forces  $L(ax_it) = L(ax_i)$ . We claim that given any finite number of such maximal  $ax_i, -ax_1, \dots, ax_n$ —then there is an element  $u \neq 0$  in  $Rax_1R$  such that  $ax_1u = \dots = ax_nu = 0$ . Suppose such a  $u$  has been found such that  $ax_1u = \dots = ax_{n-1}u = 0$ . If  $ax_nu = 0$  then we are done. So suppose any  $u \neq 0 \in Rax_1R$  annihilating  $ax_1, \dots, ax_{n-1}$  does not annihilate  $ax_n$ . We claim there is such a  $u \neq 0$  such that  $uax_n \neq 0$ ; for if  $uax_n = 0$  for all  $u \in Rax_1R$  which annihilate  $ax_1, \dots, ax_{n-1}$ , then since  $uR$  also does the trick,  $uRax_n = (0)$ , whence  $(Rax_nu)^2 = (0)$ . But then  $Rax_nu$  is a nilpotent left-ideal, so must be  $(0)$ . This forces  $ax_nu = 0$  the desired result. Thus we may suppose that  $uax_n \neq 0$ . Now  $(uax_n)^t = 0$  for some  $t$ , since  $x_nua \in A$  is nilpotent. If  $(uax_n)^t = 0$ ,  $(uax_n)^{t-1} \neq 0$ , then  $uax_n(uax_n)^{t-1} = 0$ . Since  $u \in L(ax_n(uax_n)^{t-1})$  but  $u \notin L(ax_n)$ , by the remark at the beginning of this paragraph,  $ax_n(uax_n)^{t-1} = 0$ . Thus  $(uax_n)^{t-1} \neq 0$  annihilates  $ax_1, \dots, ax_{n-1}$  and  $ax_n$ ; moreover since  $u \in Rax_1R$ ,  $(uax_n)^{t-1}$  also is.

Consider the ascending chain of left-ideals  $Rax_1, Rax_1 + Rax_2, \dots$ ,

---

Received by the editors February 27, 1961.

$Rax_1 + Rax_2 + \cdots + Rax_n, \cdots$ . It must terminate; thus for some  $n$ ,  $Rax_i \subset Rax_1 + \cdots + Rax_n$  for all  $ax_i$  such that  $L(ax_i)$  is maximal. Now  $ax_1 r$  is either 0 or a maximal  $ax_i$ ; whence  $Rax_1 R \subset Rax_1 + \cdots + Rax_n$ . But there is a  $u \neq 0$  in  $Rax_1 R$  such that  $ax_1 u = \cdots = ax_n u = 0$ . Thus  $Rax_1 Ru \subset Rax_1 u + \cdots + Rax_n u = (0)$ . Therefore, since  $Ru \neq 0 \subset Rax_1 R$ ,  $(Ru)^2 \subset Rax_1 R Ru = (0)$ . We have produced a nonzero nilpotent left-ideal in  $R$ ! This contradiction proves the theorem.

Clearly the proof would work if  $A$  were a nil right-ideal, for if  $0 \neq a \in A$ , then  $Ra$  would be a nonzero nil left-ideal of  $R$ .

#### REFERENCES

1. A. W. Goldie, *Semi-prime rings with maximum condition*, Proc. London Math. Soc. 10 (1960), 201-220.
2. Nathan Jacobson, *The structure of rings*, Amer. Math. Soc. Colloq. Publ. Vol. 37, Amer. Math. Soc., Providence, R. I., 1956.

CORNELL UNIVERSITY AND  
THE UNIVERSITY OF CHICAGO

---

## A REMARK ON DIRECT PRODUCTS OF MODULES

STEPHEN U. CHASE

It is a standard result of abelian group theory that the direct product of an infinite number of infinite cyclic groups is not free [3, p. 48]. Various generalizations of this fact to modules over an arbitrary ring are contained in [1]. In this note we restrict our attention to integral domains, and in this setting present a generalization of the aforementioned result which is similar to, but in some sense stronger than, those of [1]. We then apply our theorem to show that any factor group  $A$  of a direct product of infinite cyclic groups possesses the following amusing property: If  $A$  is a subgroup of a direct sum of reduced torsion-free abelian groups, then it must be contained in the direct sum of a finite number of them.

Throughout this discussion,  $R$  will be an integral domain which is not a field. All  $R$ -modules considered will be assumed to be unitary.

**LEMMA.** *Let  $A$  be a torsion-free  $R$ -module, and set  $A' = \bigcap_{\lambda \in R^*} \lambda A$ . Then  $A'$  is divisible. In particular, if  $A$  is reduced (i.e., has no divisible submodules) then  $\bigcap_{\lambda \in R^*} \lambda A = 0$ .<sup>1</sup>*

Received by the editors February 24, 1961

<sup>1</sup>  $R^*$  means  $R - \{0\}$ .