

A REMARK ON COMMUTABLE FUNCTIONS AND CONTINUOUS ITERATIONS

MAREK KUCZMA

The purpose of this note is to apply some recent results of B. Choczewski to a question concerning commutable functions and continuous iterations. Namely, we shall prove the following:

THEOREM. *Let $f(x)$ be a function of class C^2 in an interval (a, b) , such that $f(x) > x$ in (a, b) , $f(b) = b$, $\lim_{x \rightarrow a+0} f(x) = a$, $f'(x) > 0$ in (a, b) , and $f'(b) < 1$.¹ Moreover let us assume that the function $(d^2/dx^2)f^{-1}(x)$ satisfies a Lipschitz condition in a left neighbourhood of the point b ($f^{-1}(x)$ denotes the inverse function of $f(x)$, which exists since $f'(x) > 0$).*

Then there exists a unique one-parameter family of functions $\phi_s(x)$, $s \in (-\infty, \infty)$, satisfying the following conditions for all s :

- (A) $\phi_s(x)$ is of class C^2 in (a, b) .
- (B) $\phi_s(x) \in (a, b)$ for $x \in (a, b)$, $\phi_s(b) = b$.
- (C) $\phi_s(x)$ commutes with $f(x)$, i.e.

$$(1) \quad f[\phi_s(x)] = \phi_s[f(x)].$$

The family of the functions $\phi_s(x)$ is also, in a sense, the "best" family of continuous iterations of the function $f(x)$. The functions $\phi_s(x)$ may be obtained as the limit of the sequence defined by

$$(2) \quad \phi_{s,n}(x) = f^{-n}(\phi_{s,0}[f^n(x)]), \quad x \in (a, b), \quad n = 1, 2, 3, \dots,$$

where $f^k(x)$ denotes the k th iterate of the function $f(x)$, i.e., $f^0(x) = x$, $f^{k+1}(x) = f[f^k(x)]$, $f^{k-1}(x) = f^{-1}[f^k(x)]$, $k = 0, \pm 1, \pm 2, \dots$, and $\phi_{s,0}(x)$ is an arbitrary function of class C^2 in (a, b) such that

$$(3) \quad \phi_{s,0}(x) \in (a, b) \text{ for } x \in (a, b), \quad \phi_{s,0}(b) = b, \quad \phi'_{s,0}(b) = [f'(b)]^s.$$

Furthermore, if the function $f(x)$ is of class C^r ($r > 2$) in (a, b) , then the functions $\phi_s(x)$ are also of class C^r in (a, b) .

PROOF. In order to simplify the notation we write

$$h(x) = \frac{df}{f^{-1}(x)}.$$

Received by the editors October 30, 1961.

¹ The number a may also equal $-\infty$; but b must be finite, which undoubtedly is a defect of the present theorem. The condition $\lim_{x \rightarrow a+0} f(x) = a$ may be omitted, but then the functions $\phi_s(x)$ occurring in the thesis of the theorem may not be defined in the whole interval (a, b) , but only in a left neighbourhood of the point b . The conditions $f(x) > x$ in (a, b) , $f'(b) < 1$, may be replaced by $f(x) < x$ in (a, b) , $f'(b) > 1$. And naturally, the endpoints, a and b , of the interval in question may be interchanged.

Then equation (1) takes the form

$$(4) \quad \phi(x) = h(\phi[f(x)]).$$

Moreover, we have

$$(5) \quad h'(b)[f'(b)]^2 = f'(b) < 1.$$

Consequently, as a direct consequence of a theorem of B. Choczewski [2], it follows that for every number $s \in (-\infty, \infty)$ there exists exactly one function $\phi_s(x)$ satisfying equation (4), and fulfilling conditions (A), (B), and the condition²

$$(6) \quad \phi'_s(b) = [f'(b)]^s.$$

(Condition (C) is fulfilled, since the functions $\phi_s(x)$ satisfy equation (4), which is equivalent to (1)).

It follows from the uniqueness of ϕ_s that $\phi_k(x) = f^k(x)$ for integral k . In particular we have $\phi_1(x) = f(x)$.

The paper by B. Choczewski has not yet been published; and since the method employed by him is essential to our further considerations, we shall present here a very short outline of his arguments, naturally omitting all details.³

Let \mathfrak{R} be the space of all functions $\phi(x)$ of class C^2 in the interval $\langle b-\eta, b \rangle$ (where η is a sufficiently small, fixed positive number, depending only on $f(x)$ and s), fulfilling the conditions:

$$(7) \quad \begin{aligned} \phi(x) \in \langle a + \alpha, b \rangle \text{ for } x \in \langle b - \eta, b \rangle, \quad \phi(b) = b, \quad \phi'(b) = [f'(b)]^s \\ |\phi''(x)| \leq K \quad \text{for } x \in \langle b - \eta, b \rangle, \end{aligned}$$

(where α and K are suitably chosen positive numbers, depending only on $f(x)$ and s), and endowed with the metric

$$\rho[\phi, \psi] = \sup_{\langle b-\eta, b \rangle} |\phi''(x) - \psi''(x)|,$$

(with which \mathfrak{R} becomes a complete metric space). Consider the transformation

$$\Phi[\phi] = h(\phi[f(x)]).$$

This transformation (under a suitable choice of η , α and K) maps \mathfrak{R}

² Of course, for $s \in (-\infty, \infty)$ the values of $[f'(b)]^s$ run over the set of all positive numbers. However, in order that $\phi_s(x)$ form a family of continuous iterations of $f(x)$ (cf. (9) below), it is convenient to define $\phi_s(x)$ with the aid of relation (6).

³ In his paper [2] B. Choczewski deals with the more general equation $\phi(x) = H(x, \phi[f(x)])$.

into itself. Furthermore, in view of (5), we have, for $\phi, \psi \in \mathfrak{R}$

$$\rho[\Phi[\phi], \Phi[\psi]] < \vartheta \rho[\phi, \psi], \quad \vartheta < 1.$$

Consequently, Φ is a contraction mapping, and by the theorem of Banach-Caccioppoli, has exactly one fixed point in \mathfrak{R} , i.e. there exists exactly one function $\phi_s(x)$ of class C^2 in $\langle b-\eta, b \rangle$, satisfying equation (4) and conditions (7). This function can be uniquely extended onto the whole interval (a, b) in such a manner that it will satisfy equation (4).

Hence it follows (see also [3, Theorem V]) that the function $\phi_s(x)$ can be obtained as the limit of the sequence defined by the recurrence formula

$$(8) \quad \phi_{s,n+1}(x) = h(\phi_{s,n}[f(x)]), \quad n = 0, 1, 2, \dots,$$

where $\phi_{s,0}(x)$ is an arbitrary function which, when restricted to $\langle b-\eta, b \rangle$, belongs to \mathfrak{R} , and thus e.g. an arbitrary function of class C^2 in (a, b) , fulfilling conditions (3). One can easily verify that formula (2) follows immediately from (8).

From the theorems of B. Choczewski [2] it further follows that the function $\phi_s(x)$ has the same regularity properties as $f(x)$.

To complete the proof of our theorem it remains only to show that the functions $\phi_s(x)$ form a family of continuous iterations of $f(x)$, i.e. that they satisfy the relation

$$(9) \quad \phi_s[\phi_t(x)] = \phi_{s+t}(x), \quad x \in (a, b), \quad s, t \in (-\infty, \infty).$$

Now, the function $\phi_s[\phi_t(x)]$, just like $\phi_s(x)$ and $\phi_t(x)$, is of class C^2 in (a, b) and evidently commutes with $f(x)$. We have also $\phi_s[\phi_t(x)] \in (a, b)$ for $x \in (a, b)$, $\phi_s[\phi_t(b)] = b$, $(d/dx)\phi_s[\phi_t(x)]|_{x=b} = \phi'_s[\phi_t(b)]\phi'_t(b) = \phi'_s(b)\phi'_t(b) = [f'(b)]^{s+t}$. But the function with these properties is unique. Thus $\phi_s[\phi_t(x)] = \phi_{s+t}(x)$, which was to be proved.

REMARK. The above theorem is closely related to some results of L. Berg [1]. The results obtained by Berg are, in some respect, stronger than ours—the condition $b < \infty$ does not occur, and the case $f'(b) = 1$ is also discussed. On the other hand, in most of the theorems in [1] assumptions of analyticity, or at least of multiple differentiability of $f(x)$, as well as the assumption that $f''(x)$ has a constant sign, occur; while in our case the assumptions regarding the regularity of $f(x)$ are weaker.

Similar equations have been also investigated by S. Sternberg [5]. He has obtained similar results, however in a different way.

It follows from [4] that in condition (A) it is not sufficient to re-

quire only the continuity of $\phi_s(x)$: for equation (1) admits continuous solution depending on an arbitrary function. Unfortunately, however, our method does not allow us to decide whether in addition to $\phi_s(x)$ there also exist other functions of class C^1 in (a, b) and fulfilling conditions (B) and (C). (For the linear function; $f(x) = b + q(x = b)$ such functions do not exist; cf. [1, §1].)

REFERENCES

1. L. Berg, *Iterationen von beliebiger Ordnung*, Z. Angew. Math. Mech. **40** (1960), 215–229.
2. B. Choczewski, *Investigation of the existence and uniqueness of differentiable solutions of a functional equation*, Ann. Polon. Math. (to appear).
3. J. Kordylewski and M. Kuczma, *On the functional equation $F(x, \phi(x), \phi[f(x)]) = 0$* , Ann. Polon. Math. **7** (1959), 21–32.
4. M. Kuczma, *On continuous solutions of a functional equation*, Ann. Polon. Math. **8** (1960), 209–214.
5. S. Sternberg, *Local C^n transformations of the real line*, Duke Math. J. **24** (1957), 97–102.

UNIwersytet Jagielloński, Kraków, Poland