

ON THE PRODUCT OF DIRECTED GRAPHS

M. H. McANDREW

1. Introduction. The concept of the product of two graphs [1, p. 23] has application in the theory of games and the theory of automata. Weichsel [2] deduced a necessary and sufficient condition for the product of two connected undirected graphs to be connected; he further proves that the product graph, if not connected, has exactly two components. In this paper we deduce the corresponding results for directed graphs, namely that if G_1, G_2, \dots, G_r are strongly connected directed graphs then $G_1 \times G_2 \times \dots \times G_r$ has exactly

$$\frac{d(G_1) \cdot d(G_2) \cdot \dots \cdot d(G_r)}{\text{lcm}(d(G_1), d(G_2), \dots, d(G_r))}$$

components where $d(G)$ is as defined below.

2. Definitions. We define a *directed graph* G to be a set of nodes $N(G)$ and a set of edges $E(G)$, the elements of $E(G)$ being ordered pairs (p, q) with $p, q \in N$.

We write $p \Rightarrow q$ if $(p, q) \in E$.

We define a *path* \mathcal{P} of G as a finite sequence $\{p_i\}$, $(0 \leq i \leq n, n \geq 1)$ such that $p_i \Rightarrow p_{i+1}$, $(0 \leq i \leq n-1)$. We define $n(\mathcal{P}) = n$, the number of edges in \mathcal{P} and write $\mathcal{P}(p, q)$ for a path with $p_0 = p, p_n = q$.

By $p \in \mathcal{P}$ we mean \mathcal{P} is the path $\{p_i\}$ and for some $i, p_i = p$.

We write $p \rightarrow q$ if either $p = q$ or there is a path $\mathcal{P}(p, q)$.

We say p, q are *strongly connected* if $p \rightarrow q$ and $q \rightarrow p$. This is clearly an equivalence relation; we define a component C of G to be a subgraph whose nodes, $N(C)$, are just the nodes in an equivalence class with respect to this relation and whose edges are those edges (p, q) of G with $p, q \in N(C)$.

We say G is *strongly connected* if each pair of nodes of G are strongly connected.

If $\mathcal{P}_1(p_1, p_2)$ and $\mathcal{P}_2(p_2, p_3)$ are paths of G , we define $\mathcal{P}_1 + \mathcal{P}_2$ to be the path $\mathcal{P}_3(p_1, p_3)$ obtained by concatenating the sequences of nodes in \mathcal{P}_1 and \mathcal{P}_2 . This sum is clearly associative and $n(\mathcal{P}_3) = n(\mathcal{P}_1) + n(\mathcal{P}_2)$.

We define a cycle \mathcal{C} to be a path $\mathcal{P}(p, p)$.

We define $d(G) = \text{gcd}(n(\mathcal{C}))$, taken over all cycles of G .

We define the product $G \times H$ of two graphs G, H as follows

(i) $N(G \times H) = \{(p, q) \mid p \in N(G) \text{ and } q \in N(H)\}$,

Received by the editors May 26, 1962.

(ii) $E(G \times H) = \{((p_1, q_1), (p_2, q_2)) \mid (p_1, p_2) \in E(G) \text{ and } (q_1, q_2) \in E(H)\}$.

It may be readily seen that the adjacency matrix of $G \times H$ is just the Kronecker product of the adjacency matrices of G and H . Since the product is clearly associative we may define unambiguously $G_1 \times G_2 \times \dots \times G_r$.

3. Preliminary lemmas.

LEMMA 1. *Let $\mathcal{P}_1(p, q), \mathcal{P}_2(p, q)$ be paths of a strongly connected graph G . Then*

$$d(G) \mid (n(\mathcal{P}_1) - n(\mathcal{P}_2)).$$

PROOF. Since G is strongly connected there is a path $\mathcal{P}_3(q, p)$. Then $\mathcal{P}_1 + \mathcal{P}_3$ and $\mathcal{P}_2 + \mathcal{P}_3$ are cycles; hence $d(G) \mid n(\mathcal{P}_i + \mathcal{P}_3) = n(\mathcal{P}_i) + n(\mathcal{P}_3)$, ($i = 1, 2$). The result of the lemma now follows.

Let G be a strongly connected graph and let Z_d be the ring of residue classes mod $d = d(G)$. We may define a function $f: N(G) \rightarrow Z_d$ as follows: choose some node p_0 of G . Then for any $p \in N(G)$ define $f(p) = n(\mathcal{P}(p_0, p))$ for any path $\mathcal{P}(p_0, p)$. Since G is strongly connected there is some such path and Lemma 1 ensures that $f(p)$ is single valued.

LEMMA 2. *Let G be a strongly connected graph and f be defined as above. Then for any path $\mathcal{P}(p, q)$,*

$$n(\mathcal{P}) \equiv f(p) - f(q) \pmod{d(G)}.$$

PROOF. Let $\mathcal{P}_1(p_0, p)$ be a path and let $\mathcal{P}_2(p_0, q) = \mathcal{P}_1(p_0, p) + \mathcal{P}(p, q)$. By definition of f ,

$$n(\mathcal{P}_1) \equiv f(p) \pmod{d(G)}$$

and

$$n(\mathcal{P}_2) \equiv f(q) \pmod{d(G)}.$$

The result of the lemma now follows since $n(\mathcal{P}_2) = n(\mathcal{P}) + n(\mathcal{P}_1)$.

LEMMA 3. *Let G be a strongly connected graph and let $d = d(G)$. Then there exists a finite set of cycles $\mathcal{C}_1, \dots, \mathcal{C}_t$ such that*

$$\gcd_{1 \leq i \leq t} (n(\mathcal{C}_i)) = d.$$

PROOF. We may construct the set $\mathcal{C}_1 \dots \mathcal{C}_t$ inductively. Suppose $\mathcal{C}_1, \dots, \mathcal{C}_r$ already chosen with $\gcd_{1 \leq i \leq r} (n(\mathcal{C}_i)) = d_r > d$. Then there exists some \mathcal{C} with $d_r \nmid n(\mathcal{C})$. Let this be \mathcal{C}_{r+1} ; we now have $d_r > d_{r+1}$.

Since $\{d_r\}$ is a strictly monotonically decreasing sequence it must reach d in a finite number of steps.

LEMMA 4. *Let G be a strongly connected graph; let $d=d(G)$. Let $p_1, p_2 \in N(G)$. Then there is an integer n_0 such that for all $n > n_0$ there is a path $\mathcal{O}(p_1, p_2)$ with*

$$(1) \quad n(\mathcal{O}) = nd + f(p_2) - f(p_1).$$

PROOF. Let $\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_t$ be a set of cycles with $\gcd_{1 \leq i \leq t} (n(\mathcal{C}_i)) = d$. The existence of such a set is assured by Lemma 3. Let $n_i = n(\mathcal{C}_i)$. By definition of the gcd, there exist integers u_1, u_2, \dots, u_t such that

$$(2) \quad \sum_{i=1}^t u_i n_i = d.$$

Let $\mathcal{C}_i = \mathcal{O}_i(q_i, q_i)$. Choose some set of paths $\mathcal{O}'_0(p_1, q_1), \mathcal{O}'_1(q_1, q_2), \dots, \mathcal{O}'_t(q_t, p_2)$. Let $n(\sum_{i=0}^t \mathcal{O}'_i) = \sum_{i=0}^t n(\mathcal{O}'_i) = K$. Then by Lemma 2,

$$(3) \quad K = ad + f(p_2) - f(p_1)$$

for some integer a .

Let $u = \max_i (|u_i|)$. Let $s = \sum_{i=1}^t n_i/d$; s is integral since $d|n_i$ for every i . We will show that $n_0 = a + s + s^2u$ satisfies the requirements of the lemma. Let $n > n_0$. Then for some integral b, c

$$(4) \quad n - a = bs + c$$

with

$$(5) \quad 0 \leq c < s.$$

Let

$$(6) \quad v_i = b + cu_i.$$

Then

$$\begin{aligned} v_i &\geq \left(\frac{n-a}{s} - 1 \right) - us, && \text{from (4) and (5)} \\ &\geq \frac{n_0 - a}{s} - 1 - us, && \text{since } n > n_0 \\ &= 0 && \text{by definition of } n_0. \end{aligned}$$

I.e., the v_i are non-negative integers. Hence we may construct a path

$$\begin{aligned} \mathcal{O}(p_1, p_2) &= \mathcal{O}'_0(p_1, q_1) + v_1 \mathcal{O}_1(q_1, q_1) + \mathcal{O}'_1(q_1, q_2) + \dots \\ &\quad + v_t \mathcal{O}_t(q_t, q_t) + \mathcal{O}'_t(q_t, p_2). \end{aligned}$$

Then

$$\begin{aligned}
n(\mathcal{P}) &= \sum_{i=0}^t n(\mathcal{P}'_i) + \sum_{i=1}^t v_i n(\mathcal{P}_i) \\
&= K + \sum_{i=1}^t v_i n_i \\
&= ad + f(p_2) - f(p_1) + b \sum_{i=1}^t n_i + \sum_{i=1}^t cu_i n_i, \text{ from (3) and (6)} \\
&= ad + f(p_2) - f(p_1) + bsd + cd, \text{ by (2) and the definition of } s \\
&= nd + f(p_2) - f(p_1).
\end{aligned}$$

This completes the proof of the lemma.

4. Main theorems.

THEOREM 1. *Let G and H be strongly connected graphs. Let $d_1 = d(G)$, $d_2 = d(H)$, $d_3 = \gcd(d_1, d_2)$ and $D = \text{lcm}(d_1, d_2)$. Then, in $G \times H$,*

(i) $(p, q) \rightarrow (p', q')$ if and only if

$$f_G(p) - f_H(q) \equiv f_G(p') - f_H(q') \pmod{d_3}.$$

(ii) *The number of components of $G \times H$ is d_3 .*

(iii) *If C is any component of $G \times H$ then $d(C) = D$.*

PROOF. (i) If $(p, q) \rightarrow (p', q')$ then there exists a path $\{(p_i, q_i)\}$ with $(p_0, q_0) = (p, q)$ and $(p_n, q_n) = (p', q')$. Then $\{p_i\}$ and $\{q_i\}$ are paths $\mathcal{P}, \mathcal{P}'$ of G and H respectively with $n(\mathcal{P}) = n(\mathcal{P}') = n$. Hence, by Lemma 1,

$$n \equiv f_G(p') - f_G(p) \pmod{d_1}$$

and

$$n \equiv f_H(q') - f_H(q) \pmod{d_2}.$$

Therefore, since $d_3 | d_1, d_2$,

$$(7) \quad f_G(p') - f_G(p) \equiv f_H(q') - f_H(q) \pmod{d_3}.$$

Conversely, if (7) is satisfied, let n_1 be a value of n_0 satisfying the conclusions of Lemma 4 for G, p, p' ; let n_2 be defined similarly for H, q, q' . Let a, b be integers such that $ad_1 - bd_2 = d_3$. Let

$$(8) \quad \frac{f_G(p) - f_H(q) - f_G(p') + f_H(q')}{d_3} = u,$$

which is integral by hypothesis. Choose r so that

$$(au + rd_2) \geq n_1$$

and

$$(bu + rd_1) \geq n_2.$$

Then by Lemma 4 there is a path $\mathcal{P}_1(p, p')$ of G of length

$$n(\mathcal{P}_1) = (au + rd_2)d_1 + f_G(p') - f_G(p)$$

and a path $\mathcal{P}_2(q, q')$ of H of length

$$n(\mathcal{P}_2) = (bu + rd_1)d_2 + f_H(q') - f_H(q).$$

Then

$$\begin{aligned} n(\mathcal{P}_2) - n(\mathcal{P}_1) &= (ad_1 - bd_2)u - ud_3, & \text{by (8)} \\ &= 0. \end{aligned}$$

Hence $\mathcal{P}_1 \equiv \{p_i\}$ and $\mathcal{P}_2 \equiv \{q_i\}$ are paths of the same length and so $\{(p_i, q_i)\}$ is a path of $G \times H$ whose end-nodes are just (p, q) and (p', q') , i.e., $(p, q) \rightarrow (p', q')$ in $G \times H$.

(ii) Let $l(p, q) = f_G(p) - f_H(q)$, reduced mod d_3 . Then, by (i), nodes (p, q) and (p', q') are strongly connected if and only if $l(p, q) = l(p', q')$. For any residue $x \pmod{d_3}$ there is certainly some node (p, q) with $l(p, q) = x$. Hence the number of components of $G \times H$ is just d_3 , each component consisting of just these nodes with the same value of $l(p, q)$.

(iii) Let $(p, q), (p', q')$ be nodes of $G \times H$ in the same component, C . Let $d_0 = d(C)$. By Lemmas 1 and 4 the lengths of paths $\mathcal{P}(p, p')$ in G consist of all but a finite number of the integers

$$(9) \quad nd_1 + f_G(p') - f_G(p) \quad \text{for } n > 0.$$

Similarly the lengths of paths $\mathcal{P}(q, q')$ in H consist of all but a finite number of the integers

$$(10) \quad nd_2 + f_H(q') - f_H(q) \quad \text{for } n > 0.$$

The integers common to the arithmetic series (9) and (10) form an arithmetic series of difference $D = \text{lcm}(d_1, d_2)$. Hence the lengths of paths $\mathcal{P}((p, q), (p', q'))$ in C consist of all but a finite number of the integers of this series; but applying Lemmas 1 and 4 to $G \times H$ the lengths of these paths consist of all but a finite number of the integers

$$nd(C) + f_C(p', q') - f_C(p, q), \quad (n > 0).$$

Therefore $d(C) = D = \text{lcm}(d_1, d_2)$.

THEOREM 2. *Let G_1, G_2, \dots, G_s be strongly connected graphs. Let $d_i = d(G_i)$ ($i = 1, 2, \dots, s$). Then in $G_1 \times G_2 \times \dots \times G_s$:*

- (i) *The number of components is $\prod d_i / \text{lcm}(d_i)$.*
 (ii) *Each component C has $d(C) = \text{lcm}(d_i)$.*

PROOF. The results are true trivially for $s=1$. We shall complete the proof by induction. Suppose the results are true for $s=r$; i.e., $H \equiv G_1 \times G_2 \times \cdots \times G_r$ has m components, each component C having $d(C) = d_H$, where

$$d_H = \text{lcm}_{1 \leq i \leq r} (d_i)$$

and

$$m = \frac{\prod_{i=1}^r d_i}{d_H}.$$

In $H \times G_{r+1}$ nodes (p, q) and (p', q') are strongly connected only if p and p' are strongly connected in H . Hence the components of $H \times G_{r+1}$ consist of just the components of $C \times G_{r+1}$ for all components C of H . To each such product $C \times G_{r+1}$ we may apply the results of Theorem 1. Each component, C' , of $C \times G_{r+1}$ has

$$\begin{aligned} d(C') &= \text{lcm}(d_H, d_{r+1}) \\ &= \text{lcm}(\text{lcm}(d_1, d_2, \dots, d_r), d_{r+1}) \\ &= \text{lcm}(d_1, d_2, \dots, d_{r+1}). \end{aligned}$$

The number of components of $C \times G$ is

$$\text{gcd}(d_H, d_{r+1}) = \frac{d_H \cdot d_{r+1}}{\text{lcm}(d_H, d_{r+1})}.$$

Hence the number of components of $H \times G_{r+1}$

$$\begin{aligned} &= m \frac{d_H \cdot d_{r+1}}{\text{lcm}(d_H, d_{r+1})} \\ &= \frac{\prod_{i=1}^r d_i}{d_H} \cdot \frac{d_H \cdot d_{r+1}}{\text{lcm}(d_H, d_{r+1})} \\ &= \frac{\prod_{i=1}^{r+1} d_i}{\text{lcm}(d_1, d_2, \dots, d_{r+1})}. \end{aligned}$$

Hence the results of the theorem are true for $s=r+1$. The theorem

now follows by induction.

We may apply the results of Theorems 1 and 2 to undirected graphs by replacing each edge (p, q) of the undirected graph by a pair of edges $(p, q), (q, p)$ and constructing an equivalent directed graph. For such a graph G , $d(G)$ is 1 or 2 according as G has or has not an odd cycle. Theorem 1 of [1] now follows from Theorem 1 above. From Theorem 2 follows directly

THEOREM 3. *Let G_1, G_2, \dots, G_r be undirected graphs. If exactly s of these graphs have an odd cycle then m , the number of components of $G_1 \times G_2 \times \dots \times G_r$ is given by*

$$m = \begin{cases} 1 & s \geq r - 1 \\ 2^{r-s} & s \leq r - 1. \end{cases}$$

BIBLIOGRAPHY

1. Claude Berge, *Théorie des graphes et ses applications*, Dunod, Paris, 1958.
2. P. M. Weichsel, *The Kronecker product of graphs*, Proc. Amer. Math. Soc. **13** (1962), 47-52.

INTERNATIONAL BUSINESS MACHINES CORPORATION