

A NOTE ON AN INEQUALITY OF M. MARCUS
AND M. NEWMAN

HENRYK MINC¹

The permanent of an n -square matrix $A = (a_{ij})$ is denoted by $p(A)$ and is defined by

$$(1) \quad p(A) = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i\sigma(i)},$$

where S_n is the symmetric group of degree n . A real matrix S is said to be doubly stochastic if its entries are nonnegative and its row sums and column sums are all 1.

In a recent paper [2] Marcus and Newman proved that if A is a symmetric positive semi-definite doubly stochastic matrix then

$$(2) \quad p(A) \geq n!/n^n$$

with equality if and only if $A = J_n$, the matrix all of whose entries are $1/n$. This result is a partial answer to a conjecture of van der Waerden [3] stating that if S is any doubly stochastic matrix then $p(S) \geq p(J_n) = n!/n^n$ with equality if and only if $S = J_n$.

In the present note we extend the result of Marcus and Newman to all positive semi-definite hermitian matrices which have $e = (1, 1, \dots, 1)$ as a characteristic vector and prove it by a new method. We first establish a lemma which is a weakened version of Theorem 2 in [1] and use it in conjunction with the following inequality also due to Marcus and Newman (Theorem 5 in [2]).

If A and B are complex n -square matrices then

$$(3) \quad |p(AB)|^2 \leq p(AA^*)p(B^*B)$$

with equality if and only if either (a) a row of A or a column of B is zero, or (b) there exist a diagonal matrix G and a permutation matrix P such that $A^* = BGP$.

Let Φ_n denote the set of positive semi-definite hermitian n -square matrices which have e as a characteristic vector. In other words, Φ_n contains all nonnegative multiples of positive semi-definite hermitian n -square matrices whose row sums are all 1. Thus if $H \in \Phi_n$ then $HJ_n = \lambda_1 J_n$ for some nonnegative λ_1 .

Presented to the Society, November 17, 1962; received by the editors September 17, 1962.

¹ This research was supported by the Air Force Office of Scientific Research.

LEMMA. For any H in Φ_n there exists a matrix B in Φ_n such that $B^2 = H$.

PROOF. Let all the row sums of H be equal to $\lambda_1 \geq 0$. Then λ_1 is a characteristic root of H and e is a corresponding characteristic vector. There exists therefore a unitary matrix U with e/\sqrt{n} as its first column vector and such that

$$U^*HU = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n),$$

where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the nonnegative characteristic roots of H . Let $D = \text{diag}(\sqrt{\lambda_1}, \sqrt{\lambda_2}, \dots, \sqrt{\lambda_n})$ and let $B = UDU^*$. Clearly B is positive semi-definite hermitian and $B^2 = H$. It remains to prove that e is a characteristic vector of B . Let $X^{(1)}$ denote the first column vector of a matrix X . Then

$$U^*e = U^*\sqrt{n}U^{(1)} = \sqrt{n}(U^*U)^{(1)} = \begin{pmatrix} \sqrt{n} \\ 0 \\ \vdots \\ 0 \end{pmatrix}.$$

Hence $XU^*e = \sqrt{n}X^{(1)}$ for any n -square matrix X . In particular

$$Be = UDU^*e = \sqrt{n}(UD)^{(1)} = \sqrt{n}UD^{(1)} = \sqrt{(\lambda_1 n)}U^{(1)} = \sqrt{\lambda_1}e.$$

THEOREM. If $H \in \Phi_n$ and the row sums of H are all equal to λ_1 then

$$p(H) \geq n!(\lambda_1/n)^n$$

with equality if and only if either a row of H is zero or H is a nonnegative multiple of J_n .

PROOF. By the lemma, $H = B^2$ where $B \in \Phi_n$ and therefore $B = B^*$ and $BJ_n = \sqrt{\lambda_1}J_n$. Hence, by (3),

$$|p(BJ_n)|^2 \leq p(BB^*)p(J_n^*J_n),$$

i.e.,

$$(p(\sqrt{\lambda_1}J_n))^2 \leq p(H)p(J_n)$$

and therefore

$$p(H) \geq \lambda_1^n p(J_n) = n!(\lambda_1/n)^n.$$

Equality will occur (see conditions for equality in (3)) if and only if either (a) a row of B , and therefore a row of H , is zero, or (b) there exist a diagonal matrix G and a permutation matrix P such that

$B = J_n G P$. In case (b), $H = BB^* = J_n G^2 J_n$ and clearly $J_n G^2 J_n = (\text{tr}(G^2)/n) J_n$.

COROLLARY. *If S is a symmetric positive semi-definite doubly stochastic n -square matrix then*

$$p(S) \geq n!/n^n$$

with equality if and only if $S = J_n$.

REFERENCES

1. Marvin Marcus and Henryk Minc, *Some results on doubly stochastic matrices*, Proc. Amer. Math. Soc. **13** (1962), 571-579.
2. Marvin Marcus and Morris Newman, *Inequalities for the permanent function*, Ann. of Math. (2) **75** (1962), 47-62.
3. B. L. van der Waerden, *Aufgabe 45*, Jber. Deutsch. Math.-Verein. **35** (1926), 117.

UNIVERSITY OF FLORIDA