

**ON THE CORRECTNESS OF PARTIAL DIFFERENTIAL  
OPERATORS AND THE VON NEUMANN CONDITION  
FOR STABILITY OF FINITE DIFFERENCE  
OPERATORS<sup>1</sup>**

D. G. ARONSON

1. **Introduction.** Let  $P(\xi)$  be an  $N \times N$  matrix whose elements are polynomials of degree  $\leq m$  in  $\xi = (\xi_1, \dots, \xi_n)$  and let

$$L \equiv P(D) - E\partial/\partial t,$$

where  $D = (\partial/\partial x_1, \dots, \partial/\partial x_n)$  and  $E$  is the  $N \times N$  identity matrix. The operator  $L$  is said to be *correct in the sense of Petrovskiĭ* [3] if there exists a constant  $K > 0$  such that

$$(1.1) \quad \max_j \operatorname{Re} \mu_j(\xi) \leq K \quad \text{for all } \xi \in E^n,$$

where the  $\mu_j$  ( $j = 1, 2, \dots, N$ ) are the eigenvalues of  $P(i\xi)$ . We shall give a necessary and sufficient condition for the correctness of  $L$  in terms of finite difference approximations to  $L$ .

Let  $h$  and  $\tau$  be arbitrary positive numbers, and let

$$R_{h,\tau} = \{(jh, l\tau) \mid j \in I^n, l = 0, 1, \dots\},$$

where  $I^n$  is the set of all  $n$ -tuples of integers. As approximations to  $L$  we shall consider the two-parameter family of *two-level finite difference operators*

$$L_{h,\tau}u(x, t) \equiv \sum_r B_r^{(0)}u(x + rh, t) + \sum_r B_r^{(1)}u(x + rh, t + \tau)$$

for  $(x, t) \in R_{h,\tau}$ , where the  $B_r^{(i)}$  are  $N \times N$  matrices which depend only on  $h$  and  $\tau$ ,  $r \in I^n$ , and  $B_r^{(i)} = 0$  outside some fixed finite neighborhood of  $r = 0$ . The family of finite difference operators  $\{L_{h,\tau}\}$  is said to be *consistent with  $L$*  if

$$(1.2) \quad (L_{h,\tau} - L)u(x, t) \rightarrow 0 \quad \text{as } h, \tau \rightarrow 0$$

for all sufficiently smooth  $u$  [7].

For any  $\alpha \in E^n$ ,  $L_{h,\tau}u = 0$  has a solution of the form  $u(x, t) = e^{i\alpha^T x} q(t)$  if and only if  $q$  satisfies

Received by the editors July 16, 1962.

<sup>1</sup> This work was supported in part by the Office of Naval Research and in part by the National Science Foundation under a Postdoctoral Fellowship.

$$\left( \sum_r B_r^{(0)} e^{i\alpha^T r h} \right) q(t) + \left( \sum_r B_r^{(1)} e^{i\alpha^T r h} \right) q(t + \tau) = 0$$

for  $t = l\tau$ . Let

$$H_j(h, \tau, \alpha) = \sum_r B_r^{(j)} e^{i\alpha^T r h} \quad (j = 0, 1).$$

The family of finite difference operators  $\{L_{h,\tau}\}$  is said to satisfy the *von Neumann condition for stability* if there exists a positive rational number  $a$ , and positive constants  $h_0$  and  $\lambda_0$  such that for all  $\alpha \in E^n$  and  $h \leq h_0$ , and for every subfamily  $\{L_{h,\lambda}\}$  of  $\{L_{h,\tau}\}$  with  $\lambda \equiv \tau h^{-a} = \text{constant} \leq \lambda_0$  the eigenvalues  $\gamma_j = \gamma_j(\lambda, h, a, \alpha)$  of  $-H_1^{-1}H_0$  all satisfy

$$(1.3) \quad |\gamma_j| \leq 1 + c\tau,$$

where  $c$  is a positive constant which may depend on  $a, h_0$  and  $\lambda$ . The von Neumann condition is necessary for the stability of solutions of the initial value problem for  $\{L_{h,\tau}\}$  (see [7] and §3 below).

In §2 we shall prove

**THEOREM 1.** *The differential operator  $L$  is correct in the sense of Petrovskiĭ if and only if there exists at least one family of two-level finite difference operators  $\{L_{h,\tau}\}$  which is consistent with  $L$  and which satisfies the von Neumann condition for stability.*

In view of the connection between correctness in the sense of Petrovskiĭ and the von Neumann condition for stability it is reasonable to expect to be able to prove finite difference analogues of known results concerning the initial value problem with correct  $L$  for the initial value problem for a family  $\{L_{h,\tau}\}$  which satisfies the von Neumann condition. We consider this question in §3, where we show that for explicit two-level finite difference operators the von Neumann condition is sufficient for stability in the sense of Rjaben'kiĭ and Filippov [8].

**2. Proof of Theorem 1.** Assume that  $L$  is correct in the sense of Petrovskiĭ, i.e., that (1.1) holds. Let  $\delta = (\delta_1, \dots, \delta_n)$ , where  $\delta_j u(x, t) = u(x + e_j h, t) - u(x - e_j h, t)$  and  $e_j$  is the  $j$ th unit coordinate vector in  $E^n$ , and let  $\Delta u(x, t) = u(x, t + \tau) - u(x, t)$ . The family of two-level finite difference operators

$$(2.1) \quad L_{h,\tau} \equiv P(\delta/2h) - \tau^{-1}E\Delta$$

is clearly consistent with  $L$ . We claim that  $\{L_{h,\tau}\}$  satisfies the von

Neumann condition. In this case  $H_1 = -\tau^{-1}E$  and  $H_0 = \tau^{-1}E + P(i \sin \alpha h/h)$ , where  $\sin \alpha h = (\sin \alpha_1 h, \dots, \sin \alpha_n h)$ . Thus  $-H_1^{-1}H_0 = E + \tau P(i \sin \alpha h/h)$  and it follows that

$$(2.2) \quad |\gamma_j|^2 = 1 + \tau \{ 2 \operatorname{Re} \mu_j(\sin \alpha h/h) + \tau |\mu_j(\sin \alpha h/h)|^2 \}.$$

By a theorem of Gel'fand and Šilov [3, p. 83], there exists a constant  $c_1 > 0$  such that

$$(2.3) \quad \max_j |\mu_j(\xi)| \leq c_1(1 + |\xi|)^{m_0},$$

where  $|\xi|^2 = \sum_{k=1}^n \xi_k^2$ ,  $m_0 = \max_{1 \leq k \leq N} (p_k/k)$ , and  $p_k$  is the degree of the coefficient of  $\mu^{N-k}$  in the characteristic polynomial of  $P(i\xi)$ . If we use (1.1), (2.3) and the fact that  $|\sin \alpha h| \leq n^{1/2}$  in (2.2) we obtain

$$|\gamma_j|^2 \leq 1 + \tau \{ 2K + c_1^2 \tau^2 h^{-2m_0} (h + n^{1/2})^{2m_0} \}.$$

Thus if we set  $\lambda = \tau h^{-2m_0}$  and  $h_0 = 1$ , the von Neumann condition holds for arbitrary  $\lambda_0 > 0$ .

Now assume that we have a family of finite difference operators  $\{L_{h,\tau}\}$  which is consistent with  $L$  and for which the von Neumann condition holds. Choose any  $\lambda \leq \lambda_0$  and consider the subfamily  $\{L_{h,\lambda}\}$ . Write  $-H_1^{-1}H_0$  in the form

$$(2.4) \quad -H_1^{-1}H_0 = E + \tau P_{h,\lambda}(\alpha).$$

We assert that for each fixed  $\alpha \in E^n$

$$(2.5) \quad \lim_{h \rightarrow 0} P_{h,\lambda}(\alpha) = P(i\alpha).$$

Let  $u_1(x) = e^{i\alpha^T x} E$  and  $u_2(x, t) = t u_1(x)$ . If we apply (1.2) first to  $u_1$  and then to  $u_2$  we obtain  $-(H_0 + H_1)u_1 = P(i\alpha)u_1 + A_1$  and  $\tau H_1 u_2 = E + A_2$ , where  $A_j \rightarrow 0$  as  $h \rightarrow 0$  for each fixed  $\alpha \in E^n$ . In particular, it follows that  $(\tau H_1)^{-1} = E + A_3$ , where  $A_3 \rightarrow 0$  as  $h \rightarrow 0$  for fixed  $\alpha$ . Thus  $-\tau^{-1}(E + H_1^{-1}H_0)u_1 = P(i\alpha)u_1 + A_3 P(i\alpha)u_1 + A_1 + A_3 A_1$ , which proves the assertion.

Let  $\mu_j^{(h,\lambda)}(\alpha)$  denote the eigenvalues of  $P_{h,\lambda}(\alpha)$ . In view of (2.4) we have  $\gamma_j = 1 + \tau \mu_j^{(h,\lambda)}$ . Since the von Neumann condition holds, it follows from (1.3) that

$$|\gamma_j|^2 = 1 + \tau(2 \operatorname{Re} \mu_j^{(h,\lambda)} + \tau |\mu_j^{(h,\lambda)}|^2) \leq 1 + \tau c(2 + h_0^a \lambda c)$$

for all  $h \leq h_0$  and  $\alpha \in E^n$ , where  $c = c(a, h_0, \lambda)$ . In particular,

$$(2.6) \quad \operatorname{Re} \mu_j^{(h,\lambda)}(\alpha) \leq c \left( 1 + \frac{1}{2} h_0^a \lambda c \right) \equiv \tilde{c}(a, h_0, \lambda)$$

for all  $h \leq h_0$  and  $\alpha \in E^n$ . On the other hand, (2.5) implies that  $\mu_j^{(h,\lambda)}(\alpha) \rightarrow \mu_j(\alpha)$  as  $h \rightarrow 0$  for each fixed  $\alpha \in E^n$ . Thus it follows from (2.6) that  $\operatorname{Re} \mu_j(\alpha) \leq \zeta(a, h_0, \lambda)$  for all  $\alpha \in E^n$ , i.e.,  $L$  is correct in the sense of Petrovskii.

REMARK. Although, as we have proved,  $\tau h^{-2m_0} = \text{constant}$  implies the von Neumann condition for the difference operators (2.1), in certain special cases an exponent smaller than  $2m_0$  will suffice. For example, if  $L$  is parabolic in the sense of Petrovskii [3], i.e., if  $\operatorname{Re} \mu_j(\xi) \leq -\rho |\xi|^m$  for some  $\rho > 0$ , then  $m_0 = m$  but the von Neumann condition holds for (2.1) if  $\tau h^{-m}$  is suitably restricted [2]. On the other hand, if for  $n = 1$

$$P(i\xi) = \begin{pmatrix} 0 & i\xi \\ i\xi & 0 \end{pmatrix}$$

(i.e.,  $Lu = 0$  is the one-dimensional wave equation) then  $m_0 = 1$  and  $\tau h^{-2} = \text{constant}$  is also necessary for the von Neumann condition. In this case, and in general, the situation can be improved by using a more sophisticated difference approximation in place of (2.1) (see [7, p. 168]).

**3. Stability.** We now restrict our attention to the class of *explicit* finite difference operators

$$\mathfrak{L}_{h,\tau} u(x, t) \equiv \sum_{\tau} B_{\tau} u(x + \tau h, t) - \tau^{-1} \Delta u(x, t).$$

For example, the difference operators (2.1) are clearly explicit. Moreover, the two-level approximation to the one-dimensional wave operator discussed in [7, formula (8.5), p. 168] is also explicit since it can be written in the form

$$(3.1) \quad \mathfrak{L}_{h,\tau} = \begin{pmatrix} 0 & \frac{c}{2h} \delta \\ \frac{c}{2h} \delta & \frac{c^2 \tau}{4h^2} \delta^2 \end{pmatrix} - \frac{1}{\tau} \begin{pmatrix} \Delta & 0 \\ 0 & \Delta \end{pmatrix}.$$

We shall assume that  $\{\mathfrak{L}_{h,\tau}\}$  is consistent with a fixed differential operator  $L \equiv P(D) - E\partial/\partial t$  and that the von Neumann condition holds for  $\{\mathfrak{L}_{h,\tau}\}$ . (Thus  $L$  is correct.) Let  $g = g(x)$  be a given  $N$ -vector defined on  $E^n$ . We consider the two-parameter family of initial value problems

$$(3.2) \quad \begin{aligned} \mathfrak{L}_{h,\tau} u(x, t) &= 0 && \text{for } (x, t) \in R_{h,\tau}(T); \\ u(jh, 0) &= g(jh) && \text{for } j \in I^n, \end{aligned}$$

where

$$R_{h,\tau}(T) = \{(jh, l\tau) \mid j \in I^n, l = 0, 1, \dots, T/\tau - 1\}$$

for arbitrary  $T > 0$  and  $\tau$  chosen so that  $T/\tau = \text{integer}$ . Since  $\mathcal{L}_{h,\tau}$  is explicit and  $\tau$  ranges over a finite subset of  $I^n$ , it is clear that the initial value problem has a unique solution for every positive  $h$  and  $\tau$ .

Let  $\lambda$  be a positive constant and consider the one-parameter subfamily  $\{\mathcal{L}_{h,\lambda}\} \subset \{\mathcal{L}_{h,\tau}\}$  characterized by  $\tau h^{-a} = \lambda$  for some fixed positive rational number  $a$ . The one-parameter family of initial value problems (3.2) corresponding to  $\{\mathcal{L}_{h,\lambda}\}$  is said to be *stable* if it is uniformly well-posed for all sufficiently small  $h$ , i.e., if there exist norms  $\|\cdot\|_0$  and  $\|\cdot\|_1$ , and positive constants  $h_0$  and  $\sigma = \sigma(a, h_0, \lambda)$ , such that for every  $h \leq h_0$  the solution  $u = u_{h,\lambda}(x, t)$  satisfies  $\|u\|_1 \leq \sigma \|g\|_0$ . The problem of stability of (3.2) has been studied by various authors. In the work of Lax and Richtmyer [7] and Kreiss [5; 6],  $\|g\|_0$  is the  $L_2(E^n)$ -norm and  $\|u\|_1 = \sup_{[0, T]} (\int_{E^n} |u|^2 dx)^{1/2}$ . Kreiss [5] gives a necessary and sufficient condition for stability in this sense. John [4] and Aronson [1; 2] use the uniform norm for  $\|g\|_0$  and a norm involving the first  $m - 1$  differences of  $u$  for  $\|u\|_1$ . In all of these works the von Neumann condition is necessary but, in general, not sufficient for stability. In this paper, as in the work of Rjaben'kiĭ and Filippov [8],  $\|u\|_1$  will be the uniform norm and  $\|g\|_0$  will involve various derivatives (or differences) of  $g$ . We shall show that the von Neumann condition is sufficient for stability in this sense.

For explicit difference operator  $\mathcal{L}_{h,\tau}$  we have  $-H_1^{-1}H_0 = E + \tau P_{h,\tau}(\alpha)$ , where

$$(3.3) \quad P_{h,\tau}(\alpha) = \sum_r B_r e^{i\alpha^T r h}.$$

Since  $\{\mathcal{L}_{h,\tau}\}$  is consistent with  $L$ , we have

$$(3.4) \quad P_{h,\tau}(\alpha) \rightarrow P(i\alpha) \quad \text{as } h, \tau \rightarrow 0$$

for each fixed  $\alpha \in E^n$ . Write  $P(\xi) = \sum_{l=0}^m P_l(\xi)$ , where the elements of  $P_l(\xi)$  are homogeneous polynomials of degree  $l$  in  $\xi$ . It follows from (3.3) and (3.4) that

$$(3.5) \quad \frac{(ih)^l}{l!} \sum_r B_r (\alpha^T r)^l \rightarrow \begin{cases} P_l(i\alpha) & \text{for } l = 0, 1, \dots, m, \\ 0 & \text{for } l = m + 1, m + 2, \dots \end{cases}$$

as  $h, \tau \rightarrow 0$  for fixed  $\alpha \in E^n$ . The matrices  $B_r = 0$  for  $r$  outside a finite neighborhood of  $r = 0$ . Let  $r^{(1)}, \dots, r^{(s)}$  be the set of all  $r \in I^n$  such that  $r \neq 0$  and  $B_r \neq 0$ . Using Cramer's rule, it is not difficult to verify

from (3.5) that  $h^s B_r$  is bounded as  $h, \tau \rightarrow 0$  for all  $r$ . Let  $q$  be the smallest integer for which  $h^q B_r$  is bounded as  $h, \tau \rightarrow 0$  for all  $r$ . Clearly  $q \leq s$ . It is possible to have  $q < s$ . For the operator (2.1) we have  $s = 2m$  and  $q = m$ , and for the operator (3.1) we have  $s = 4$  and  $q = 2$ . In general, if  $\mathfrak{L}_{h,\tau}$  involves only centered  $x$ -differences  $\delta$ , then  $q = s/2$ . It follows from Taylor's theorem and (3.3) that<sup>2</sup>

$$|P_{h,\tau}(\alpha)| \leq \sum_{l=0}^{q-1} |\alpha|^l \left| \frac{(ih)^l}{l!} \sum_r B^r \left( \frac{\alpha^T r}{|\alpha|} \right)^l \right| + \frac{|\alpha|^q}{q!} \sum_r h^q |B_r| |\tau|^q.$$

Thus in view of (3.5) and the definition of  $q$ , there exist positive constants  $h_0, \tau_0$ , and  $c_1 = c_1(h_0, \tau_0, q, s)$  such that

$$(3.6) \quad |P_{h,\tau}(\alpha)| \leq c_1(1 + |\alpha|)^q \quad \text{for all } h \leq h_0, \tau \leq \tau_0 \text{ and } \alpha \in E^n.$$

Let  $\mathcal{Q}_\nu$ , where  $\nu$  is a non-negative integer, denote the class of all  $N$ -vectors  $g = g(x)$  defined on  $E^n$  which are such that  $g \in C^r(E^n)$  and  $D^k g \in L(E^n)$  for  $|k| \leq \nu$ . If  $g \in \mathcal{Q}_\nu$  we shall write

$$\|g\|_\nu = \max_{|k| \leq \nu} \left( \int_{E^n} |D^k g| dx \right).$$

Let

$$\bar{R}_{h,\tau}(T) = \{(jh, l\tau) \mid j \in I^n, l = 0, 1, \dots, T/\tau\}.$$

For  $u = u(x, t)$  defined on  $\bar{R}_{h,\tau}(T)$  we shall write

$$\|u\| = \max_{\bar{R}_{h,\tau}(T)} |u(x, t)|.$$

We shall prove

**THEOREM 2.** *Let the family  $\{\mathfrak{L}_{h,\tau}\}$  of explicit finite difference operators be consistent with a fixed differential operator  $L$  and satisfy the von Neumann condition. For any  $\lambda \leq \lambda_0$  let  $u = u_{h,\lambda}(x, t)$  denote the solution of (3.2) for  $\mathfrak{L}_{h,\lambda}$ . Then if  $g \in \mathcal{Q}_\nu$ , for  $\nu = q(N-1) + n + 1$ , there exists a constant  $\sigma = \sigma(a, h_0, \lambda, q, s, N, T)$  such that*

$$\|u\| \leq \sigma \|g\|_\nu \quad \text{for all } h \leq h_0.$$

**PROOF.** Let  $\hat{g}(\alpha)$  denote the Fourier transform of  $g(x)$ . Since  $g \in \mathcal{Q}_\nu$ , it is easy to verify that  $|\hat{g}(\alpha)| \leq (n+1)^\nu (1 + |\alpha|)^{-\nu} \|g\|_\nu$ . Assume that there is a constant  $c_2 > 0$  such that

$$(3.7) \quad \left| \{E + \tau P_{h,\lambda}(\alpha)\}^l \right| \leq c_2(1 + |\alpha|)^{q(N-1)}$$

<sup>2</sup> If  $u = (u_1, \dots, u_N)$  then  $|u| = \sum_{j=1}^N |u_j|$ . If  $A = (a_{ij})$  is an  $N \times N$  matrix then  $|A| = N \max_{i,j} |a_{ij}|$ . Note that  $|Au| \leq |A| |u|$ .

for all  $h \leq h_0$ ,  $\alpha \in E^n$  and  $l\tau \leq T$ . Let

$$u(x, l\tau) = (2\pi)^{-n/2} \int_{E^n} \{E + \tau P_{h,\lambda}(\alpha)\} {}^l g(\alpha) e^{i\alpha^T x} d\alpha.$$

Then

$$|u(x, l\tau)| \leq (2\pi)^{-n/2} (n+1)^{r_2} \|g\|_r \int_{E^n} (1 + |\alpha|)^{-(n+1)} d\alpha,$$

and  $u(jh, l\tau)$  is clearly the solution of (3.2). Thus we must show that (3.7) holds.

By a theorem of Schur (quoted in [7, p. 67]) there exists a unitary matrix  $U = U_{h,\lambda}(\alpha)$  such that  $U^* P_{h,\lambda}(\alpha) U = \tilde{P}_{h,\lambda}(\alpha) = (\tilde{p}_{ij}^{(h,\lambda)}(\alpha))$ , where  $\tilde{p}_{ii}^{(h,\lambda)} = \mu_i^{(h,\lambda)}$ ,  $\tilde{p}_{ij}^{(h,\lambda)} = 0$  for  $i > j$ , and  $|\tilde{p}_{ij}^{(h,\lambda)}| \leq N^2 \max_{k,l} |\mu_{kl}^{(h,\lambda)}|$ . Thus  $\{E + \tau P_{h,\lambda}(\alpha)\} {}^l U = U \{E + \tau \tilde{P}_{h,\lambda}(\alpha)\} {}^l U^*$ . Write  $E + \tau \tilde{P}_{h,\lambda} = \Gamma + \tau \bar{P}_{h,\lambda}$ , where  $\Gamma$  is the diagonal matrix with elements  $\gamma_j = 1 + \tau \mu_j^{(h,\lambda)}$ . In view of (3.6),  $|\bar{P}_{h,\lambda}| \leq c_1 N^3 (1 + |\alpha|)^q$ . Moreover, according to (1.3),  $|\gamma_j| \leq 1 + c\tau$ . Thus for  $l > N - 1$ , taking account of the structure of  $\Gamma$  and  $\bar{P}_{h,\lambda}$ , we find

$$|(E + \tau \tilde{P}_{h,\lambda})^l| \leq \sum_{j=0}^{N-1} (N\tau)^j C_{l,j} (1 + c\tau)^{l-j} \{c_1 N^3 (1 + |\alpha|)^q\}^j \equiv M.$$

Since  $C_{l,j} \leq l^j/j!$  and  $l\tau \leq T$  we obtain

$$M \leq N^{N-1} (1 + |\alpha|)^{q(N-1)} e^{cT} \sum_{j=0}^{N-1} (c_1 T N^3)^j / j! \equiv c_3 (1 + |\alpha|)^{q(N-1)},$$

where  $c_3 = c_3(a, h_0, \lambda, q, s, N, T)$ . Therefore  $|(E + \tau P_{h,\lambda})^l| \leq c_3 N^2 (1 + |\alpha|)^{q(N-1)}$  for  $l \geq N$ . On the other hand, it follows from (3.6) that for  $l \leq N - 1$ ,  $|(E + \tau P_{h,\lambda})^l| \leq (N + h_0^q c_1 \lambda)^{N-1} (1 + |\alpha|)^{q(N-1)} \equiv c_4 (1 + |\alpha|)^{q(N-1)}$ , where  $c_4 = c_4(a, h_0, \lambda, q, s, N)$ . Therefore (3.7) holds with  $c_2 = \max(c_3 N^2, c_4)$  and the proof of the theorem is complete.

REMARKS. As Rjaben'kiĭ and Filippov [8] point out, (3.7) is the finite difference analogue of Petrovskii's "Bedingung A." Thus we may paraphrase Theorem 2 as follows: The von Neumann condition implies "Bedingung A" which in turn implies stability in the indicated sense. If, in addition to the correctness implied by the hypothesis of Theorem 2, we make further assumptions about the structure of  $L$  we can, of course, obtain much stronger results (see, e.g., [1; 2] for results concerning parabolic equations and systems). On the other hand, Theorem 2 can be generalized for data of polynomial growth. This, however, would require the deep methods of Gel'fand and Šilov [3] and is beyond the scope of this note.

Stability considerations are important for two reasons. First because, in general, they yield information about the effects of rounding errors on the numerical solution of (3.2), and second because in the appropriate setting, as was shown by Lax [7], stability is necessary and sufficient for the convergence of solutions of (3.2) to the solution of the corresponding initial value problem for  $L$ . Theorem 2 is not appropriate for the former application. To obtain a better result for this purpose, we assume that  $g$  has compact support and let

$$\|g\|_{h,\nu} = \max_{x \in E^n, |k| \leq \nu} |h^{-|k|} \Delta^k g(x)|,$$

where  $\Delta = (\Delta_1, \dots, \Delta_n)$  and  $\Delta_j g(x) = g(x + e_j h) - g(x)$ . For such  $g$ , Rjaben'kiĭ and Filippov [8] have shown that (3.7) implies that  $\|u\| \leq \sigma \|g\|_{h,\nu}$ , where  $\sigma$  depends on the support of  $g$ . Since we have shown that the von Neumann condition implies (3.7) it follows that under the hypothesis on  $L$  in Theorem 2,  $\|u\| \leq \sigma \|g\|_{h,\nu}$ . Thus, in particular, if the values of  $g$  are known only to within  $\pm \epsilon$ , then the error from this source in  $u$  is bounded by  $2^{\nu} \sigma \epsilon h^{-\nu}$ .

#### REFERENCES

1. D. G. Aronson, *The stability of finite difference approximations to second order linear parabolic differential equations*, Duke Math. J. **30** (1963), 117-128.
2. ———, *On the stability of certain finite difference approximations to parabolic systems of differential equations*, Numer. Math. **5** (1963), 118-137, 290.
3. I. M. Gel'fand and G. E. Šilov, *Generalized functions*, Part 3, Gosudarstv. Izdat. Fiz.-Mat. Lit., Moscow, 1958. (Russian)
4. Fritz John, *On integration of parabolic equations by difference methods*, Comm. Pure Appl. Math. **5** (1952), 155-211.
5. Heinz-Otto Kreiss, *Über die Approximative Lösung von linearen partiellen Differentialgleichungen mit Hilfe von Differenzgleichungen*, Kungl. Tekn. Högsk. Handl. Stockholm **128** (1958), 1-19.
6. ———, *Über die Lösung des Cauchyproblems für lineare partielle Differentialgleichungen mit Hilfe von Differenzgleichungen*. 1, Acta Math. **101** (1959), 179-199.
7. R. D. Richtmyer, *Difference methods for initial-value problems*, Interscience Tracts in Pure and Applied Mathematics. Tract 4. Interscience, New York, 1957.
8. V. S. Rjaben'kiĭ and A. F. Filippov, *On stability of difference equations*, GITTL, Moscow, 1956 (Russian); German transl., *Über die Stabilität von Differenzgleichungen*, Berlin, 1960.

STANFORD UNIVERSITY AND  
UNIVERSITY OF MINNESOTA