

THE EXISTENCE OF PROPER SOLUTIONS OF A SECOND ORDER ORDINARY DIFFERENTIAL EQUATION

J. D. SCHUUR

We shall consider the equation

$$(1) \quad \begin{aligned} x'' &= f(t, x, x'): f(t, x, y) \text{ continuous on } D \\ &= \{ (t, x, y): 0 \leq t < +\infty, -\infty < x, y < +\infty \}. \end{aligned}$$

We further assume: (i) $f(t, x, y)$ is such that the solutions of (1) are uniquely determined by initial conditions; (ii) $xf(t, x, 0) > 0$ if $t > 0, x \neq 0$; (iii) $f(t, 0, 0) = 0$ for all $t \geq 0$; (iv) if $u > v$, then $f(t, u, y) \geq f(t, v, y)$, if $r > s$, then $f(t, x, r) \geq f(t, x, s)$.

From (ii), a positive solution of (1) has no relative maxima, a negative solution has no relative minima. If $x(0) > 0$ and $x(c) = 0, x'(c) < 0$ for some $c > 0$, then $x(t) < 0$ for all $t > c$. The behavior of positive solutions and negative solutions is similar, so we shall consider only the former. If we totally disregard nonpositive solutions, we can weaken (ii) to $f(t, x, 0) > 0$ for $x, t > 0$.

A solution, $x(t)$, of (1) is *proper* if $x(t)$ exists and is positive for all $t \geq 0$. If $f(t, x, y)$ satisfies (i)–(iv), we shall show that given $A > 0$ there exists a unique proper solution of (1), $x(t)$, such that $x(0) = A$ and $x'(t) < 0$ for all $t \geq 0$. We shall use the topological method of T. Ważewski ([2] or [1, pp. 179–182]). This approach generalizes a result (and simplifies the proof) of P. K. Wong [3, Theorem 1.1].

Let $u(t)$ and $v(t)$ be two solutions of (1). If $u(t)$ and $v(t)$ are defined for $0 \leq t < a \leq +\infty$, then

$$(2) \quad \begin{aligned} u'(t) - v'(t) &= u'(0) - v'(0) \\ &+ \int_0^t [f(r, u(r), u'(r)) - f(r, v(r), v'(r))] dr. \end{aligned}$$

If $u(t)$ and $v(t)$ are defined for $0 \leq t \leq b < +\infty$, then

$$(3) \quad \begin{aligned} u(t) - v(t) &= u(0) - v(0) + [u'(0) - v'(0)]t \\ &+ \int_0^t \int_0^s [f(r, u(r), u'(r)) - f(r, v(r), v'(r))] dr ds. \end{aligned}$$

THEOREM 1. *If $u(t)$ and $v(t)$ are two proper solutions of (1) with $u(0) = v(0) = A > 0, u'(\infty) = v'(\infty) = 0$, then $u(t) = v(t)$ for all $t \geq 0$.*

Received by the editors November 1, 1965.

PROOF. Let $u(t)$ and $v(t)$ satisfy the hypotheses and assume $v'(0) < u'(0)$. Suppose there exists a $c > 0$ such that $v(t) < u(t)$ and $v'(t) < u'(t)$ for $0 < t < c$ and either $v(c) = u(c)$ or $v'(c) = u'(c)$. From (2) and (iv), $u'(c) - v'(c) \geq u'(0) - v'(0) > 0$; and from (3) and (iv), $u(c) - v(c) \geq [u'(0) - v'(0)]c > 0$. Therefore, $v(t) < u(t)$ and $v'(t) < u'(t)$ for $0 \leq t < +\infty$ and $0 = u'(\infty) - v'(\infty) \geq [u'(0) - v'(0)] > 0$, a contradiction. Thus, $v'(0) \geq u'(0)$ —and likewise $u'(0) \geq v'(0)$ —so $u(t) = v(t)$ for all $t \geq 0$.

LEMMA. Given $A > 0, c > 0$ there exists a $d(A, c) > 0$ such that if $x(t)$ is a solution of (1) with $0 < x(0) \leq A$ and $x'(0) \leq -d$, then $x(t) = 0$ for some $t \in (0, c)$.

PROOF. Let $T = \{(t, x) : 0 \leq t \leq c, 0 \leq x \leq A - t(A/c)\}$ and let H be the hypotenuse of T . We will show that solutions satisfying the hypotheses do not cross H for $0 \leq t \leq c$.

Let $M = \max\{f(t, x, 0) : (t, x) \in T\}$, $M > 0$ by (ii), and let $d = (A/c) + Mc$. Let $x(t)$ be a solution of (1) with $0 < x(0) \leq A$ and $x'(0) \leq -d$. For t small, $x(t)$ lies in T and below H . If $x(t)$ strikes H for some $t \in (0, c]$, then there exists an $e \in (0, c)$ such that $x'(e) = -(A/c)$, and $x'(t) < -(A/c)$, $(t, x(t)) \in T$ for $0 \leq t < e$. Then

$$\begin{aligned} x'(e) &= x'(0) + \int_0^e f(s, x(s), x'(s)) ds \\ &\leq -(A/c) - Mc + Me < -(A/c). \end{aligned}$$

THEOREM 2. Given $A > 0$ there exists exactly one proper solution, $x(t)$, of (1) such that $x(0) = A$ and $x'(t) < 0$ for all $t \geq 0$.

PROOF. Theorem 1 shows the uniqueness of such solutions.

Write (1) as a system

$$(4) \quad \begin{aligned} x' &= y, \\ y' &= f(t, x, y); \end{aligned}$$

we seek a solution, $(x(t), y(t))$, of (4) such that $x(0) = A > 0, y(0) < 0$, and $x(t) > 0, y(t) < 0$ for all $t \geq 0$. We will now use the method and terminology of Ważewski. Let $T = \{(t, x, y) : t \geq 0, x > 0, y < 0\}$; $Q = \{(t, x, y) : t \geq 0, x > 0, y = 0\}$; and $R = \{(t, x, y) : t \geq 0, x = 0, y < 0\}$. For solutions of (4) with $t > 0$, the set of egress of T is $S = Q \cup R$ (by (i), the t -axis contains no points of egress). Every point in Q is a point of strict egress since $y' = f(t, x, 0) > 0$ and every point in R is a point of strict egress since $x' = y < 0$ on R . Thus, the set of egress equals the set of strict egress.

Now let $X = \{(t, x, y) : t=0, x=A, -d \leq y \leq 0\}$ (where d is determined in the lemma); $Y = \{(t, x, y) : t=0, 0 \leq x \leq A, y=-d\}$; and $Z = X \cup Y$. Then $S \cap Z$ is a retract of S but not of Z and by Ważewski's theorem there exists a point $P = (0, a, b) \in Z$ such that the solution of (4) with $x(0) = a, y(0) = b$ remains in T for all $t \geq 0$. And by the lemma, $P \notin Y$.

REFERENCES

1. L. Cesari, *Asymptotic behavior and stability problems in ordinary differential equations*, Springer-Verlag, Berlin, 1959.
2. T. Ważewski, *Une méthode topologique de l'examen du phénomène asymptotique relativement aux équations différentielles ordinaires*, Rend. Accad. Lincei 3 (1947), 210-215.
3. P. K. Wong, *Existence and asymptotic behavior of proper solutions of a class of second-order nonlinear differential equations*, Pacific J. Math. 13 (1963), 737-760.

MICHIGAN STATE UNIVERSITY