

# ČEBYŠEV SUBSPACES OF FINITE DIMENSION IN $L_1$

R. R. PHELPS<sup>1</sup>

**1. Introduction.** If  $E$  is a real normed linear space, a subspace  $M$  of  $E$  is said to be a Čebyšev subspace if to each  $y$  in  $E$  there exists a unique nearest element  $x$  in  $M$ , that is, an element  $x$  in  $M$  such that  $\|y-x\| < \|y-z\|$  for  $z \in M, z \neq x$ . The problem of characterizing such subspaces for the classical Banach spaces of functions is an interesting one. The first (and best) result of this nature was the classical theorem of Haar [6] characterizing the finite dimensional Čebyšev subspaces of  $C[0, 1]$ . Much recent work has been done on this and related questions by A. L. Garkavi [3], [4], [5], [6] and the author [9], [10], stimulated in part by the discovery [9] of a certain duality between uniqueness of best approximation and uniqueness of Hahn-Banach extensions of functionals. So far, however, there has been no characterization given of the finite dimensional Čebyšev subspaces of  $L_1(T, \Sigma, \mu)$  ( $T$   $\sigma$ -finite). This is partly due to the fact that if  $(T, \Sigma, \mu)$  is nonatomic, then  $L_1(T, \Sigma, \mu)$  contains no finite dimensional Čebyšev subspaces, a result due to M. S. Krein and B. Ya. Levin [1, Chapter IV] for  $T = [0, 1]$ ,  $\mu$  Lebesgue measure, and to Henry Dye [9] in the general case. In fact, Garkavi [4] has extended these arguments to show that  $L_1(T, \Sigma, \mu)$  contains a Čebyšev subspace of dimension  $n < \infty$  if and only if  $(T, \Sigma, \mu)$  contains at least  $n$  atoms. This result, of course, does not tell which subspaces are Čebyšev subspaces, a question which we consider in the present paper. Our characterization is not entirely satisfactory, in that it is given in terms of the elements of the annihilator  $M^\perp$  of  $M$  in  $L_\infty$ , rather than in terms of the elements of  $M$  itself, but we see no way to avoid this. ( $M^\perp = \{f: f \in L_\infty(T, \Sigma, \mu) \text{ and } \int_T fg d\mu = 0 \text{ for each } g \in M\}$ .) The proof makes essential use of an interpolation theorem (Lemma 2) which is actually a reformulation of Liapunov's theorem on the range of a vector measure.

The theorem itself has two corollaries of interest; one immediate, the other not so immediate. The first one is simply the restatement of the theorem for the case when  $L_1(T, \Sigma, \mu)$  is actually the space  $l_1$ .

**COROLLARY 1.** *Suppose that  $M$  is an  $n$ -dimensional subspace of  $l_1$ . Then  $M$  is a Čebyšev subspace if and only if for each  $z = \{z_k\}$  in  $M^\perp \subset l_\infty, z \neq 0$ , there are at least  $n$  integers  $k$  such that  $|z_k| < \|z\|_\infty$ .*

Received by the editors November 20, 1964.

<sup>1</sup> Supported by Grant GP 378 of the National Science Foundation, U.S.A.

The second corollary characterizes the subspaces  $M$  of finite co-dimension in  $E = C(T)$  ( $T$  compact Hausdorff) which have *property U*: Every continuous linear functional on  $M$  has a unique norm-preserving extension to  $E$ . The relation between this property and the Čebyšev property arises from the fact [9] that a subspace  $M$  of a normed linear space  $E$  has property *U* if and only if  $M^\perp$  is a Čebyšev subspace of  $E^*$ . Again, our characterization is not as “intrinsic” as we would like, but an example shows that the only obvious improvement is impossible.

If  $f \in L_\infty(T, \Sigma, \mu)$ , denote by  $S(f)$  the measurable set (defined to within a set of measure zero) on which  $|f| < \|f\|$ . Our main theorem is the following:

**THEOREM 1.** *Suppose that  $M$  is an  $n$ -dimensional subspace of  $L_1(T, \Sigma, \mu)$ . Then  $M$  fails to be a Čebyšev subspace if and only if there exists  $f$  in  $M^\perp \subset L_\infty(T, \Sigma, \mu)$ ,  $f \neq 0$ , such that  $S(f)$  is purely atomic and contains at most  $n - 1$  atoms.*

The proof of Theorem 1 depends on several lemmas, the first of which actually characterizes Čebyšev subspaces of arbitrary dimension in  $L_1$ . The lemma itself is new, but the ideas in the proof have all been used previously.

For a measurable function  $g$ , let  $Z(g) = \{t: g(t) = 0\}$ ; for elements of  $L_1$  this is defined only to within a set of measure zero, and set theoretic operations with both  $Z(g)$  and  $S(f)$  are to be interpreted to be modulo sets of measure zero.

**LEMMA 1.** *A closed linear subspace  $M$  of  $L_1$  is not a Čebyšev subspace if and only if there exist  $f$  in  $M^\perp$ ,  $\|f\| = 1$ , and  $g$  in  $M$ ,  $g \neq 0$ , such that  $S(f) \subset Z(g)$ .*

**PROOF.** If  $M$  is not a Čebyšev subspace there exist  $h$  in  $L_1 \sim M$  and  $g$  in  $M$ ,  $g \neq 0$ , such that  $1 = \|h\| = \|h + g\| = d(h, M)$ . By the Hahn-Banach theorem there exists  $f$  in  $M^\perp \subset L_\infty$  such that  $\|f\| = 1 = \int f h d\mu = \int f(h + g) d\mu$ . It follows that  $S(f) \subset Z(h) \cap Z(h + g) \subset Z(g)$ .

To prove the converse, suppose that  $f$  and  $g$  exist as above, and define  $h = f|g|$ . We will show that  $\int f h d\mu = \|h\|$  and that  $\|h\| = \|h - g\|$ . (Since, for any  $g'$  in  $M$ ,  $\|h - g'\| \geq \int f(h - g') d\mu = \int f h d\mu = \|h\|$ , this will show that  $h$  has 0 and  $g$  as distinct nearest points in  $M$ .) First,

$$\int f h d\mu = \int f^2 |g| d\mu = \int |g| d\mu = \int |h| d\mu = \|h\|.$$

Second, we note that

$$|h - g| = |(f|g| - g)| = f(f|g| - g) = |g| - fg,$$

so that

$$\|h - g\| = \int |h - g| d\mu = \int (|g| - fg)d\mu = \int |g| d\mu = \|h\|,$$

and the proof is complete.

We can now easily prove half of Theorem 1. Suppose that  $M$  is an  $n$ -dimensional subspace of  $L_1$  and that there exists  $f \neq 0$  in  $M^\perp \subset L_\infty$  such that  $S(f)$  is the union of  $k$  atoms,  $0 \leq k \leq n - 1$ . Let  $N = \{g: g \in L_1 \text{ and } g = 0 \text{ a.e. on } S(f)\}$ . Evidently  $N$  has finite codimension  $k < n$ , so that there exists  $g \neq 0$  in  $M \cap N$ , and hence  $M$  is not a Čebyšev subspace, by Lemma 1.

The proof of the converse is more involved. If  $M$  is an  $n$ -dimensional subspace of  $L_1$  which is not a Čebyšev subspace, then there exist  $f$  in  $M^\perp$ ,  $\|f\| = 1$ , and  $g$  in  $M$ ,  $g \neq 0$ , such that  $S(f) \subset Z(g)$ . We will first replace  $f$  by a function  $f'$  in  $M^\perp$ ,  $\|f'\| = 1$ , for which  $S(f') \subset S(f)$  and  $S(f')$  is purely atomic; after that, a further replacement will yield  $f''$  in  $M^\perp$ ,  $\|f''\| = 1$ , such that  $S(f') \supset S(f'')$  and the latter has at most  $n - 1$  atoms. To this end, write  $S(f) = S \cup A$ , where  $S$  contains no atoms, and  $A$  is purely atomic. (Since  $T$  is  $\sigma$ -finite,  $A$  is the union of at most countably many atoms.) The function  $f'$  will be defined to be equal to  $f$  on  $T \sim S$  and to have absolute value 1 on  $S$ , so that  $S(f') \subset A$ . To this end, choose  $g_2, \dots, g_n$  such that  $g_1 = g, g_2, \dots, g_n$  form a basis for  $M$ . If we can produce  $f'$  with  $|f'| = 1$  a.e. on  $S$ ,  $f' = f$  on  $T \sim S$  and  $\int_S f' g_i d\mu = \int_S f g_i d\mu, i = 2, \dots, n$ , then we will certainly have  $\int_T f' g_i d\mu = \int_T f g_i d\mu = 0$  for each  $i$ , so that  $f' \in M^\perp$ . Since we are only concerned with defining  $f'$  on  $S$ , the next lemma shows that such a function exists.

LEMMA 2. *Suppose that the measure space  $(S, \Sigma, \mu)$  contains no atoms, that  $f$  is a measurable function on  $S$  with  $|f| \leq 1$  a.e. and that  $g_1, g_2, \dots, g_k$  are in  $L_1(S, \Sigma, \mu)$ . Then there exists a measurable function  $f'$  on  $S$  with  $|f'| = 1$  a.e. such that  $\int_S f' g_i d\mu = \int_S f g_i d\mu, i = 1, 2, \dots, k$ .*

PROOF. Such a function  $f'$  will exist if and only if there exists a measurable set  $B$  (which will be the set on which  $f' = 1$ ) such that  $\int_S f g_i = \int_B g_i - \int_{S \sim B} g_i = 2 \int_B g_i - \int_S g_i$ , and this is equivalent to  $\int_B g_i = 1/2 \int_S (1 + f) g_i, i = 1, 2, \dots, k$ . Now, by Liapunov's theorem [7], [8] if we define, for  $C$  in  $\Sigma, \phi(C) = (\int_C g_1, \int_C g_2, \dots, \int_C g_k)$ , then the image  $K$  (in  $R^k$ ) of  $\Sigma$  under the map  $\phi$  is a compact convex set. On the other hand, if we define, for  $|g| \leq 1$ ,

$$\psi(g) = \left( 1/2 \int_S (1+g)g_1, 1/2 \int_S (1+g)g_2, \dots, 1/2 \int_S (1+g)g_k \right),$$

then  $\psi$  is a weak\* continuous affine map from the unit ball  $U$  of  $L_\infty$  into  $R^k$ , hence  $\psi(U)$  is also compact and convex. Our problem is to find  $B$  in  $\Sigma$  such that  $\phi(B) = \psi(f)$ , and for this it suffices to show that  $\psi(f) \in K$ . This will certainly be true if  $\psi(U) \subset K$ , and for *this* it is sufficient that  $\psi(\text{ext } U) \subset K$ , where  $\text{ext } U$  is the set of extreme points of  $U$ . Now, as is well known (and easy to prove), if  $g \in \text{ext } U$ , then  $|g| = 1$  a.e., i.e. there exists  $C$  in  $\Sigma$  such that  $g = 2\chi_C - 1$ . It follows that  $\psi(g) = (\int_C g_1, \dots, \int_C g_k) = \phi(C) \in K$ , which completes the proof of the lemma.

[It is interesting (but not pertinent to the questions at hand) to note that one can easily deduce the Liapunov theorem from this lemma. We make two observations which help in the proof. If  $\mu_1, \mu_2, \dots, \mu_n$  define a vector measure on  $(T, \Sigma)$ , then they may be identified with their Radon-Nykodym derivatives  $g_1, \dots, g_n$  with respect to  $\mu = |\mu_1| + |\mu_2| + \dots + |\mu_n|$ . Also, if  $0 \leq h \leq 1$  is an element of  $L_\infty(T, \Sigma, \mu)$ , then by applying the lemma to  $f = 2h - 1$ , one obtains a set  $S$  in  $\Sigma$  such that  $\mu_k(S) = \int_S h g_k d\mu$ ,  $k = 1, 2, \dots, n$ .]

It remains to define  $f''$  so that  $\|f''\| = 1$ ,  $f'' \in M^\perp$ ,  $f'' = f'$  on  $T \sim A$  and  $|f''| = 1$  on all but (at most)  $n - 1$  atoms in  $A$ . It suffices, then, to define  $f''$  on  $A$  such that  $|f''| = 1$  on all but  $n - 1$  atoms and such that  $\int_A f'' g_i = \int_A f' g_i$ , where  $g_1 = g, g_2, \dots, g_n$  form a basis for  $M$ . Since all the functions involved are measurable, they are a.e. constant on atoms, so their restrictions to  $A$  may be considered to be sequences. If  $A$  has only finitely many atoms, the rest of the proof follows by induction from the next lemma, so we will assume that  $A = \bigcup_{k=1}^\infty A_k$  ( $A_k$  essentially disjoint atoms) and define the obvious isometry between  $L_1(A)$  and the space  $l_1$  by  $\alpha(g)_k = g(A_k)\mu(A_k)$ . Thus, there is no loss in generality in assuming that the  $g$ 's (which we now write as  $g^1, g^2, \dots, g^n$ ) are elements of  $l_1$  and that  $f' \in l_\infty$ , with  $|f'_k| < 1$  for all  $k$ . Our problem requires us to produce  $f''$  in  $l_\infty$  with  $|f''_k| = 1$  (for all but at most  $n - 1$  integers  $k$ ) such that  $\sum f''_k g_k^i = \sum f'_k g_k^i$ ,  $i = 2, 3, \dots, n$ . We first prove a simple lemma concerning *finite* sequences.

**LEMMA 3.** *Suppose that  $u^1, u^2, \dots, u^{n-1}$  are elements of  $R^n$  and that  $z \in R^n$ , with  $|z_k| < 1$  for each  $k$ . Then there exists  $z'$  in  $R^n$ , with  $|z'_k| \leq 1$  for each  $k$ ,  $|z'_k| = 1$  for at least one  $k$ , and  $\sum z'_k u_k^i = \sum z_k u_k^i$ ,  $i = 1, 2, \dots, n - 1$ .*

**PROOF.** The subspace spanned by  $u^1, \dots, u^{n-1}$  has dimension at most  $n - 1$ , so there exists  $a$  in  $R^n$ ,  $a \neq 0$ , such that  $(\lambda a, u^i = 0$  for

in  $R$ ,  $i=1, 2, \dots, n-1$ . For small  $\lambda \neq 0$ ,  $|z_k + \lambda a_k| < 1$ , so we can choose  $\lambda_0$  such that  $|z_k + \lambda_0 a_k| = 1$  for at least one  $k$  and  $|z_j + \lambda_0 a_j| \leq 1$  for  $j \neq k$ . Clearly,  $z' = z + \lambda_0 a$  satisfies the above requirements.

Returning to the proof of the theorem, let  $S(f')$  denote the set of integers  $k$  such that  $|f'_k| < 1$  and suppose  $S(f')$  contains  $n$  or more integers. Write these in their natural order and consider the elements of  $R^n$  obtained by restricting  $f', g^2, \dots, g^n$  to the set  $S_1$  consisting of the first  $n$  integers in  $S(f')$ . (Recall that  $g^1 = 0$  on  $S(f')$ .) It follows from the above lemma (applied to these elements of  $R^n$ ) that we can choose  $h^1 \in l_\infty$ ,  $\|h^1\| \leq 1$ , such that  $h^1_k = f'_k$  outside of  $S_1$ ,  $h^1 \in M^\perp$ , and  $|h^1_k| = 1$  for at least one  $k$  in  $S_1$ . Let  $S_2$  be the first  $n$  elements of  $S(h^1)$  and choose  $h^2$  such that  $\|h^2\| \leq 1$ ,  $h^2 = h^1$  outside  $S_2$ ,  $h^2 \in M^\perp$  and  $|h^2_k| = 1$  for at least one  $k$  in  $S_2$ . Continuing in this manner we obtain a sequence  $\{h^i\}$  of elements of  $M^\perp$  such that  $\|h^i\| = 1$  and for each integer  $k$  (with at most  $n-1$  exceptions) there exists  $j_k$  such that  $h^j_k = h^{j_k}_k$  for  $j \geq j_k$  and  $|h^{j_k}_k| = 1$ . Since  $\{h^i\}$  lies in a weak\* compact set, it has a subsequence which converges to an element  $f''$  in  $M^\perp$ . It follows easily that  $S(f'')$  contains at most  $n-1$  integers, and the proof of Theorem 1 is complete.

We can now prove our corollary concerning property  $U$ .

**COROLLARY 2.** *Suppose that  $M$  is a subspace of finite codimension  $n$  in  $C(T)$ ,  $T$  compact Hausdorff. Then  $M$  fails to have property  $U$  if and only if there exists a bounded Baire measurable function  $f$  on  $T$  such that  $|f| \leq 1$ ,  $\int_T f d\mu = 0$  for each  $\mu$  in  $M^\perp$ , and  $|f(t)| < 1$  for at most  $n-1$  points  $t$  of  $T$ .*

**PROOF.** Suppose that such a function exists, and let  $\mu_1, \dots, \mu_n$  be a basis for  $M^\perp$ . Let  $\mu = |\mu_1| + |\mu_2| + \dots + |\mu_n|$  and write  $\mu_i = f_i \mu$ , where  $f_i = d\mu_i/d\mu$  is the Radon-Nykodym derivative of  $\mu_i$  with respect to  $\mu$ . The functions  $f_i$  generate an  $n$ -dimensional subspace  $M_0$  of  $L_1(T, \Sigma, \mu)$  ( $\Sigma$  the Baire subsets of  $T$ ) and it is clear that if  $M_0$  is not a Čebyšev subspace of  $L_1(T, \Sigma, \mu)$ , then  $M^\perp$  is not a Čebyšev subspace of  $C(T)^*$ . (Note that  $L_1(T, \Sigma, \mu)$  is isometric with the subspace of measures in  $C(T)^*$  which are absolutely continuous with respect to  $\mu$ , and that this isometry maps  $M_0$  onto  $M$ .) Now,  $f \in L_\infty(T, \Sigma, \mu)$ ,  $f \in M_0^\perp$ ,  $\|f\| = 1$  and  $S(f)$  consists of at most  $n-1$  points, so by Theorem 1,  $M_0$  is not a Čebyšev subspace and this half of the proof is complete. To prove the converse, suppose that  $M$  is non- $U$ , so that  $M^\perp$  is not a Čebyšev subspace of  $C(T)^*$ . Let  $\nu \in C(T)^*$ ,  $\nu \notin M^\perp$ , have two distinct nearest points in  $M^\perp$ , and choose a basis  $\mu_1, \dots, \mu_n$  for  $M^\perp$ . Let  $\mu = |\nu| + |\mu_1| + \dots + |\mu_n|$  and write  $\nu = h\mu$ ,

$\mu_i = f_i \mu$ ,  $i = 1, \dots, n$ . Thus, the subspace  $M_0$  in  $L_1(T, \Sigma, \mu)$  generated by  $f_1, \dots, f_n$  is not a Čebyšev subspace, since  $k$  has two nearest points in  $M_0$ , so by Theorem 1 there exists a function  $f_0$  in  $L_\infty(T, \Sigma, \mu)$  of norm 1 with  $\int f_0 f_i d\mu = \int f_0 d\mu_i = 0$  for  $i = 1, \dots, n$  and  $S(f_0)$  consists of  $k$  atoms,  $0 \leq k \leq n-1$ . Now  $f_0$  is defined on  $T$  only to within a Baire set  $B$  of  $\mu$ -measure zero. Define  $f$  on  $T$  by setting  $f = f_0$  on  $T \sim B$ ,  $f = 1$  on  $B$ . Since  $\mu(B) = 0$ , we have  $|\mu_i|(B) = 0$  for each  $i$ , so  $\int f d\mu_i = 0$ . The atoms (if any) in  $S(f_0)$  are, of course, simply points of  $T$ , and the proof is complete.

It is obvious that the first ("if") part of this proof is valid under the (stronger) hypothesis that  $f$  be an element of  $M$  with  $|f(t)| = 1$  for all but at most  $n-1$  points; and it would be nice indeed if the existence of such a function were necessary (as well as sufficient) for  $M$  to be non- $U$ . This is not the case, however, as the following example will show. (Recall that the space  $c$  of all convergent sequences  $\{x_n\}$  may be regarded as the space of all continuous functions  $f$  on the points  $\{1/n\} \cup \{0\}$ , with  $f(1/n) = x_n$ ,  $f(0) = \lim x_n$ .) Our theorem on  $l_1$  says the following about  $c$ :

*If  $M$  has finite codimension  $n$  in  $c$ , then  $M$  is non- $U$  if and only if there exists  $z$  in  $M^{\perp} \subset l_\infty$  such that  $\|z\| = 1$  and  $|z_k| < 1$  for at most  $n-1$  integers  $k$ .*

Thus (in particular), if there exists  $z$  in  $M$  such that  $\|z\| = 1$  and  $S(z)$  has at most  $n-1$  points, then  $M$  is non- $U$ . The converse of *this* assertion is false, however. In the example below, let  $M$  be the hyperplane  $\{x \in c: \sum x_i y_i = 0\}$ . There exists  $z$  in  $(Ry)^\perp = M^{\perp}$  such that  $|z_k| = 1$  for all  $k$  (so  $M$  is non- $U$ ), but there is no  $z$  in  $M$  with this property.

**EXAMPLE.** *There exists a sequence  $y$  in  $l_1$  such that  $(y, z) = \sum y_n z_n = 0$  for some  $z$  in  $l_\infty$  with  $|z_n| = 1$  for all  $n$ , but there is no  $z$  in  $c$  with this property.*

**PROOF.** Let  $y_n = (-1)^{n+1} 2^{-n}$ ,  $n = 1, 2, 3, \dots$ . If  $z_1 = 1$ ,  $z_n = (-1)^n$ ,  $n \geq 2$ , then  $\sum y_n z_n = 0$ . If  $x \in c$ , then  $(x, y) = y_1 \lim x_n + \sum_{n=2}^\infty x_n y_n$ . If  $|x_n| = 1$  for all  $n$ , then there exists  $n_0$  such that  $x_k = \lim x_n (= \pm 1)$  for  $k \geq n_0$ . This fact, together with some simple computations, shows that  $(x, y) = 0$  is impossible for such an element  $x$ .

What about the *existence* of subspaces of finite codimension in  $C(T)$  which have property  $U$ ? The question is answered by the following result.

**COROLLARY 3.** *If  $t_1, t_2, \dots, t_n$  are distinct points of  $T$ , then the subspace  $M = \{f \in C(T): f(t_i) = 0, i = 1, 2, \dots, n\}$  has codimension  $n$  in  $C(T)$  and has property  $U$ .*

PROOF. This is immediate from Corollary 2 and the fact that in this case,  $M^\perp$  is generated by the point masses at  $t_1, \dots, t_n$ .

In conclusion, we note that the hypothesis that  $(T, \Sigma, \mu)$  be  $\sigma$ -finite was not essential. Indeed, suppose that  $M$  is a subspace of  $L_1(T, \Sigma, \mu)$  generated by  $g_1, g_2, \dots, g_n$  and that  $g_0 \in L_1(T, \Sigma, \mu)$ ,  $g_0 \notin M$ . Let  $T_0$  be the union of the supports of the  $g$ 's i.e.,  $T_0 = T \sim \bigcap_{k=1}^n Z(g_k)$ ; then  $T_0$  is  $\sigma$ -finite. Furthermore, if we denote by  $M_0$  the subspace of  $L_1(T_0, \Sigma, \mu)$  obtained by restricting  $M$  to  $T_0$ , then  $g_0$  has a unique nearest point in  $M$  if and only if the restriction of  $g_0$  to  $T_0$  has a unique nearest point in  $M_0$ . Using this remark, it is easy to extend Theorem 1 to the general case.

*Added in proof.* A simple one-page proof of Liapunov's theorem, using the Krein-Milman theorem, has been given by Joram Lindenstrauss, *A short proof of Liapunoff's convexity theorem*, J. Math. Mech. (to appear).

#### BIBLIOGRAPHY

1. N. I. Ahieser and M. G. Krein, *Some questions in the theory of moments*, Transl. Math. Monographs, Vol. 2, Amer. Math. Soc., Providence, R. I., 1962.
2. A. L. Garkavi, *On best approximation by elements of infinite dimensional subspaces of a certain class*, Mat. Sb. (104) 62 (1963), 104-120. (Russian)
3. ———, *Approximative properties of subspaces of finite defect in the space of continuous functions*, Dokl. Adad. Nauk SSSR 155 (1964), 513-516; English transl., Soviet Math. Dokl. 5 (1964), 440-443.
4. ———, *On uniqueness of the solutions of the L-problem of moments*, Izv. Akad. Nauk SSSR 28 (1964), 553-570. (Russian)
5. ———, *On Čebyšev and almost-Čebyšev subspaces*, Izv. Akad. Nauk SSSR 28 (1964), 799-818. (Russian)
6. A. Haar, *Die Minkowskische Geometrie und die Annäherung an stetige Funktionen*, Math. Ann. 78 (1918), 294-311.
7. P. R. Halmos, *The range of a vector measure*, Bull. Amer. Math. Soc. 54 (1948), 416-421.
8. A. A. Liapunov, *On completely additive vector-functions*, Izv. Akad. Nauk SSSR 4 (1940), 465-478. (Russian. French summary)
9. R. R. Phelps, *Uniqueness of Hahn-Banach extensions and unique best approximation*, Trans. Amer. Math. Soc. 95 (1960), 238-255.
10. ———, *Čebyšev subspaces of finite codimension in  $C(X)$* , Pacific J. Math. 13 (1963), 647-655.

UNIVERSITY OF WASHINGTON