

GENERALIZATION OF DEHN'S RESULT ON THE CONJUGACY PROBLEM

SEYMOUR LIPSCHUTZ¹

1. Introduction. In 1911 Max Dehn [2] solved the conjugacy problem for the fundamental group G_k of a two-dimensional orientable closed manifold of genus k . Now G_k can be finitely presented in terms of $2k$ generators and one defining relation as follows:

$$G_k = gp(a_1, b_1, a_2, b_2, \dots, a_k, b_k; a_1^{-1}b_1^{-1}a_1b_1 \dots a_k^{-1}b_k^{-1}a_kb_k = 1).$$

If $k > 1$ then G_k is the free product of two free groups with an infinite cyclic group amalgamated. Hence the following theorem generalizes Dehn's result.

THEOREM. *Let G be the free product of free groups F_λ with an infinite cyclic group amalgamated. Then G has a solvable conjugacy problem.*

We remark that the word problem has already been solved for G (e.g. Lipschutz [4]) and that Greendlinger [3] extended Dehn's result to a class of groups which overlap those of our theorem.

A theorem of Solitar [5] which gives conditions for the solution of the conjugacy problem for free products with amalgamations is used in the proof of our theorem.

2. Preliminary results. Let F be a free group. We will let u, v and w denote freely reduced nonempty words in F ; the letters a, b, c, \dots will denote a fixed set of generators of F and hence the letters of the words in F . We use the notation

$$\begin{aligned} l(u) & \text{ for the length of } u; \\ v \equiv u_1u_2 & \text{ if } v \text{ is identical to the letters of } u_1 \text{ followed by the letters} \\ & \text{ of } u_2. \end{aligned}$$

Recall that $u = a_1a_2 \dots a_n$ is cyclically reduced iff $a_n \neq a_1^{-1}$.

LEMMA 1. *Suppose u is cyclically reduced and $v \neq u^{-1}w$. If the first letter in the product*

$$(1) \quad uvv^n$$

is absorbed when (1) is free reduced, then $n < 0$ and u and v commute.

PROOF. We will perform cancellations in (1) from left to right.

Received by the editors October 2, 1965.

¹ This work has been sponsored by the National Science Foundation Grant GP 4255.

Since u is cyclically reduced, cancellations can only take place in two places, on either side of v . Let u_1 denote the first u in (1) and let $u_1 = a_1 a_2 \cdots a_m$. Then (1) can be written

$$a_1 a_2 \cdots a_m u v u^n.$$

Now a_1 can be absorbed only if a_m is absorbed. Since $v \neq u^{-1} w$, a_m cannot be absorbed by a letter of v and, furthermore, a_m can be absorbed only after all the letters of v are absorbed. We now consider the step in the free reduction of (1) just before a_m is absorbed. If $n \geq 0$ and a_m is absorbed by the letter a_i of $u_2 = u = a_1 a_2 \cdots a_m$, then (1) must be reduced to the form

$$a_1 a_2 \cdots a_n a_i a_{i+1} \cdots a_n u^r.$$

Thus if a_1 is absorbed then $(a_i a_{i+1} \cdots a_n)^2 = 1$. But there are no elements of finite order in a free group; hence if a_1 is absorbed, then $n < 0$ as claimed.

Now, for $n < 0$, let a_m be absorbed by the letter a_i^{-1} of $u_2^{-1} = u^{-1} = a_m^{-1} a_{m-1}^{-1} \cdots a_1^{-1}$. Then (1) is reduced to the form

$$a_1 a_2 \cdots a_m a_i^{-1} a_{i-1}^{-1} \cdots a_1^{-1} u^r, \quad r < 0.$$

If $i = m$, then $u v u^{n-r-1} = 1$ which implies that v is a power of u and so commutes with u . On the other hand, if $i < m$ and a_1 is absorbed, then

$$(a_1 a_2 \cdots a_i)(a_{i+1} \cdots a_m)(a_i^{-1} \cdots a_1^{-1})(a_n^{-1} \cdots a_{i+1}^{-1}) = 1.$$

Hence $a_1 a_2 \cdots a_i$ and $a_{i+1} \cdots a_m$ commute and so are powers of the same element, say w . Thus u is a power of w . Furthermore

$$u v u a_m^{-1} \cdots a_{i+1}^{-1} = 1$$

and so v is also a power of w . Accordingly, u and v commute.

COROLLARY 1. *Let u be cyclically reduced and let $m > l(v) + 2$. Then either*

$$m < l(u^m v u^n) + l(v) + 2$$

or u and v commute.

PROOF. We can write v in the form $v \equiv u^{-s} v^*$ where $0 \leq s \leq l(v)$ and $v^* \neq u^{-1} w$. Then $u^m v u^n = u^{m-s} v^* u^n$. Let u_1 denote the second u preceding v^* . Suppose the first letter of u_1 is not absorbed when $u^m v u^n$ is freely reduced; then

$$l(u^m v u^n) = l(u^{m-s} v^* u^n) > m - s - 2 \geq m - l(v) - 2.$$

On the other hand, if the first letter of u_1 is absorbed then, by Lemma 1, u and v^* commute and so are powers of some element w . Hence v is also a power of w and so u and v commute as claimed.

REMARK. This corollary extends a lemma of Baumslag [1, p. 428].

LEMMA 2. *Given u and v , one can decide whether or not u is a power of v or is conjugate to a power of v ; and such a power is unique.*

PROOF. If $u = w^{-1}u^mw = (w^{-1}uw)^m$, then

$$|m| \leq l(u) = l((w^{-1}uw)^m).$$

Thus we only need to test a finite number of cases. The lemma now follows from the fact that the conjugacy problem is solved for free groups, and the fact that different powers of an element in a free group are in different conjugate classes.

3. Main results.

LEMMA 3. *Let G be the free product of free groups F_λ with an infinite cyclic group amalgamated. Let h be a generator of the amalgamated cyclic group and suppose h is cyclically reduced in each factor. Let u and v be elements of G with free product length n , say $u = u_1u_2 \cdots u_n$ and $v = v_1v_2 \cdots v_n$ where u_i and v_j appear in individual factors. Then we can decide if there exists an m such that*

$$(2) \quad h^m u_1 u_2 \cdots u_n h^{-m} = v_1 v_2 \cdots v_n.$$

PROOF. Since the word problem has been solved for G , we can decide if h and u_1 commute. Suppose h and u_1 do not commute. By properties of free products with amalgamations, if (2) holds then $v_1^{-1}h^m u_1$ belongs to the amalgamated cyclic group; say $v_1^{-1}h^m u_1 = h^{-r}$. Then

$$h^m u_1 h^r = v_1.$$

Hence, by Corollary 1, $|m| < l(v_1) + l(u_1) + 2$. The lemma now follows, in the case that h and u_1 do not commute, from the fact that the word problem has been solved for G and that there are only a finite number of m to consider.

Suppose h and u_1 do commute. Then $v_1^{-1}u_1$ belongs to the amalgamated cyclic group; say $v_1^{-1}u_1 = h^s$ and so $u_1 = v_1 h^s$. (By Lemma 2, s is unique and can be found.) Substituting in (2) and using the fact that u_1 and h commute gives

$$v_1 h^m (h^s u_2) u_3 \cdots u_n h^{-m} = v_1 v_2 \cdots v_n$$

or

$$h^m(h^*u_2)u_3 \cdots u_n h^{-m} = v_2 v_3 \cdots v_n.$$

The lemma now follows from induction.

PROOF OF THEOREM. Let h be a generator of the amalgamated cyclic group. The theorem is proven if we can decide whether or not any two free product cyclically reduced elements u and v in G are conjugate. We can also assume without loss in generality that h is cyclically reduced in the factors in which u and v appear (by considering appropriate generating sets of these factors).

Case I. u and v have different free product lengths. Then u and v are not conjugate.

Case II. u and v have length 1; say $u \in F_i$ and $v \in F_j$. By Solitar's theorem, either (a) $i = j$ and u and v are conjugate elements of F_i ; or (b) there exist a sequence of elements h_1, h_2, \dots, h_s in the amalgamated subgroup such that u and h_1 are conjugate elements of F_i , h_1 and h_2 are conjugate elements of F_{λ_1} , h_2 and h_3 are conjugate elements of $F_{\lambda_2}, \dots, h_s$ and v are conjugate elements of F_j . But the elements of a cyclic subgroup of any factor are in different conjugate classes; hence $h_1 = h_2 = \dots = h_s = h^m$ for some integer m .

Now in case (a) we can decide if u and v are conjugate elements of F_i since the conjugacy problem has been solved for free groups. In case (b) we can decide, by Lemma 2, if u is conjugate to a power h^m of h in F_i and if v is conjugate to the same power h^m in F_j . Consequently, in Case II, we can decide if u and v are conjugate elements in G .

Case III. u and v have length $n > 1$; say $u = u_1 \cdots u_n$ and $v = v_1 \cdots v_n$ where u_i and v_j appear in factors. By Solitar's theorem, u and v are conjugate iff for some i and some j ,

$$h^m u_i \cdots u_n u_1 \cdots u_{i-1} h^{-m} = v_j \cdots v_n v_1 \cdots v_{j-1}.$$

By Lemma 3, we can decide if the above is true. Since there are only n^2 cases to consider, we can decide if u and v are conjugate elements of G .

BIBLIOGRAPHY

1. G. Baumslag, *On generalized free products*, Math. Z. **78** (1962), 423-438.
2. M. Dehn, *Über unendliche diskontinuierliche Gruppen*, Math. Ann. **71** (1911), 116-144.
3. M. Greendlinger, *On Dehn's algorithm for the conjugacy and word problems, with applications*, Comm. Pure Appl. Math. **13** (1960), 641-677.
4. S. Lipschutz, *An extension of Greendlinger's results on the word problem*, Proc. Amer. Math. Soc. **15** (1964), 37-43.
5. W. Magnus, A. Karass and D. Solitar, *Combinatorial group theory*, Wiley, New York, 1965.