

## ON A BOUNDED INCREASING POWER SERIES

P. B. KENNEDY AND P. SZÜSZ

Recently [1] H. S. Shapiro has shown that if  $k$  is a positive integer,  $\beta = 1/(2k)$  and  $\alpha = 1 - \beta$ , and if  $a_0 = 0$ ,  $a_n = n^{-\alpha} \cos n^\beta$  for  $n = 1, 2, 3, \dots$ , then the function

$$(1) \quad f(x) = \sum_0^{\infty} a_n x^n$$

has bounded variation on  $[0, 1)$  but the series

$$(2) \quad \sum_0^{\infty} a_n$$

is divergent. Given any  $\epsilon > 0$  we may by choosing  $k$  large enough ensure that

$$(3) \quad a_n = O(n^{-1+\epsilon}).$$

However the series (2) is certainly Abel summable and so the stronger condition

$$(4) \quad a_n = O(n^{-1})$$

would imply the convergence of (2), by Littlewood's Tauberian theorem. Thus Shapiro's example shows that we cannot weaken (4) to a condition of the form (3) in Littlewood's theorem by making the compensating assumption that  $f$  has bounded variation on  $[0, 1)$ .

In this note we prove a stronger negative result than that of Shapiro.

*THEOREM.* *Let  $\phi(n)$  be positive for all positive integers  $n$ , and let  $\phi(n) \uparrow \infty$  as  $n \rightarrow \infty$ . Then there is a function  $f$  of the form (1) which is increasing and bounded on  $[0, 1)$  and for which (2) is divergent although*

$$(5) \quad |a_n| < n^{-1}\phi(n) \quad \text{for all } n \geq 1.$$

*PROOF.* Let  $\{n_k\}$  ( $k = 1, 2, 3, \dots$ ) be an increasing sequence of integers satisfying

$$(6) \quad \phi(n_k) > k^2 \quad (k \geq 1).$$

Put

---

Received by the editors June 28, 1965.

$$(7) \quad a_n = \begin{cases} k^2/n & \text{if } k^2 n_k \leq n < (k^2 + 1)n_k, \quad k = 1, 2, 3, \dots \\ -k^2/n & \text{if } (k^2 + 1)n_k \leq n < (k^2 + 2)n_k, \quad k = 1, 2, 3, \dots \\ 0 & \text{for all other values of } n. \end{cases}$$

(7) defines  $a_n$  uniquely for all  $n \geq 0$ ; and if the summation is over the range  $k^2 n_k \leq n < (k^2 + 1)n_k$ , then

$$\sum a_n = k^2 \sum n^{-1} > k^2 n_k / \{(k^2 + 1)n_k\} \rightarrow 1,$$

so that (2) is divergent. Moreover for each value of  $n$ , either  $a_n = 0$  or else by (7) there is a positive integer  $k$  such that

$$n \mid a_n \mid = k^2, \quad k^2 n_k \leq n,$$

and so by (6) and the fact that  $\phi$  is increasing,

$$n \mid a_n \mid < \phi(n_k) \leq \phi(n),$$

which proves (5).

It remains to show that  $f$ , defined by (1), is increasing and bounded on  $[0, 1)$ . In fact, by (7),  $f(x) = \sum_1^\infty f_k(x)$ , where

$$f_k(x) = k^2 \sum_1 x^n/n - k^2 \sum_2 x^n/n,$$

$\sum_1$  being taken over the range  $k^2 n_k \leq n < (k^2 + 1)n_k$ , and  $\sum_2$  over the range  $(k^2 + 1)n_k \leq n < (k^2 + 2)n_k$ . It is easy to see that  $f'_k(x) \geq 0$  for all  $x$  on  $[0, 1)$ , and so  $f'(x) \geq 0$ , whence  $f$  is increasing on  $[0, 1)$ . Also, when  $x = 1$ ,  $\sum_1$  consists of  $n_k$  terms, each at most as big as  $1/(k^2 n_k)$ , and  $\sum_2$  consists of  $n_k$  terms, each at least as big as  $1/\{(k^2 + 2)n_k\}$ , and so

$$f_k(1) \leq 1 - k^2/(k^2 + 2) \sim 2/k^2.$$

Therefore  $f(1 - 0) = \sum_1^\infty f_k(1) < +\infty$ . Thus  $f$  is increasing and bounded on  $[0, 1)$ , and this proves the theorem.

#### REFERENCE

1. H. S. Shapiro, *A remark concerning Littlewood's Tauberian theorem*, Proc. Amer. Math. Soc. **16** (1965), 258-259.

UNIVERSITY OF YORK, ENGLAND AND  
PENNSYLVANIA STATE UNIVERSITY