

ARCS IN INVERSE LIMITS ON $[0, 1]$ WITH ONLY ONE BONDING MAP

WILLIAM S. MAHAVIER

1. Introduction. In this note, by a continuum we mean a non-degenerate, compact, connected metric space. It is known ([2] or [5]) that a continuum is chainable if and only if it is homeomorphic to the inverse limit of a sequence of maps from $[0, 1]$ onto $[0, 1]$. Indeed, the mappings may be required to be piecewise linear. Henderson has shown that a pseudo arc can be obtained as an inverse limit on $[0, 1]$ with only one bonding map [3], and the author has shown that not every chainable continuum can be so obtained [4]. We now show that if M is an inverse limit on $[0, 1]$ with only one bonding map, and the bonding map is piecewise monotone, then every subcontinuum of M contains an arc. Thus the set of all such continua is a proper subset of the set of chainable continua which are inverse limits on $[0, 1]$ with only one bonding map.

2. Definitions and notation. If each term of the sequence $g = \{g_i\}$ maps $[0, 1]$ onto $[0, 1]$, then the inverse limit of g , denoted by $\lim g$, is the subspace of the infinite cartesian product $[0, 1]^\infty$ consisting of all number sequences $\{x_i\}$ such that for each $i > 0$, $g_i(x_{i+1}) = x_i$. If f maps $[0, 1]$ onto $[0, 1]$, then $\lim f$ denotes $\lim g$ where $g = f, f, \dots$. By an interval we mean a nondegenerate closed subinterval of $[0, 1]$, and the statement that A is an inverse sequence (for f) means that A is a sequence A_1, A_2, \dots such that if $i > 0$, A_i is degenerate or an interval and $f(A_{i+1}) = A_i$. The set of all points of $\lim f$ such that for each $i > 0$, $x_i \in A_i$ is denoted by $\lim(f, A)$. By a subinverse sequence of A is meant an inverse sequence $B = \{B_i\}$ such that if $i > 0$, $B_i \subseteq A_i$. If each of I and I' is an interval, the statement that f maps I onto I' efficiently means that $f(I) = I'$ and no interior point of I maps onto an endpoint of I' .

3. Arcs in chainable continua. Our object in this section is to prove the following:

THEOREM. *If f is a piecewise monotone function from $[0, 1]$ onto $[0, 1]$ then each subcontinuum of $\lim f$ contains an arc.*

PROOF. Suppose there is a subcontinuum of $\lim f$ which contains no arc. Then there is an inverse sequence $A = \{A_i\}$ such that $\lim(f, A)$ contains no arc. There is an increasing number sequence $x_0 = 0, x_1,$

Received by the editors June 3, 1968.

$\dots, x_n = 1$ such that if $0 < i \leq n$, $f|_{[x_{i-1}, x_i]}$ is monotone. Let M denote the set of images under f of the numbers in this sequence and let ϵ denote a positive number such that if each of I and I' is an interval, and f maps I onto I' efficiently, and the length of I' is less than ϵ , then f is monotone on I . Let K denote the subset of $[0, 1]$ to which a number x belongs if and only if there is a point $\{p_i\}$ in $\lim f$ such that $p_1 = x$ and such that either (1) for some $i > 0$, $p_i \in M$ or (2) for some $i > 0$, $f^{-1}(p_i)$ contains an interval. K is countable and so there is a finite collection G' of nonoverlapping intervals filling up $[0, 1]$ such that each interval in G' is of length less than $\epsilon/2$, and no endpoint of an interval in G' , other than 0 or 1, is a number in K . Let G denote the intervals in G' which have neither 0 nor 1 as an endpoint. We now state an easily established lemma.

LEMMA 1. *If g maps $[0, 1]$ onto $[0, 1]$ and $\alpha = \{\alpha_i\}$ is an inverse sequence for g , and I is a subinterval of a term α_n of α , then there is a subinverse sequence $\{\beta_i\}$ of α such that $\beta_n = I$ and for each $i > n$, g maps β_{i+1} onto β_i efficiently.*

Applying Lemma 1, there is a subinverse sequence $A^0 = \{A^0(i)\}$ of A and an integer n_0 such that for each $i \geq n_0$, f maps $A^0(i+1)$ efficiently onto $A^0(i)$. If there is an n such that for $i > n$, f is monotone on $A^0(i)$, then $\lim(f, A^0)$ is an arc (see [1]). Since $\lim(f, A^0)$ is not an arc, there is a subsequence of A^0 each term of which is of length at least ϵ and so there is an interval g_1 in G which is a subset of each term of some subsequence of A^0 . Applying Lemma 1 again, there is a subinverse sequence $A^1 = \{A^1(i)\}$ of A^0 and an integer n_1 such that $A^1(n_1) = g_1$ and such that if $i \geq n_1$, f maps $A^1(i+1)$ efficiently onto $A^1(i)$. As before we note that since $\lim(f, A^1)$ is not an arc, there is an interval g_2 in G which is a subset of each term of some subsequence of A^1 . We continue to establish the existence of a sequence A^0, A^1, A^2, \dots , an increasing integer sequence n_1, n_2, \dots and a sequence g_1, g_2, \dots of members of G such that for each $i > 0$, (1) A^i is a subinverse sequence of A^{i-1} , (2) $A^i(n_i) = g_i$, (3) if $j \geq n_i$, f maps $A^i(j+1)$ efficiently onto $A^i(j)$ and (4) g_i is a subset of each term of a subsequence of A^{i-1} . There is an integer u and an integer $m < u$ such that $g_u = g_m$. The sequence A^m is a subinverse sequence of A^0 , $A^m(n_m) = g_m$, and g_m is a subset of each term of a subsequence of A^m . Further, as a consequence of the definition of G we have that if $j \geq n_m$, f is not constant on any interval which contains an endpoint of $A^m(j+1)$ and it follows that $A^m(j+1)$ is a component of $f^{-1}[A^m(j)]$. Let N denote an integer such that $A^m(n_m + N)$ contains $A^m(n_m)$. We shall show that if $j > 0$, $A^m(n_m + N + j)$ contains $A^m(n_m + j)$. Either $A^m(n_m + N + 1)$

$\supseteq A^m(n_m+1)$ or they do not intersect since $A^m(n_m+1)$ is a connected subset of $f^{-1}[A^m(n_m+N)]$ and $A^m(n_m+N+1)$ is a component of $f^{-1}[A^m(n_m+N)]$. Let k denote a positive integer such that $A^m(n_m+N+k)$ contains g_m . Since $f^N(g_m) \subseteq g_m$, then $f^N[A^m(n_m+N+k)]$ intersects g_m and thus intersects $A^m(n_m+N+k)$. But

$$f^N[A^m(n_m+N+k)] = A^m(n_m+k),$$

so $A^m(n_m+N+k)$ intersects $A^m(n_m+k)$ which implies that $A^m(n_m+N+1)$ intersects, and thus contains, $A^m(n_m+1)$. It follows inductively that if $j > 0$, $A^m(n_m+N+j)$ contains $A^m(n_m+j)$. Now let $g = f^N$ and for each $j > 0$ let $B(j) = A^m[n_m+(j-1)N]$. Then g is a piecewise monotone map of $[0, 1]$ onto $[0, 1]$, $B = \{B(i)\}$ is an inverse sequence (for g), and for each $j > 0$, $B(j+1) \supseteq B(j)$ and g maps $B(j+1)$ efficiently onto $B(j)$. Further, $\lim(g, B)$ is homeomorphic to a subcontinuum of $\lim(f, A)$ and so the following lemma applies to complete our proof.

LEMMA 2. *If g maps $[0, 1]$ onto $[0, 1]$ and is piecewise monotone and $B = \{B_i\}$ is an inverse sequence (for g) such that for each $i > 0$, B_i is a subinterval of B_{i+1} and g maps B_{i+1} efficiently onto B_i , then $\lim(g, B)$ contains an arc.*

PROOF. For each $i > 0$, let $B_i = [a_i, b_i]$. The sequence $\{a_i\}$ is non-increasing and converges to a number a . Similarly, $\{b_i\}$ converges to a number b . Further, there is a number x such that the point (x, x, \dots) is in $\lim(g, B)$. Suppose that there is a positive integer n such that $g(b_{n+2}) = a_{n+1}$ and $g(a_{n+1}) = a_n$. According as $g(b_{n+3}) = b_{n+2}$ or $g(b_{n+3}) = a_{n+2}$, we have that either $g([b_{n+2}, b_{n+3}])$ contains x and is nondegenerate or $g([a_{n+3}, a_{n+1}])$ contains x and is nondegenerate. But g is piecewise monotone and there do not exist infinitely many mutually disjoint intervals such that if I is one of them, $g(I)$ contains x and is nondegenerate. So there is an integer N' such that if $n > N'$, then it is not true that $g(b_{n+2}) = a_{n+1}$ and $g(a_{n+1}) = a_n$. It follows that there is an integer N such that either (1) if $n > N$, $g(a_{n+1}) = a_n$, or (2) if $n > N$, $g(a_{n+1}) = b_n$. Suppose first that (1) holds and that $b \neq x$. There is a number u , $x < u < b$, such that g is monotone on $[u, b]$ and an integer k such that if $j \geq k$, $b_j > u$. If $g(u) \leq u$, there is a point $\{u_i\}$ in $\lim g$ such that $u_1 = u$ and for $i > 0$, $u_i \leq u_{i+1} < b_{k+i}$. Then for $i > 0$, $g([u_{i+1}, b_{k+i}]) = [u_i, b_{k+i-1}]$ and g is monotonic on $[u_{i+1}, b_{k+i}]$. It follows that the set of all points $\{p_i\}$ of $\lim g$ such that for $i > 0$, $p_{k+i} \in [u_{i+1}, b_{k+i}]$ is an arc which is a subset of $\lim(g, B)$. If $g(u) > u$, then there is a number s and a number t such that (1) $g(s) = s$, (2) $x \leq s < t < u$, (3) if $y \in [s, t]$, $g(y) \geq y$ and (4) g is monotonic on $[s, t]$. There

is a point $\{t_i\}$ in $\lim g$ such that $t_1 = t$ and for each $i > 0$, $s < t_{i+1} \leq t_i$. For each $i > 0$, $g([s, t_{i+1}]) = [s, t_i]$ and g is monotonic on $[s, t_{i+1}]$, and it follows that the set of all points $\{p_i\}$ of $\lim g$ such that for $i \geq 0$, $p_{k+i} \in [s, t_{i+1}] \subseteq [a_{k+i}, b_{k+i}]$ is an arc which is a subset of $\lim(g, B)$. Thus we have that $\lim(g, B)$ contains an arc in case there is an integer N such that if $n > N$, $g(a_{n+1}) = a_n$ and $b \neq x$. An analogous argument suffices in case $a \neq x$ and it remains to show that $\lim(g, B)$ contains an arc in case there is an integer N such that if $n > N$, $g(a_{n+1}) = b_n$. In this case we consider the piecewise monotone map g^2 of $[0, 1]$ onto $[0, 1]$ and the inverse sequence $B' = B_1, B_3, B_5, \dots$. There is an integer k such that if $i > k$, $g^2(a_{2i+1}) = a_{2i-1}$, so an argument analogous to that given above applies to show that $\lim(g^2, B')$ contains an arc. But $\lim(g, B)$ is homeomorphic to $\lim(g^2, B')$ and this completes our proof.

REFERENCES

1. C. E. Capel, *Inverse limit spaces*, Duke Math. J. 21 (1954), 233–245.
2. H. Freudenthal, *Entwicklungen von Räumen und ihren Gruppen*, Compositio Math. 4 (1937), 145–234.
3. G. W. Henderson, *The pseudo-arc as an inverse limit with one binding map*, Duke Math. J. 31 (1964), 421–425.
4. W. S. Mahavier, *A chainable continuum not homeomorphic to an inverse limit on $[0, 1]$ with only one bonding map*, Proc. Amer. Math. Soc. 18 (1967), 284–286.
5. S. Mardesić and J. Segal, *ϵ -mappings and inverse limits*, Trans. Amer. Math. Soc. 109 (1963), 146–164.

EMORY UNIVERSITY