

## THE FUNCTIONAL EQUATION OF SOME DIRICHLET SERIES

BRUCE C. BERNDT<sup>1</sup>

**ABSTRACT.** The functional equation for two classes of Dirichlet series is established. These Dirichlet series involve primitive characters and can be regarded as generalizations of Dirichlet's  $L$ -functions or of Epstein's zeta-functions. One class is also a generalization of some series studied by Stark.

Our objective here is to derive the functional equation for two classes of Dirichlet series. Both classes of series may be regarded as generalizations of Epstein's zeta-functions or of Dirichlet's  $L$ -functions. One class is also a generalization of Stark's  $L$ -functions [5], and so it will be convenient to use the notation in [5].

Let  $x$  denote an  $m$ -dimensional vector and  $Q(x)$  a positive definite quadratic form in  $m$  variables. We let  $(x, y)$  denote the inner product of two  $m$ -dimensional vectors  $x$  and  $y$ . Write  $2Q(x) = (xF, x)$ , where  $F$  is a symmetric  $m \times m$  matrix of real numbers. The discriminant  $d > 0$  of  $Q$  is defined to be the determinant of  $F$ . The inverse  $Q^{-1}$  of  $Q$  is defined by  $2Q^{-1}(x) = (x d F^{-1}, x)$ . The Epstein zeta-functions are defined for  $\sigma = \operatorname{Re} s > m/2$  by

$$(1) \quad Z(s, Q; g, h) = \sum_n' e^{2\pi i(h, n)} \{Q(n + g)\}^{-s},$$

where  $g$  and  $h$  are  $m$ -dimensional real vectors and the sum is over all integral vectors  $n$  with  $Q(n + g) \neq 0$ . Epstein [3, p. 207] showed that  $Z(s, Q; g, h)$  satisfies the functional equation (the definitions of the discriminant and inverse differ in [3] and [5])

$$(2) \quad \left(\frac{d^{1/m}}{2\pi}\right)^s \Gamma(s) Z(s, Q; g, h) \\
 = e^{-2\pi i(\sigma, h)} \left(\frac{d^{(m-1)/m}}{2\pi}\right)^{m/2-s} \Gamma\left(\frac{1}{2}m - s\right) Z\left(\frac{1}{2}m - s, Q^{-1}; h, -g\right).$$

We now give our generalizations of (1). Throughout the sequel all characters are primitive. For  $\sigma > m/2$  define

Received by the editors September 17, 1970.

*AMS 1969 subject classifications.* Primary 1041; Secondary 1040.

*Key words and phrases.* Epstein zeta-function, Dirichlet  $L$ -function, primitive character, functional equation, quadratic form.

<sup>1</sup> Research supported by NSF Grant 21335.

$$Z(s, \chi, Q; g, h) = \sum'_n \prod_{j=1}^m \chi_j(n_j) e^{2\pi i(n, h)} \{Q(n + g)\}^{-s},$$

where  $n = (n_1, \dots, n_m)$ ,  $g$  and  $h$  are as above,  $\chi_j$  is a character (mod  $k_j$ ),  $1 \leq j \leq m$ , and  $k_j$  is any positive integer such that a primitive character (mod  $k_j$ ) exists. Secondly, for  $\sigma > m/2$  define

$$L(s, \chi, Q; g, h) = \sum'_n \chi(Q(n)) e^{2\pi i(n, h)} \{Q(n + g)\}^{-s},$$

where now  $Q$  has integral coefficients. We shall use the Gauss sum

$$G(r, \chi) = \sum_{j=1}^{k-1} \chi(j) e^{2\pi i r j / k},$$

where  $r$  is an integer and  $\chi$  is a character (mod  $k$ ). Furthermore, we write  $G(\chi) = G(1, \chi)$ .

**THEOREM 1.**  *$Z(s, \chi, Q; g, h)$  can be continued to a meromorphic function of  $s$  that is either entire or has but one pole, a simple one at  $s = m/2$ . Furthermore,*

$$\begin{aligned} (3) \quad & \prod_{j=1}^m G(\bar{\chi}_j) \left(\frac{d^{1/m}}{2\pi}\right)^s \Gamma(s) Z(s, \chi, Q; g, h) \\ & = e^{-2\pi i(\sigma, h)} \left(\frac{d^{(m-1)/m}}{2\pi}\right)^{m/2-s} \Gamma(\tfrac{1}{2}m - s) Z(\tfrac{1}{2}m - s, \bar{\chi}, Q^{-1}/k; h, -g/k). \end{aligned}$$

Here,  $Q^{-1}/k$  denotes  $Q^{-1}$  but with  $x_j$  replaced by  $x_j/k_j$ ,  $1 \leq j \leq m$ , and  $g/k = (g_1/k_1, \dots, g_m/k_m)$ .

**PROOF.** Since  $\chi$  is primitive, we have the factorization theorem [2, pp. 67–68],  $G(r, \bar{\chi}) = \chi(r)G(\bar{\chi})$ . Thus, for  $\sigma > m/2$ ,

$$\begin{aligned} (4) \quad & \prod_{j=1}^m G(\bar{\chi}_j) Z(s, \chi, Q; g, h) \\ & = \sum'_n \prod_{j=1}^m G(n_j, \bar{\chi}_j) e^{2\pi i(n, h)} \{Q(n + g)\}^{-s} \\ & = \sum_{j_1=1}^{k_1-1} \cdots \sum_{j_m=1}^{k_m-1} \bar{\chi}_1(j_1) \cdots \bar{\chi}_m(j_m) Z(s, Q; g, h + j/k), \end{aligned}$$

where  $j/k = (j_1/k_1, \dots, j_m/k_m)$ . Since  $Z(s, Q; g, h)$  can be analytically continued to an entire function if  $h$  is nonintegral or to a function analytic everywhere except for a simple pole at  $s = m/2$  if  $h$  is integral [3, p. 207], the first part of the theorem follows directly from (4). We

now employ (2) to obtain, for all  $s$ ,

$$\begin{aligned}
 \prod_{j=1}^m G(\bar{\chi}_j) Z(s, \chi, Q; g, h) &= d^{(m-1)/2-s} (2\pi)^{2s-m/2} \frac{\Gamma(\frac{1}{2}m-s)}{\Gamma(s)} \\
 (5) \quad &\cdot \sum_{j_1=1}^{k_1-1} \cdots \sum_{j_m=1}^{k_m-1} \bar{\chi}_1(j_1) \cdots \bar{\chi}_m(j_m) \\
 &\cdot e^{-2\pi i(\sigma, h+j/k)} Z(\frac{1}{2}m-s, Q^{-1}; h+j/k, -g).
 \end{aligned}$$

For  $\sigma < 0$ , put  $r_i = n_i k_i + j_i$ ,  $1 \leq j_i \leq k_i$ ,  $-\infty < n_i < \infty$ ,  $1 \leq i \leq m$ , where  $n$  is the summation index for  $Z(\frac{1}{2}m-s, Q^{-1}; h+j/k, -g)$ . The right-hand side of (5) then becomes

$$\begin{aligned}
 d^{(m-1)/2-s} (2\pi)^{2s-m/2} e^{-2\pi i(\sigma, h)} \frac{\Gamma(\frac{1}{2}m-s)}{\Gamma(s)} \\
 \cdot \sum_r' \prod_{j=1}^m \bar{\chi}(r_j) e^{-2\pi i(r, \sigma/k)} \{Q^{-1}(r/k + h)\}^{s-m/2}.
 \end{aligned}$$

(3) now easily follows.

If  $m = 1$ ,  $g = h = 0$ , and  $\chi(-1) = 1$ , we have derived the functional equation of Dirichlet's  $L$ -functions. A similar proof of the functional equation for Dirichlet  $L$ -functions originated with Hurwitz [4]. Slight variants of his proof can be found in [2, p. 73] and [1, pp. 232-233].

Define the character  $\chi'$  by  $\chi'(r) = \chi(r)\chi_1(r)^m$ , where  $\chi$  is a character (mod  $k$ ) and  $\chi_1$  is a character defined by

$$\begin{aligned}
 \chi_1(r) &= (k' | r), & k' &= (-1)^{(k-1)/2} k, & k &\text{ odd,} \\
 & & &= -k, & k &\equiv 0 \pmod{4}, \\
 & & &= 4k, & k &\equiv 2 \pmod{4},
 \end{aligned}$$

where  $(k' | r)$  denotes the Kronecker symbol.

**THEOREM 2.** *Let  $(d, k) = 1$  with either  $m$  even or  $k$  odd. Suppose  $\chi'$  is a primitive character (mod  $k$ ). Then  $L(s, \chi, Q; g, h)$  can be continued to a meromorphic function of  $s$  that is either entire or has but one pole, a simple one at  $s = m/2$ . Furthermore,*

$$\begin{aligned}
 (6) \quad \left(\frac{kd^{1/m}}{2\pi}\right)^\alpha \Gamma(s) L(s, \chi, Q; g, h) &= \alpha e^{-2\pi i(\sigma, h)} \left(\frac{kd^{(m-1)/m}}{2\pi}\right)^{m/2-s} \Gamma(\frac{1}{2}m-s) \\
 &\cdot L(\frac{1}{2}m-s, \bar{\chi}', Q^{-1}; hk, -g/k),
 \end{aligned}$$

where  $|\alpha| = 1$ .

For the value of  $\alpha$  see [5, p. 37]. If  $g = h = 0$ , Theorem 2 is due to Stark [5]. Stark also shows [5, Lemma 4] that  $Q^{-1}$ , indeed, has integral coefficients.

PROOF. We shall let  $\sum_{j=1}^k = \sum_{j_1=1}^k \cdots \sum_{j_m=1}^k$ . From [5, Theorem 1],

$$\sum_{j=1}^k \chi(Q(j)) e^{2\pi i(j,n)/k} = \alpha k^{m/2} \bar{\chi}'(Q^{-1}(n)).$$

Thus, for  $\sigma > m/2$ ,

$$\begin{aligned} \alpha k^{m/2} L(s, \bar{\chi}', Q^{-1}; g, h) &= \sum_{j=1}^k \chi(Q(j)) \sum_n' e^{2\pi i(n, h+j/k)} \{Q^{-1}(n + g)\}^{-s} \\ &= \sum_{j=1}^k \chi(Q(j)) Z(s, Q^{-1}; g, h + j/k). \end{aligned}$$

The first part of the theorem now follows as in the proof of Theorem 1. Now use (2) to obtain, for all  $s$ ,

$$\begin{aligned} \alpha k^{m/2} L(s, \bar{\chi}', Q^{-1}; g, h) &= d^{1/2-s} (2\pi)^{2s-m/2} \frac{\Gamma(\frac{1}{2}m - s)}{\Gamma(s)} \sum_{j=1}^k \chi(Q(j)) e^{-2\pi i(\sigma, h+j/k)} \\ &\quad \cdot Z(\frac{1}{2}m - s, Q; -(h + j/k), g). \end{aligned} \tag{7}$$

For  $\sigma < 0$ , put  $r_i = n_i k - j_i$ ,  $1 \leq j_i \leq k$ ,  $-\infty < n_i < \infty$ ,  $1 \leq i \leq m$ , where  $n$  is the summation index of  $Z(\frac{1}{2}m - s, Q; -(h + j/k), g)$ . Since  $\chi(Q(r)) = \chi(Q(j))$ , the right-hand side of (7) becomes

$$d^{1/2-s} (2\pi)^{2s-m/2} \frac{\Gamma(\frac{1}{2}m - s)}{\Gamma(s)} e^{-2\pi i(\sigma, h)} k^{m-2s} L(\frac{1}{2}m - s, \chi, Q; -hk, g/k).$$

If we now replace  $s$  by  $\frac{1}{2}m - s$ ,  $-hk$  by  $g$ , and  $g/k$  by  $h$ , we arrive at (6).

REFERENCES

1. Tom M. Apostol, *Dirichlet L-functions and character power sums*, J. Number Theory **2** (1970), 223-234.
2. Harold Davenport, *Multiplicative number theory*, Markham, Chicago, Ill., 1967. MR **36** #117.
3. Paul Epstein, *Zur Theorie allgemeiner Zetafunktionen*. II, Math. Ann. **63** (1907), 205-216.
4. Adolf Hurwitz, *Einige Eigenschaften der Dirichlet'schen Funktionen  $F(s) = \sum (D/n) \cdot 1/n^s$ , die bei der Bestimmung der Klassenanzahlen binärer quadratischer Formen auftreten*, Z. Math. Phys. **27** (1882), 86-101.
5. H. M. Stark, *L-functions and character sums for quadratic forms*. I, Acta Arith. **14** (1967/68), 35-50. MR **37** #2707.