

## ON TOEPLITZ OPERATORS WHICH ARE CONTRACTIONS

ROBERT GOOR

ABSTRACT. We prove that a Toeplitz contraction  $T_\phi$  is completely nonunitary if  $\phi$  is not a constant. As an application, it is noted that for such  $T_\phi$ , a functional calculus can be defined for all functions  $u$  in  $H^\infty$  of the unit disk.

For  $1 \leq p \leq \infty$ , we denote by  $L^p$  the usual class of Lebesgue measurable functions on the unit circle  $\gamma$  of the complex plane. We write  $H^p$  for the closed subspace of  $L^p$  of functions whose Fourier coefficients vanish on the negative integers. We denote by  $P$  the orthogonal projection of  $L^2$  onto  $H^2$  and by  $B(H^2)$  the space of bounded operators on  $H^2$ . For  $\phi \in L^\infty$ , we consider the Toeplitz operator  $T_\phi \in B(H^2)$  defined by  $T_\phi f = P(\phi f)$  for  $f \in H^2$ . Following Sz.-Nagy and Foiaş [2], we say a contraction  $T_\phi$ ,  $\|T_\phi\| \leq 1$ , is completely nonunitary (c.n.u.) if  $T_\phi$  has no nontrivial reducing subspaces restricted to which  $T_\phi$  is unitary. We will use the fact that  $T_\phi^* = T_{\bar{\phi}}$  [1, p. 137].

THEOREM. *If  $\phi \in L^\infty$ ,  $\|\phi\|_\infty \leq 1$  and  $\phi$  is not a constant (almost everywhere), then  $T_\phi$  is c.n.u.*

PROOF. Suppose  $T_\phi$  is not c.n.u. We will show that  $\phi$  is constant. Let  $S$  be a nontrivial reducing subspace for  $T_\phi$  on which  $T_\phi$  is unitary. We may write  $S = \{f \in H^2 : \|T_\phi^n f\|_2 = \|f\|_2 = \|T_{\bar{\phi}}^n f\|_2 \text{ for } n=1, 2, \dots\}$ . Now, for  $f \in S$ ,  $\|f\|_2 = \|T_\phi f\|_2 = \|P\phi f\|_2 \leq \|\phi f\|_2 \leq \|f\|_2$  and the resulting equality gives  $\phi f \in H^2$ . Similarly,  $\phi^n f$  and  $\bar{\phi}^n f \in H^2$  for  $n \geq 0$  and  $f \in S$ . We may apply the F. and M. Riesz theorem [1, p. 82] to the equality  $\|\phi f\|_2 = \|f\|_2$  for a nonzero  $f \in S$  to conclude that  $|\phi| = 1$  almost everywhere on  $\gamma$ . Thus, we write  $S = \{f \in H^2 : \phi^n f, \bar{\phi}^n f \in H^2 \text{ for } n \geq 0\}$ .

Let  $M_z$  be the operator of multiplication by the coordinate function  $z$ . Then  $\phi^n f \in H^2$  implies  $z\phi^n f = \phi^n z f \in H^2$  and similarly for  $\bar{\phi}^n z f$ , i.e.  $M_z S \subset S$ . By Beurling's theorem [1, p. 79], there is a function  $\psi \in H^\infty$ ,  $|\psi| = 1$  almost everywhere, such that  $S = \psi H^2$ . Since  $1 \in H^2$ ,  $\psi$  is in  $S$ . Note that  $\phi\psi = \psi f$  for some  $f \in H^2$  (since  $\phi\psi \in S = \psi H^2$ ). Hence for

---

Received by the editors November 15, 1971.

AMS 1970 subject classifications. Primary 47B35, 47A20, 47A60; Secondary 46J15.

Key words and phrases. Toeplitz operator, contraction, completely nonunitary, reducing subspace, Beurling theorem, F. and M. Riesz theorem.

© American Mathematical Society 1972

$n \geq 0$ , we have

$$\int_{\gamma} \phi z^n dz = \int_{\gamma} (\phi \psi)(\bar{\psi} z^n) dz = \int_{\gamma} (\psi f)(\bar{\psi} z^n) dz = \int_{\gamma} f z^n dz = 0$$

which implies that  $\phi \in H^{\infty}$ . Similarly,  $\bar{\phi} \in H^{\infty}$  implies  $\phi$  is a constant.

**COROLLARY.** *If  $\|\phi\|_{\infty} \leq 1$  and if  $\phi$  is not constant, then the map  $u \rightarrow u(T_{\phi})$  from  $H^{\infty}$  into  $B(H^2)$  defined by*

$$u(T_{\phi}) = \lim_{r \rightarrow 1-0} \sum_{k=0}^{\infty} C_k r^k T_{\phi}^k,$$

where  $u(\lambda) = \sum_{k=0}^{\infty} C_k \lambda^k \in H^{\infty}$  is a norm decreasing homomorphism of the algebra  $H^{\infty}$  into  $B(H^2)$ .

**PROOF.** Apply the above theorem and Theorem 2.1, Chapter III of [2].

#### REFERENCES

1. P. Halmos, *A Hilbert space problem book*, Van Nostrand, Princeton, N.J., 1967. MR 34 #8178.
2. B. Sz.-Nagy and C. Foiaş, *Analyse harmonique des opérateurs de l'espace de Hilbert*, Masson, Paris; Akad. Kiadó, Budapest, 1967. MR 37 #778.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48104