

A REPRESENTATION FORMULA FOR HARMONIC FUNCTIONS

CHIN-HUNG CHING AND CHARLES K. CHUI

ABSTRACT. We give a formula to reconstruct certain entire harmonic functions from their values on some lattice points.

1. Introduction and results. In [1], Boas proved the following uniqueness theorem for harmonic functions.

THEOREM A. *Let $u(z)$ be a real-valued entire harmonic function of exponential type less than π such that $u(m)=0$ and $u(m+i)=0$ for $m=0, \pm 1, \pm 2, \dots$. Then $u(z) \equiv 0$.*

He also asked if it is possible to reconstruct an entire harmonic function of exponential type less than π from its values on the lattice points $m, m+i$. We have

THEOREM 1. *Let $u(z)$ be a real-valued entire harmonic function of exponential type $\tau \leq \pi$ such that $u(x)$ and $u(x+i)$ are in $L^2(-\infty, \infty)$. Then*

$$\begin{aligned}
 (1) \quad u(z) &= u(x + iy) \\
 &= \sum_{n=-\infty}^{\infty} \frac{u(n)}{2\pi} \int_{-\pi}^{\pi} \frac{\sinh t(1-y)}{\sinh t} e^{it(x-n)} dt \\
 &\quad + \sum_{n=-\infty}^{\infty} \frac{u(n+i)}{2\pi} \int_{-\pi}^{\pi} \frac{\sinh ty}{\sinh t} e^{it(x-n)} dt \\
 &\quad + c_1 e^{\pi x} \sin \pi y + c_2 e^{-\pi x} \sin \pi y
 \end{aligned}$$

where the series converges uniformly in every strip $|y| \leq K < \infty$. Furthermore, if $\tau < \pi$ then $c_1 = c_2 = 0$, and if $\tau = \pi$ then

$$c_1 = \lim_{x \rightarrow \infty} \frac{e^{-\pi x} u(x + iy)}{\sin \pi y} \quad \text{and} \quad c_2 = \lim_{x \rightarrow \infty} \frac{e^{-\pi x} u(-x + iy)}{\sin \pi y}$$

for any $y, 0 < y < 1$.

Received by the editors September 20, 1972 and, in revised form, October 27, 1972.
 AMS (MOS) subject classifications (1970). Primary 31A10; Secondary 30A62.

Key words and phrases. Harmonic functions, entire functions of exponential type, Paley-Wiener theorem, Plancherel theorem, representation formula.

Hence, we have the following

COROLLARY. *Let $u(z)$ be a real-valued entire harmonic function of exponential type equal to π such that $u(x)$ and $u(x+i)$ are in $L^2(-\infty, \infty)$, $u(m)=0$ and $u(m+i)=0$ for $m=0, \pm 1, \pm 2, \dots$. Then*

$$u(z) = c_1 e^{\pi x} \sin \pi y + c_2 e^{-\pi x} \sin \pi y$$

for some real constants c_1 and c_2 .

Of course, the above results hold for complex-valued $u(z)$.

2. Proof of Theorem 1. Let $f(z)$ be an entire function with $\text{Re } f = u$ and let $F(z) = f(z) + [f(\bar{z})]^-$. Then $F(z)$ is an entire function of exponential type at most π . By the Paley-Wiener theorem, we have

$$F(z) = \int_{-\pi}^{\pi} e^{izt} \phi(t) dt.$$

Since $F(x) = 2u(x)$ is in $L^2(-\infty, \infty)$, $\phi(t)$ is in $L^2(-\pi, \pi)$ (cf. [2]). Hence,

$$\sum_{n=-\infty}^{\infty} u^2(n) = \frac{1}{4} \sum_{n=-\infty}^{\infty} F^2(n) = \frac{1}{4} \int_{-\pi}^{\pi} |\phi(t)|^2 dt < \infty.$$

Similarly, we have $\sum u^2(n+i) < \infty$. Using Schwarz's inequality and the Plancherel theorem, we obtain

$$\begin{aligned} & \left| \sum_{n=-\infty}^{\infty} \left| \frac{u(n)}{2\pi} \int_{-\pi}^{\pi} \frac{\sinh t(1-y)}{\sinh t} e^{it(x-n)} dt \right|^2 \right. \\ & \leq \left\{ \frac{1}{4} \pi^2 \sum_{n=-\infty}^{\infty} u^2(n) \right\} \left\{ \sum_{n=-\infty}^{\infty} \left| \int_{-\pi}^{\pi} \left\{ \frac{\sinh t(1-y)}{\sinh t} e^{itx} \right\} e^{-itn} dt \right|^2 \right\} \\ & \leq c_1 \int_{-\pi}^{\pi} \left| \frac{\sinh t(1-y)}{\sinh t} e^{itx} \right|^2 dt \\ & \leq c_2 \int_{-\pi}^{\pi} \left| \frac{\sinh t(1-y)}{t} \right|^2 dt \leq c_3 |1-y|^2 e^{2\pi|y|} \end{aligned}$$

for all $x+iy$. Similarly, we have

$$\sum_{n=-\infty}^{\infty} \left| \frac{u(n+i)}{2\pi} \int_{-\pi}^{\pi} \frac{\sinh ty}{\sinh t} e^{it(x-n)} dt \right| \leq c_4 |y| e^{\pi|y|}.$$

Hence, the series in (1) converges uniformly in every strip $|y| \leq K < \infty$ to a real-valued entire harmonic function $w(z)$ with

$$(2) \quad |w(x+iy)| \leq (\alpha + \beta |y|) e^{\pi|y|}$$

for some constants α, β and for all $x+iy$.

Since $F(z)$ is an entire function of exponential type $\leq \pi$ and its restriction to the real axis is in $L^2(-\infty, \infty)$, we have

$$\begin{aligned} u(x) &= \frac{1}{2} F(x) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \frac{\sin \pi(x-n)}{x-n} F(n) \\ &= \frac{1}{\pi} \sum_{n=-\infty}^{\infty} \frac{\sin \pi(x-n)}{x-n} u(n) \\ &= \sum_{n=-\infty}^{\infty} \frac{u(n)}{2\pi} \int_{-\pi}^{\pi} e^{it(x-n)} dt = w(x) \end{aligned}$$

for all real x . Similarly, we can also conclude that $u(x+i)=w(x+i)$.

Now, let $h(z)$ be an entire function with $\operatorname{Re} h = u - w$. From (2) and Carathéodory's inequality, we have

$$(3) \quad h(z) = O(e^{(\pi+\varepsilon)|z|})$$

for $0 < \varepsilon < 1$. Let $H(z) = h(z) + [h(\bar{z})]^-$. Then $H(x) = 2 \operatorname{Re} h(x) = 0$ for all real x so that $H(z) \equiv 0$ or

$$(4) \quad h(z) = -[h(\bar{z})]^-.$$

Similarly, let $G(z) = h(z+i) + [h(\bar{z}+i)]^-$. Then $G(x) = 2 \operatorname{Re} h(x+i) = 0$ for all real x so that $h(z+i) = -[h(\bar{z}+i)]^-$ for all z . Hence, by combining this with (4), we see that $h(z)$ has period $2i$. Now, it is well known that an entire function with period $2i$ satisfying (3) must be an exponential sum of the form $h(z) = \alpha + \beta e^{\pi z} + \gamma e^{-\pi z}$. That is, we have

$$\begin{aligned} u(x+iy) - w(x+iy) \\ = a + b_1 e^{\pi x} \cos \pi y + c_1 e^{\pi x} \sin \pi y + b_2 e^{-\pi x} \cos \pi y + c_2 e^{-\pi x} \sin \pi y \end{aligned}$$

for some real constants a, b_1, b_2, c_1, c_2 . Since $u(x) = w(x)$ for all real x , $a = b_1 = b_2 = 0$. From (2), we can conclude that

$$c_1 = \lim_{x \rightarrow \infty} e^{-\pi x} \frac{u(x+iy) - w(x+iy)}{\sin \pi y} = \lim_{x \rightarrow \infty} e^{-\pi x} \frac{u(x+iy)}{\sin \pi y}$$

for $0 < y < 1$. Similarly, for $0 < y < 1$,

$$c_2 = \lim_{x \rightarrow -\infty} e^{\pi x} \frac{u(x+iy)}{\sin \pi y}.$$

It is obvious that $c_1 = c_2 = 0$ if $\tau < \pi$.

REFERENCES

1. R. P. Boas, Jr., *A uniqueness theorem for harmonic functions*, J. Approximation Theory **5** (1972), 425–427.
2. ———, *Entire functions*, Academic Press, New York, 1954. MR **16**, 914.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MELBOURNE, PARKVILLE, AUSTRALIA

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION,
TEXAS 77843