

**A SHORT PROOF OF VAN DER WAERDEN'S THEOREM
 ON ARITHMETIC PROGRESSIONS**

R. L. GRAHAM AND B. L. ROTHSCHILD

ABSTRACT. A short proof is given for the classical theorem of van der Waerden which asserts that for any partition of the integers into a finite number of classes, some class contains arbitrarily long arithmetic progressions.

Let $[a, b]$ denote the set of integers x with $a \leq x \leq b$. We call $(x_1, \dots, x_m), (x'_1, \dots, x'_m) \in [0, l]^m$ *l-equivalent* if they agree up through their last occurrences of l . For any $l, m \geq 1$, consider the statement

For any r , there exists $N(l, m, r)$ so that for any function $C: [1, N(l, m, r)] \rightarrow [1, r]$, there exist positive a, d_1, \dots, d_m such that $C(a + \sum_{i=1}^m x_i d_i)$ is constant on each l -equivalence class of $[0, l]^m$.

FACT 1. $S(l, m)$ for some $m \geq 1 \Rightarrow S(l, m+1)$.

PROOF. For a fixed r , let $M = N(l, m, r)$, $M' = N(l, 1, r^M)$ and suppose $C: [1, MM'] \rightarrow [1, r]$ is given. Define $C': [1, M'] \rightarrow [1, r^M]$ so that $C'(k) = C'(k')$ iff $C(kM - j) = C(k'M - j)$ for all $0 \leq j < M$. By the inductive hypothesis, there exist a' and d' such that $C'(a' + xd')$ is constant for $x \in [0, l-1]$. Since $S(l, m)$ can apply to the interval $[a'M+1, (a'+1)M]$, then by the choice of M , there exist a, d_1, \dots, d_m with all sums $a + \sum_{i=1}^m x_i d_i$, $x_i \in [0, l]$, in $[a'M+1, (a'+1)M]$ and with $C(a + \sum_{i=1}^m x_i d_i)$ constant on l -equivalence classes. Set $d'_i = d_i$ for $i \in [1, m]$ and $d'_{m+1} = d'M$; then $S(l, m+1)$ holds.

FACT 2. $S(l, m)$ for all $m \geq 1 \Rightarrow S(l+1, 1)$.

PROOF. For a fixed r , let $C: [1, 2N(l, r, r)] \rightarrow [1, r]$ be given. Then there exist a, d_1, \dots, d_r such that for $x_i \in [0, l]$, $a + \sum_{i=1}^r x_i d_i \leq N(l, r, r)$ and $C(a + \sum_{i=1}^r x_i d_i)$ is constant on l -equivalence classes. By the box principle there exist $u < v$ in $[0, r]$ such that

$$C\left(a + \sum_{i=1}^u l d_i\right) = C\left(a + \sum_{i=1}^v l d_i\right).$$

Received by the editors February 1, 1973.

AMS (MOS) subject classifications (1970). Primary 05A99; Secondary 10L99.

Key words and phrases. van der Waerden's theorem, long arithmetic progressions.

Therefore $C((a + \sum_{i=1}^u ld_i) + x(\sum_{i=u+1}^v d_i))$ is constant for $x \in [0, 1]$. This proves $S(l+1, 1)$.

Since $S(1, 1)$ holds trivially, then by induction $S(l, m)$ is valid for all $l, m \geq 1$. Van der Waerden's theorem is $S(l, 1)$.

The authors point out that while previous proofs follow essentially the argument above, the one given is hopefully clearer.

REFERENCES

1. R. L. Graham and B. L. Rothschild, *Ramsey's theorem for n -parameter sets*, Trans. Amer. Math. Soc. **159** (1971), 257-292. MR **44** #1580.
2. A. W. Hales and R. I. Jewett, *Regularity and positional games*, Trans. Amer. Math. Soc. **106** (1963), 222-229. MR **26** #1265.
3. A. Ja. Hinčin, *Three pearls of number theory*, Graylock Press, Rochester, N.Y., 1952. MR **13**, 724.
4. R. Rado, *Studien zur Kombinatorik*, Math. Z. **36** (1933), 424-480.
5. B. L. van der Waerden, *Beweis einer Baudetschen Vermutung*, Nieuw Arch. Wisk. **15** (1927), 212-216.

BELL LABORATORIES, MURRAY HILL, NEW JERSEY 07974

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CALIFORNIA 90024