

ON THE EXISTENCE OF CONTACT FORMS

W. P. THURSTON AND H. E. WINKELNKEMPER

ABSTRACT. Using an old theorem of Alexander, we give a short and elementary proof that every closed, orientable 3-manifold has a contact form.

Introduction. Let M^m ($m = 2n + 1$) be a closed, smooth, orientable m -manifold; a contact form on M is a smooth 1-form ω such that $\omega \wedge (d\omega)^n \neq 0$ at each point. One knows [2] that the existence of a contact form on M^m allows a reduction of the structure group of the tangent bundle of M to $U(n)$ and hence, in particular, the odd-dimensional Stiefel-Whitney classes of M have to vanish [4]. This, however, gives no information in dimension 3 and in [3] Chern¹ asked: Does every closed, smooth, orientable 3-manifold admit a smooth contact form?

This question was first answered (in the affirmative) by Lutz [6] and Martinet [7]. They based their proof on the theorem that every closed, orientable 3-manifold can be obtained from S^3 by surgery, the same theorem used by Lickorish, Novikov and Zieschang to prove that every such manifold admits a codimension 1 foliation, i.e. a nonzero 1-form η on M such that $\eta \wedge d\eta \equiv 0$; that is, in dimension 3, the exact opposite of a contact form.

It was observed by one of us (see also [5]) that the existence theorem for codimension 1 foliations on M is an immediate consequence of the following

Theorem (Alexander [1]). *Every closed, smooth, orientable 3-manifold M^3 is diffeomorphic to $W(h) \cup_{\text{id}} \partial W \times D^2$, where D^2 is the 2-disc and W is an orientable 2-manifold with boundary and $h: W \rightarrow W$ is a diffeomorphism which restricts to the identity on ∂W ; $W(h)$ denotes the mapping torus of h , i.e. the 3-manifold with boundary obtained from $W \times I$ by identifying $(x, 0)$ with $(h(x), 1)$, $I = [0, 1]$.*

The object of this short note is to prove that the existence theorem for contact forms on 3-manifolds is also a more or less immediate and elementary consequence of Alexander's theorem.

Received by the editors July 23, 1974.

AMS (MOS) subject classifications (1970). Primary 53C15; Secondary 58A10, 58A30.

¹Also in Proc. U. S.-Japan Seminar in Differential Geometry (Kyoto, 1965), Nippon Hyoronsha, Tokyo, 1966; Question 1, p. 172. MR 35 #7268.

Copyright © 1975, American Mathematical Society

Construction of a contact form. We may assume that h fixes a neighborhood of ∂W . Let $d\theta$ be a volume form for ∂W , and let t be the collar parameter for a collar neighborhood C of ∂W , in W , invariant by h .

Let α_1 be any 1-form which equals $(1+t)d\theta$ near ∂W . By Stokes' theorem, $\int_W d\alpha_1 = 1$. If Ω is any volume form for W which is $dt \wedge d\theta$ near ∂W , then, by de Rham's theorem, $\Omega - d\alpha_1 = d\beta$, where β is a 1-form which is zero near ∂W . Let $\alpha_2 = \alpha_1 + \beta$. Then α_2 satisfies (i) $d\alpha_2$ is a volume form, and (ii) $\alpha_2 = (1+t)d\theta$ near ∂W .

The set of 1-forms satisfying (i) and (ii) is convex. Hence, there is a 1-form α on $W(h)$, such that α restricted to any fiber, satisfies (i) and (ii). Since $d\theta$ and t are defined in a natural way on $C \times S^1$, we may also specify that $\alpha = (1+t)d\theta$ in a neighborhood of $\partial W(h)$.

For a sufficient large constant K , it follows that $\omega = \alpha + Kd\phi$ is a contact form on $W(h)$, where $d\phi$ is the pullback of a volume form for the base of the bundle $W(h)$ over S^1 . In fact, $d\phi \wedge d\alpha$ is a nonsingular 3-form on $W(h)$, so $\omega \wedge d\omega = K(d\phi \wedge d\alpha) + \alpha \wedge d\alpha$ will also be nonsingular when K is large.

We will now extend ω over $\partial W(h) \times D^2$. We will use coordinates (θ, r, ϕ) , where (r, ϕ) are polar coordinates for D^2 . We may consider $C \times S^1$ to be $\partial W(h) \times (D_2^2 - D^2)$, where D_2^2 is the disk of radius two, and the present coordinates are related to the old coordinates by $\phi = \phi$, $\theta = \theta$, $r = 1 + t$, so in the new coordinates, $\omega = rd\theta + Kd\phi$ in $\partial W(h) \times (D_2^2 - D^2)$. The form $-d\theta + r^2d\phi$ is a smooth contact form near $r = 0$: this may be checked by converting to cartesian coordinates for D^2 . Consider a 1-form ω which may be written $\omega = f_1(r)d\theta + f_2(r)d\phi$. ω is a contact form away from $r = 0$, iff

$$\omega \wedge d\omega = \begin{vmatrix} f_1(r) & f_1'(r) \\ f_2(r) & f_2'(r) \end{vmatrix} d\theta \wedge dr \wedge d\phi \neq 0.$$

That is, ω is a contact form iff the position vector and the tangent vector of the curve $(f_1(r), f_2(r))$ in \mathbb{R}^2 are linearly independent, for $r \neq 0$. It is now clear that there is a curve $(f_1(r), f_2(r))$ [$0 < r \leq 2$] such that $(f_1(r), f_2(r)) = (-1, r^2)$ near $r = 0$, $(f_1(r), f_2(r)) = (r, K)$ for $1 \leq r \leq 2$, and such that $\omega = f_1(r)d\theta + f_2(r)d\phi$ is a contact form.

This completes the construction.

REFERENCES

1. J. W. Alexander, *A lemma on systems of knotted curves*, Proc. Nat. Acad. Sci. U.S.A. 9 (1923), 93-95.
2. S. S. Chern, *Pseudo-groupes continus infinies*, Géométrie Différentielle (Colloq. Internat. Centre National de la Recherche Scientifique, Strasbourg, 1953), Centre National de la Recherche Scientifique, Paris, 1953, pp. 119-136. MR 16, 112.
3. S. S. Chern, *The geometry of G-structures*, Bull. Amer. Math. Soc. 72 (1966), 167-219. MR 33 #661.

4. J. W. Gray, *Some global properties of contact structures*, *Ann. of Math.* (2) **69** (1959), 421–450. MR 22 #3016.
5. H. B. Lawson, Jr., *Foliations*, *Bull. Amer. Math. Soc.* **80** (1974), 369–418.
6. R. Lutz, *Sur quelques propriétés des formes différentielles en dimension trois*, Thèse, Strasbourg, 1971.
7. J. Martinet, *Formes de contact sur les variétés de dimension 3*, *Proc. Liverpool Singularities Sympos. II*, *Lecture Notes in Math.*, vol. 209, Springer-Verlag, Berlin and New York, 1971, 142–163.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08540

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MARYLAND 20742