

## A NAGUMO CONDITION FOR ORDINARY DIFFERENTIAL EQUATIONS<sup>1</sup>

L. K. JACKSON

**ABSTRACT.** A condition is given which guarantees that each solution of  $y^{(n)} = f(x, y, y', \dots, y^{(n-1)})$  either extends or becomes unbounded on its maximal interval of existence.

**Introduction.** The classical Nagumo condition for the second order ordinary differential equation

$$(1) \quad y'' = f(x, y, y')$$

is a growth condition on  $f(x, y, y')$  which implies that solutions of (1) either extend or become unbounded on their maximal intervals of existence. One formulation of the condition is contained in the following theorem.

**THEOREM 1.** *Assume that (1) is a scalar equation with  $f(x, y, y')$  continuous on  $(a, b) \times R^2$ . If for each  $M > 0$  and each compact interval  $[c, d] \subset (a, b)$  there is a corresponding positive continuous function  $\phi(s)$  on  $[0, \infty)$  such that  $|f(x, y, y')| \leq \phi(|y'|)$  for all  $(x, y, y')$  satisfying  $c \leq x \leq d$ ,  $|y| \leq M$  and such that  $\int_0^\infty s/\phi(s)ds = +\infty$ , then each solution of (1) either extends to  $(a, b)$  or becomes unbounded on its maximal interval of existence.*

Other formulations from which Theorem 1 follows may be found in [1, p. 428] and [2, p. 353].

This property of solutions which is stated as the conclusion in Theorem 1 along with the assumed existence of solutions of certain types of differential inequalities plays an important role in demonstrating the existence of solutions of boundary value problems not only for second order equations but for higher order equations as well; see for example, [2]–[6].

In this paper we establish a “Nagumo condition” for equations

$$(2) \quad y^{(n)} = f(x, y, y', \dots, y^{(n-1)})$$

of arbitrary order  $n$ .

**1. Preliminary results.** For a function  $g \in C^{(n)}[\alpha, \beta]$  let  $M_k = \text{Max} \{|g^{(k)}(x)|: \alpha \leq x \leq \beta\}$ ,  $0 \leq k \leq n$ . The proof of the following lemma may be found in [7].

---

Received by the editors June 18, 1975.

*AMS (MOS) subject classifications* (1970). Primary 34C10; Secondary 34B15.

*Key words and phrases.* Nagumo condition, maximal interval of existence, boundary value problem, differential inequalities.

<sup>1</sup> This work was partially supported by NSF Grant MPS75-06174.

© American Mathematical Society 1976

LEMMA 2. Let  $M_k, 0 \leq k \leq n$ , be as above. Then for each integer  $k, 0 < k < n$ ,

$$M_k \leq C_{nk} M_0^{1-k/n} (M'_n)^{k/n}$$

where

$$C_{nk} = 4e^{2k} n^k k^{-k} \quad \text{and} \quad M'_n = \text{Max}\{M_n, 2^n n! M_0(\beta - \alpha)^{-n}\}.$$

Note. The factor  $2^n$  appearing in the definition of  $M'_n$  is not present in the result as stated in [7], however, the author believes that it should be included.

DEFINITION 3. For  $p \geq 0$  let the function  $\phi_p(x)$  be defined by

$$\phi_p(x) = \begin{cases} 1 & \text{for } |x| \leq 1, \\ |x|^p & \text{for } |x| \geq 1. \end{cases}$$

LEMMA 4. Assume that the numbers  $p_i, 1 \leq i \leq n - 1$ , are fixed nonnegative numbers such that  $\sum_{i=1}^{n-1} ip_i < n$ . Then

$$\sum_{i=1}^{n-1} |x_i|^{n/i} / \prod_{i=1}^{n-1} \phi_{p_i}(x_i) \rightarrow +\infty \quad \text{as} \quad \sum_{i=1}^{n-1} |x_i|^{n/i} \rightarrow +\infty.$$

PROOF. Assume that  $\sum_{i=1}^{n-1} |x_i|^{n/i} = h > 1$ . Then  $|x_i| \leq h^{i/n}$  for each  $1 \leq i \leq n - 1$  from which it follows that

$$\phi_{p_i}(x_i) \leq h^{ip_i/n} \quad \text{and} \quad \prod_{i=1}^{n-1} \phi_{p_i}(x_i) \leq h^{(1/n)\sum_{i=1}^{n-1} ip_i}.$$

Thus  $\sum_{i=1}^{n-1} |x_i|^{n/i} = h > 1$  implies that

$$\sum_{i=1}^{n-1} |x_i|^{n/i} / \prod_{i=1}^{n-1} \phi_{p_i}(x_i) \geq h^{1-1/n\sum_{i=1}^{n-1} ip_i}$$

and the result follows from the assumption that  $\sum_{i=1}^{n-1} ip_i < n$ .

**2. The Nagumo condition.** In this section we assume that we are dealing with a fixed equation (2) of degree  $n \geq 3$  and we assume that in (2) the function  $f(x, y, y', \dots, y^{(n-1)})$  is continuous on  $I \times R^n$  where  $I$  is an interval of the reals.

THEOREM 5. Assume that the function  $f(x, y, y', \dots, y^{(n-1)})$  in (2) satisfies the condition that given any compact interval  $[c, d] \subset I$  and given any  $M > 0$  there exist numbers  $h, p_i, 1 \leq i \leq n - 1$ , such that  $h > 0, p_i \geq 0$  for  $1 \leq i \leq n - 1, \sum_{i=1}^{n-1} ip_i < n$ , and such that

$$(3) \quad |f(x, y, y', \dots, y^{(n-1)})| \leq h \prod_{i=1}^{n-1} \phi_{p_i}(y^{(i)})$$

for all  $(x, y, y', \dots, y^{(n-1)})$  satisfying  $c \leq x \leq d, |y| \leq M$  where  $\phi_p(t)$  is as given in Definition 3. Then each solution of (2) either extends to  $I$  or becomes unbounded on its maximal interval of existence.

PROOF. We prove more than the actual statement of the theorem. We prove that a solution either extends to the right or becomes unbounded on its right

maximal interval of existence and that the same result holds for left extensions. Since the proofs are the same we consider only extensions to the right.

Assume that the conditions of the theorem are satisfied but that the conclusion is false. Then there is a compact interval  $[c, d] \subset I$  and a solution  $y(x)$  of (2) on  $[c, d]$  such that  $|y(x)|$  is bounded on  $[c, d]$  and such that  $[c, d]$  is a right maximal interval of existence for  $y(x)$ . Let  $|y(x)| \leq M_0$  on  $[c, d]$  and let the numbers  $h$  and  $p_i$ ,  $1 \leq i \leq n - 1$ , be associated with  $M_0 [c, d]$  as in the theorem.

Since  $[c, d]$  is right maximal for  $y(x)$  it follows that neither  $|y^{(n)}(x)|$  nor  $|y^{(n-1)}(x)|$  are bounded on  $[c, d]$ . For  $0 \leq k \leq n$  set

$$M_k(t) = \text{Max}\{|y^{(k)}(x)|: c \leq x \leq t < d\}.$$

Let  $t_0$  be chosen so that  $(c + d)/2 < t_0 < d$  and

$$M_n(t_0) \geq 2^n n! M_0((d - c)/2)^{-n}.$$

Then for any  $t_0 \leq t < d$ ,

$$M_n(t) \geq 2^n n! M_0((d - c)/2)^{-n} \geq 2^n n! M_0(t)(t - c)^{-n}$$

and it follows that for any  $t_0 \leq t < d$

$$M'_n(t) = \text{Max}\{M_n(t), 2^n n! M_0(t)(t - c)^{-n}\} = M_n(t).$$

It follows from Lemma 2 that for  $t_0 \leq t < d$  and for each integer  $k$ ,  $0 < k < n$ ,

$$\begin{aligned} M_k(t) &\leq C_{nk} [M_0(t)]^{1-k/n} [M_n(t)]^{k/n} \\ &\leq C_{nk} M_0^{1-k/n} [M_n(t)]^{k/n} = C_{nk}^* [M_n(t)]^{k/n}. \end{aligned}$$

This implies that

$$\sum_{k=1}^{n-1} [M_k(t)]^{n/k} \leq \left( \sum_{k=1}^{n-1} (C_{nk}^*)^{n/k} \right) M_n(t) = A_n M_n(t)$$

for  $t_0 \leq t < d$ . However, it follows from inequality (3) that

$$M_n(t) \leq h \prod_{k=1}^{n-1} \phi_{p_k}(M_k(t))$$

for all  $t_0 \leq t < d$ . Hence,

$$\sum_{k=1}^{n-1} [M_k(t)]^{n/k} \leq h A_n \prod_{k=1}^{n-1} \phi_{p_k}(M_k(t))$$

for  $t_0 \leq t < d$ . This contradicts Lemma 4 since  $M_{n-1}(t) \rightarrow +\infty$  as  $t \rightarrow d$  and therefore  $\sum_{k=1}^{n-1} [M_k(t)]^{n/k} \rightarrow +\infty$  as  $t \rightarrow d$ .

If the nonnegative numbers  $p_i$ ,  $1 \leq i \leq n - 1$ , are such that  $\sum_{i=1}^{n-1} i p_i > n$ , an  $r$  with  $0 < r < 1$  can be chosen so that  $y(x) = -(x_0 - x)^r$  is a solution of

$$(4) \quad y^{(n)} = h \prod_{i=1}^{n-1} |y^{(i)}|^{p_i}$$

on  $(-\infty, x_0)$  provided a suitable value is assigned to the constant  $h$ . In fact, substituting  $y(x) = -(x_0 - x)^r$  in (4), one can verify that  $y(x)$  is a solution on  $(-\infty, x_0)$  for a suitable choice of the constant  $h$  provided

$$r = \frac{\sum_{i=1}^{n-1} ip_i - n}{\sum_{i=1}^{n-1} p_i} - 1.$$

Then, using simple induction arguments, one can show that for  $n \geq 3$  and for nonnegative numbers  $p_i, 1 \leq i \leq n - 1, \sum_{i=1}^{n-1} ip_i \geq n$  implies  $\sum_{i=1}^{n-1} p_i > 1$  and  $\sum_{i=1}^{n-1} ip_i = n$  implies  $\sum_{i=2}^{n-1} (i - 1)p_i < n - 1$ . Thus, for each  $n \geq 3$ , nonnegative numbers  $p_i, 1 \leq i \leq n - 1$ , can be chosen so that  $0 < r < 1$ . It follows that the condition  $\sum_{i=1}^{n-1} ip_i < n$  in Theorem 5 could at best possibly be improved to  $\sum_{i=1}^{n-1} ip_i \leq n$ .

Essentially the same arguments as were used in proving Theorem 5 can be used to prove the following more general theorem.

**THEOREM 6.** *Assume that  $f(x, y, y', \dots, y^{(n-1)})$  satisfies the condition that given any compact interval  $[c, d] \subset I$  and given any  $M > 0$  there is a function  $\phi(t_1, t_2, \dots, t_{n-1})$  defined for  $t_i \geq 0, 1 \leq i \leq n - 1$ , such that*

$$|f(x, y, y', \dots, y^{(n-1)})| \leq \phi(|y'|, |y''|, \dots, |y^{(n-1)}|)$$

for all  $(x, y, y', \dots, y^{(n-1)})$  satisfying  $c \leq x \leq d, |y| \leq M$ , such that  $\phi(t_1, t_2, \dots, t_{n-1})$  is nondecreasing in each variable, and such that

$$\sum_{i=1}^{n-1} t_i^{n/i} / \phi(t_1, \dots, t_{n-1}) \rightarrow +\infty \text{ as } \sum_{i=1}^{n-1} t_i^{n/i} \rightarrow +\infty.$$

Then each solution of (2) either extends to  $I$  or becomes unbounded on its maximal interval of existence. In practice Theorem 5 is probably easier to apply.

Results similar to Theorem 6 for equations of orders 3 and 4 were given in [6].

### REFERENCES

1. P. Hartman, *Ordinary differential equations*, Wiley, New York and London, 1964. MR 30 # 1270.
2. L. K. Jackson, *Subfunctions and second-order differential inequalities*, *Advances in Math.* **2** (1968), 307-363. MR 37 # 5462.
3. K. W. Schrader, *Existence theorems for second order boundary value problems*, *J. Differential Equations* **5** (1969), 572-584. MR 39 # 532.
4. G. A. Klaasen, *Differential inequalities and existence theorems for second and third order boundary value problems*, *J. Differential Equations* **10** (1971), 529-537. MR 44 # 5595.
5. W. G. Kelley, *Some existence theorems for nth-order boundary value problems*, *J. Differential Equations* **18** (1975), 158-169.
6. J. Bebernes, R. Gaines and K. Schmitt, *Existence of periodic solutions for third and fourth order ordinary differential equations via coincidence degree*, *Ann. Soc. Sci. Bruxelles Sér. I* **88** (1974), 25-36. MR 49 # 5480.
7. A. Gorny, *Contribution à l'étude des fonctions dérivables d'une variable réelle*, *Acta Math.* **71** (1939), 317-358. MR 1, 137; 400.