

ON THE ZEROS OF GENERALIZED AXIALLY SYMMETRIC POTENTIALS*

PETER A. MCCOY

ABSTRACT. Generalized axially symmetric potentials may be expanded as Fourier-Jacobi series in terms of the complete system $r^k C_k^{n/2-1}(\cos \theta)$ on axisymmetric regions $\Omega \subset E^n$ ($n \geq 3$) about the origin. The values of these potentials are characterized by the nonnegativity of sequences of determinants drawn from the Fourier coefficients in a manner analogous to the characterization of the values of analytic functions of one complex variable by the theorems of Carathéodory-Toeplitz and Schur.

1. Introduction. The singularities of generalized axially symmetric potentials (GASP) ψ are characterized from the coefficients of their Fourier-Jacobi series expansions by S. Bergman [1], [2] and R. P. Gilbert [3], [4] who found a definitive characterization by employing function theoretic means to develop a theory mirroring the classical work of Hadamard, Mandelbrojt and Fabry on the singularities of analytic functions f of one complex variable. Cauchy's classical estimate [5, p. 123] on the zeros of a polynomial, along with the theorems of Carathéodory-Toeplitz [12, p. 157] and Schur [12, p. 159] bound the zeros of f relative to convex sets from its coefficients; the possibility arises of developing a parallel theory for the values of ψ .

An extension of Cauchy's estimate applying to the zeros of harmonic polynomials in E^n was found by Morris Marden [8]. We shall illustrate how the reasoning in [8] which is based on methods found in Professor Marden's early work [6], [7] can be used to locate the zeros of GASP. Moreover, it is apparent that this reasoning plays a role in bounding the zeros of ψ analogous to the role played by the reasoning of Hadamard in the multiplication of singularities theorem upon which Gilbert's "envelope method" [1], [2] is based. *Remark:* Investigations [9]–[11] subsequent to [8] employ the A_n operator [4, p. 168] to geometrically establish equality of known sets of excluded values of the A_n associates and corresponding sets of excluded values of GASP.

2. On the zeros. In axisymmetric regions $\Omega \subset E^n$ [8] about the origin, generalized axially symmetric potentials ψ_n may be developed as a series [4, p. 168]

Received by the editors September 28, 1975 and, in revised form, December 22, 1975.

AMS (MOS) subject classifications (1970). Primary 35B05, 35C10; Secondary 30A08.

Key words and phrases. Bergman and Gilbert's integral operators, Gilbert-Hadamard theorem, Carathéodory-Toeplitz and Schur theorems, zeros of potentials.

*This paper is dedicated to Professor Morris Marden on the occasion of his retirement from the University of Wisconsin-Milwaukee.

Copyright © 1977, American Mathematical Society

$$(1) \quad \psi_n(r, \theta) = \psi_n(\mathbf{X}) = \sum_{k=0}^{\infty} B(n-2, k+1) a_k r^k C_k^{n-1/2}(\cos \theta)$$

in terms of the polar coordinates (r, θ) which depend on the cylindrical coordinates $x = x_1, \rho = (x_2^2 + \dots + x_n^2)^{1/2}$ by $x = r \cos \theta, \rho = r \sin \theta$. Because of symmetry, ψ_n assumes constant values of hypercircles $\mathbf{X} = (x, \rho)$ in Ω .

For each ψ_n with domain Ω , there is a unique associated function f ,

$$(2) \quad f(\xi) = \sum_{k=0}^{\infty} a_k \xi^k,$$

analytic on the corresponding axiconvex domain $\omega \subset \mathbf{C}$ [8] whose A_n transform [4] is defined by

$$(3) \quad \psi_n(\mathbf{X}) = \alpha_n \int_L f(\sigma) dv_n(\zeta),$$

$\sigma = x + \rho/2(\zeta + \zeta^{-1}), \alpha_n = \Gamma(n-2)/(4i)^{n-2}\Gamma(n/2-1)$ with nonnegative measure

$$(4) \quad dv_n(\zeta) = (\zeta - \zeta^{-1})^{n-3} d\zeta/\zeta$$

for ζ on the contour $L \equiv \{\zeta = e^{it} | 0 \leq t \leq \pi\}$. (The Fourier-Legendre expansion for $n = 3$ follows from the Bergman-Whittaker operator.) From the development of ψ_n in (1) we defined the matrix

$$(5) \quad M_k = \begin{pmatrix} b_0, a_1, \dots, a_k \\ 0, b_0, \dots, a_{k-1} \\ \dots \dots \dots \\ 0, 0, 0, \dots, b_0 \end{pmatrix}$$

its conjugate transpose $\overline{M}'_k (b_0 \equiv a_0 - \alpha, \alpha \in \mathbf{C})$ and the determinants

$$(6) \quad \sigma_k(\alpha) = \sigma_k(a_0, \dots, a_k, \alpha) = \det(M_k + \overline{M}'_k)$$

and

$$(7) \quad \gamma_k(\alpha) = \gamma_k(a_0, \dots, a_k, \alpha) = \det \begin{pmatrix} M_k & I_k \\ I_k & \overline{M}'_k \end{pmatrix}$$

where I_k is the $(k+1)$ by $(k+1)$ identity matrix. Initially the values of ψ_n relative to the hypersphere S_n with unit radius about the origin are considered in

THEOREM 1. *Let the generalized axially symmetric potential ψ_n be expanded as in (1). Then ψ_n is regular in the hypersphere S_n and on each hypercircle \mathbf{X} in S_n ,*

$$(8) \quad \psi_n(\mathbf{X}) \neq \alpha + \eta, \quad \forall \operatorname{Re}(\eta) < 0, (a_0 - \alpha \text{ real})$$

or

$$(9) \quad \psi_n(\mathbf{X}) \neq \eta, \quad \forall |\eta - \alpha| > 1$$

for η complex according to whether

$$(8a) \quad \sigma_0(\alpha) > 0, \sigma_1(\alpha) > 0, \dots, \sigma_k(\alpha) > 0, \dots$$

or

$$(9a) \quad \gamma_0(\alpha) > 0, \gamma_1(\alpha) > 0, \dots, \gamma_k(\alpha) > 0, \dots$$

In particular, if $\alpha > 0$ and (8a) or $|\eta| > 1$ and (9a), then ψ_n has no zeros in S_n .

PROOF. When (8a) or (9a) are valid, the associate f given by (2) is regular in the disk $D = \{|\xi| < 1\}$ so that its A_n transform represents ψ_n in S_n . When (8a) is satisfied, the Carathéodory-Toeplitz theorem guarantees that $\operatorname{Re}[f(\sigma) - \alpha] \geq 0$ when $|\sigma| < 1$ which is the case for all $x^2 + \rho^2 < 1$ and $\zeta \in L$. Moreover, for $\zeta \in L$, the measure $d\nu_n(\zeta)$ is nonnegative. Consequently, the integral $\psi_n - \alpha = A_n(f - \alpha)$, viewed as the uniform limit of a sum of vectors, each terminating in the right half plane P , is in P (see [6], [7]). This establishes (8).

In (9a), Schur's theorem guarantees the bound $|f(\sigma) - \alpha| \leq 1$, $x^2 + \rho^2 < 1$, $\zeta \in L$. The measure is nonnegative and the operator is normalized according to $A_n(1) = 1$. Hence $|\psi_n(\mathbf{X}) - \alpha| \leq A_n(1)$. Remark: Due to Fatou's theorem [12, p. 146], $\alpha > 0$ is a sufficient condition for the existence of the radial limits of ψ_n relative to S_n , with the possible exception of a set of singular circles whose intersection with a meridian plane has one dimensional measure zero.

THEOREM 2. If for some constant α , the sequence

$$(10) \quad \gamma_0(\alpha) > 0, \dots, \gamma_k(\alpha) > 0, \quad \gamma_{k+1}(\alpha) = \dots = 0,$$

then ψ_n is a linear combination of Newtonian potentials whose mass is distributed over $k + 1$ singular circles exterior to S_n such that ψ_n has no zeros in the double cone

$$(11) \quad 0 < \rho < \pm(R - x)\tan(\pi/4(k + 1)).$$

The distance R from the origin to the most remote (finite) singular circle is determined by the Gilbert-Hadamard theorem. Moreover, if ψ_n vanishes on the sphere centre at $x = x_0$ and radius $\rho = \rho_0$ with no singular circles in the concentric sphere of radius $\rho_0 \csc(\pi/2(k + 1))$ about S_n , the associate vanishes in the disk $|\xi - x_0| \leq \rho_0 \csc(\pi/2(k + 1))$.

PROOF. Because of (10), the associate f is a rational function with $(k + 1)$ distinct zeros and $(k + 1)$ poles which are symmetric in $|\xi| = 1$ [12, p. 159]. Application of the A_n operator to the partial fraction expansion of f establishes the first part. To verify that ψ_n has no zeros in the cones (11), we reason as in [8] bearing in mind the location of the zeros and poles of the associate. To establish the converse relation on the effect of the zeros of ψ on those of f , apply Marden's mean value theorem [5, p. 11] in a manner similar to its use in [9].

Relating the values of ψ_n on axisymmetric sets to the values assumed in convex sets suggests a lengthy statement on the Fourier coefficients. For brevity, we consider the version in

THEOREM 3. The generalized axially symmetric potential ψ_n as in (1) has the axisymmetric set $\Omega \subset E^n$ for its domain where the analytic function

$$(12) \quad g(\xi) = \sum_{k=0}^{\infty} g_k \xi^k / k!$$

maps the unit disk onto the corresponding axiconvex set $\omega \subset \mathbf{C}$. The constants β_k are defined by

$$(13) \quad \beta_k = \sum_{j=0}^k a_j! / k! \sum P_k^{(j)} g_1^{m_1} \dots g_k^{m_k},$$

$$(14) \quad P_k^{(j)} = k! / (1!)^{m_1} \dots (k!)^{m_k} (m_k!).$$

The summations are over $m_1 + 2m_2 + \dots + km_k = k$, and $m_1 + \dots + m_k = j$. If either sequence,

$$(15) \quad \sigma_0(\beta_0, \alpha) > 0, \dots, \sigma_k(\beta_0, \dots, \beta_k, \alpha) > 0, \dots$$

or

$$(16) \quad \gamma_0(\beta_0, \alpha) > 0, \dots, \gamma_k(\beta_0, \dots, \beta_k, \alpha) > 0, \dots,$$

then on each hypercircle $\mathbf{X} \in \Omega$, ψ_n satisfies the corresponding inequality (8) or (9).

PROOF. We write $\tilde{f} = f \circ g$. Then (15) or (16) imply that f maps D onto the respective convex sets D or $\text{Re}(\xi) \geq 0$. Consequently, $f: \omega \rightarrow D$ or $\text{Re}(\xi) \geq 0$ so that reasoning as in Theorem 1 completes the proof.

3. **A generalization to $\Delta_3 \Phi + F(r^2) = 0$.** The purpose of this closing remark is to indicate a broader range of application of [6], [7] by referring to work of S. Bergman [2] who expands axially symmetric solutions of

$$(17) \quad L\Phi \equiv \frac{\partial^2 \Phi}{\partial x^2} + \frac{\partial^2 \Phi}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial \Phi}{\partial \rho} + F(r^2)\Phi = 0$$

(F is entire) in series of generalized Bessel functions J_n^* [2, p. 435].

THEOREM 4. If

$$(18) \quad \Phi(\mathbf{X}) = \sum_{k=0}^{\infty} \frac{\Gamma(1/2)A_k}{\Gamma(k + 3/2)} \frac{J_k^*(r)}{J_k^*(1)} r^k P_k(\cos \theta)$$

is the development of a solution of $L\Phi \equiv 0$ regular in the unit sphere S_3 and there is a constant $|\alpha| > 1$ for which

$$(19) \quad \gamma_0(A_0, \alpha) > 0, \gamma_1(A_0, A_1, \alpha) > 0, \dots, \gamma_k(A_0, \dots, A_k, \alpha) > 0,$$

then on each circle $\mathbf{X} \in S_3$,

$$(20) \quad \Phi(\mathbf{X}) \neq 0.$$

PROOF. The proof relies on the operators $\Phi = D_0(H)$ [2, p. 435] and $H = B_3(f)$. Because of (19) and (essentially) Theorem 1, $|H(\mathbf{X}) - \alpha| < 1$, $|\mathbf{X}| < 1$. Therefore, since the generating function of D_0 is real, we may conclude from [7] that $|\Phi(\mathbf{X}) - \alpha| < 1$.

REMARK. More general cases can be treated using the *Method of Ascent* (see Gilbert [3]).

REFERENCES

1. S. Bergman, *Integral operators in the theory of linear partial differential equations*, Ergebnisse der Mathematik und ihrer Grenzgebiete, N. F., Heft 23, Springer-Verlag, Berlin and New York, 1961. MR 25 #5277.
2. ———, *Classes of solutions of linear partial differential equations in three variables*, Duke Math. J. 13 (1946), 419–458. MR 8, 274.
3. R. P. Gilbert, *Constructive methods for elliptic equations*, Lecture Notes in Math., vol. 365, Springer-Verlag, Berlin and New York, 1974.
4. ———, *Function theoretic methods in partial differential equations*, Math. in Science and Engineering, Vol. 54, Academic Press, New York, 1969. MR 39 #3127.
5. M. Marden, *Geometry of polynomials*, 2nd ed., Math. Surveys, No. 3, Amer. Math. Soc., Providence, R. I., 1966. MR 37 #1562.
6. ———, *A generalization of Weierstrass' and Fekete's mean-value theorems*, Bull. Amer. Math. Soc. 38 (1932), 434–441.
7. ———, *Further mean-value theorems*, Bull. Amer. Math. Soc. 39 (1933), 750–754.
8. ———, *Value distribution of harmonic polynomials in several real variables*, Trans. Amer. Math. Soc. 159 (1971), 137–154. MR 43 #5046.
9. P. A. McCoy, *Value distribution of axisymmetric potentials*, Amer. J. Math. 95 (1973), 419–428. MR 48 #6450.
10. ———, *Value distribution of linear combinations of axisymmetric harmonic polynomials and their derivatives*, Pacific J. Math. 48 (1973), 441–450.
11. ———, *Generalized axisymmetric potentials*, J. Approximation Theory 15 (1975), 256–266.
12. M. Tsuji, *Potential theory in modern function theory*, Maruzen, Tokyo, 1958. MR 22 #5712.

DEPARTMENT OF MATHEMATICS, U. S. NAVAL ACADEMY, ANNAPOLIS, MARYLAND 21401