

ON ODD DIMENSIONAL SURGERY WITH FINITE FUNDAMENTAL GROUP

JEAN-CLAUDE HAUSMANN¹

ABSTRACT. One proves that, for any finite group G and homomorphism $\omega: G \rightarrow \mathbf{Z}/2\mathbf{Z}$, the natural homomorphism $L_{2k+1}^h(\mathbf{Z}G, \omega) \rightarrow L_{2k+1}^h(\mathbf{Q}G, \omega)$ between Wall surgery groups is identically zero. Some results concerning the exponent of $L_{2k+1}^h(\mathbf{Z}G; \omega)$ are deduced.

1. Introduction. Let G be a group and $\omega: G \rightarrow \mathbf{Z}/2\mathbf{Z}$ be a homomorphism (orientation character). Let $L_n^h(G; \omega)$ be the Wall surgery obstruction group in dimension n for surgery to a homotopy equivalence. By changing $\mathbf{Z}G$ into $\mathbf{Q}G$ in the definition of L_n^h , one gets groups $L_n^h(\mathbf{Q}G; \omega)$. They contain the obstruction for surgery to an $[n - 1]/2$ -connected rational homology equivalence. There is a natural homomorphism $r: L_n^h(G; \omega) \rightarrow L_n^h(\mathbf{Q}G; \omega)$ which is used in the long exact localization sequence for surgery obstruction groups of W. Pardon [P1].

Our first result is the following:

THEOREM 1. *For any finite group G and homomorphism $\omega: G \rightarrow \mathbf{Z}/2\mathbf{Z}$, the homomorphism $r: L_{2k+1}^h(G; \omega) \rightarrow L_{2k+1}^h(\mathbf{Q}G; \omega)$ is the zero homomorphism.*

Consequently, in the odd dimensional case with a finite fundamental group, the surgery to a rational homotopy equivalence is always possible. The obstruction for surgery to a homotopy equivalence is always expressible by the linking numbers approach developed in [KM], [W1] and [C]. Since the kernel of r is annihilated by 8 [C], one has

COROLLARY 2. *For any finite group G and homomorphism $\omega: G \rightarrow \mathbf{Z}/2\mathbf{Z}$, $L_{2k+1}^h(G; \omega)$ is annihilated by 8.*

It has been conjectured that this exponent is 4, at least for a large class of finite groups. In this direction, a slight improvement of the proof of Theorem 1 gives the following results:

THEOREM 3. *Let $1 \rightarrow N \rightarrow G \xrightarrow{\Phi} B \rightarrow 1$ be an exact sequence of finite groups, and let $\omega: B \rightarrow \mathbf{Z}/2\mathbf{Z}$ be a homomorphism. Suppose that N is a 2-group. If $y \in \text{Ker}(\Phi_*: L_{2k+1}^h(G; \Phi \circ \omega) \rightarrow L_{2k+1}^h(B; \omega))$, then $4y = 0$.*

Received by the editors May 13, 1976 and, in revised form, July 30, 1976.

AMS (MOS) subject classifications (1970). Primary 18F25, 57D65.

¹ Supported in part by NSF Grant MPS72-05055 A03.

© American Mathematical Society 1977

COROLLARY 4. *For any finite 2-group G and any homomorphism $\omega: G \rightarrow \mathbf{Z}/2\mathbf{Z}$, $L_{2k+1}^h(G; \omega)$ is annihilated by 4.*

COROLLARY 5. *Let G be a finite group whose 2-Sylow subgroup is normal. Suppose that ω is trivial (the orientable case). Then $L_{2k+1}^h(G; \omega)$ is annihilated by 4.*

For instance, the assumptions of Corollary 5 are fulfilled if G is a finite nilpotent group, since a finite nilpotent group is the product of its Sylow subgroups [H].

A part of these statements is more or less known to be obtainable by different methods. Some particular cases are also deducible from published results of other authors. For instance, if k is odd, Theorem 1 is obvious since $L_{2k+1}^h(\mathbf{Q}G; \omega) = 0$ [C]. On the other hand, in the orientable case (ω trivial), Corollary 2 can be deduced from [W3, exact sequence, p. 78 and remark (4), p. 2]. Corollaries 4 and 5 are deducible from [W3] when G is abelian. Recently, W. Pardon independently found an algebraic proof of Corollary 4 and A. Bak computed $L_n^h(G; \omega)$ when G has its 2-Sylow subgroup normal and abelian [Bak2]. Finally, an announcement of Theorem 1 when G is abelian was published by R. M. Geist [G]. Our proofs are independent from these results and our approach is quite different.

In §2, we give a sufficient condition for a degree one map between a manifold pair and a Poincaré pair (of odd dimension) to be a rational homotopy equivalence. In §3, we establish a functoriality property for a part of the localization exact sequence of surgery groups ([P1] and [P2]). It would be interesting to know in which generality such a functoriality property holds.

Results of §§2 and 3 are used in §4 for proving Theorem 1; finally, Theorem 4 and Corollaries 4 and 5 are proved in §5.

I am grateful to W. Pardon for conversations, to C.T.C. Wall and A. Bak for commenting on these results, and to the referee for a great simplification of my original proof of Proposition 2.1.

2. A class of rational homology equivalences. Denote, as usual, by $\mathbf{Z}_{(p)}$ the subring of \mathbf{Q} of fractions expressible with a denominator prime to p . This section is devoted to proving the following proposition:

PROPOSITION 2.1. *Let (X, Y) be a Poincaré pair of dimension $2k + 1$ such that $\pi_1(Y) \simeq \pi_1(X)$ is a finite group G . Let $N \subset G$ be a normal p -subgroup of G (p some prime). Consider a map $f: (M; \partial M) \rightarrow (X; Y)$ of degree one (M^{2k+1} a compact manifold), with $f|_{\partial M}$ a homotopy equivalence. Suppose that f is k -connected and is a $\mathbf{Z}(G/N)$ -homology equivalence (local coefficients). Then f is a $\mathbf{Z}_{(p)}$ -homology equivalence (and thus a rational homology equivalence).*

PROOF. If B is a $\mathbf{Z}G$ -module, we denote by $K_k(M; B)$ the $\mathbf{Z}G$ -module $H_{k+1}(f; B)$ and $K_k(M)$ is used for $K_k(M; \mathbf{Z}G)$. By [CS, Lemma 1.4] one has $K_k(M; \mathbf{Z}(G/N)) = K_k(M) \otimes_{\mathbf{Z}G} \mathbf{Z}(G/N)$. This last module is \mathbf{Z} -isomorphic to $K_k(M) \otimes_{\mathbf{Z}N} \mathbf{Z} = K_k(M)/I \cdot K_k(M)$, where I is the augmentation ideal of N .

Since f is a $\mathbf{Z}(G/N)$ -homology equivalence, one has $K_k(M; \mathbf{Z}(G/N)) = 0$, and then $K_k(M) = I \cdot K_k(M)$.

Since $K_k(M)$ is finitely generated [W2, Lemma 2.3] and G is finite, $K_k(M)/pK_k(M)$ is a finite abelian p -group. The action of the finite p -group N on any finite abelian p -group L is nilpotent, i.e. $I^s \cdot L = 0$ for s large enough. (The proof goes like in [H, pp. 47 and 155].) Therefore $K_k(M) = pK_k(M)$, which is equivalent to $K_k(M) \otimes_{\mathbf{Z}} \mathbf{Z}_{(p)} = 0$. But $K_k(M) \otimes_{\mathbf{Z}} \mathbf{Z}_{(p)} = K_k(M) \otimes_{\mathbf{Z}G} \mathbf{Z}_{(p)}G = K_k(M; \mathbf{Z}_{(p)}G)$, the last isomorphism by [CS, Lemma 1.4]. Thus, $K_k(M; \mathbf{Z}_{(p)}G) = 0$ and f is a $\mathbf{Z}_{(p)}$ -homology equivalence.

3. Partial functoriality for the surgery group localization exact sequence. We will use a functorial property of the following leg of the Pardon exact sequence [P1]:

$$(3.1) \quad \begin{aligned} L_{2k+2}^h(\mathbf{Q}G, \omega) &\rightarrow L_{2k+1}^t(\mathbf{Z}G; \mathbf{Z} - \{0\}) \\ &\rightarrow L_{2k+1}^h(G, \omega) \rightarrow L_{2k+1}^h(\mathbf{Q}G, \omega). \end{aligned}$$

This functoriality property is implied by the corresponding property for this other formulation of the sequence [P2, Theorem 2.1]:

$$(3.2) \quad W_0^{-\lambda}(\mathbf{Q}G) \rightarrow W_0^{-\lambda}(\mathbf{Q}G/\mathbf{Z}G) \rightarrow W_1^{\lambda}(\mathbf{Z}G) \rightarrow W_1^{\lambda}(\mathbf{Q}G)$$

where $\lambda = (-1)^k$. (See [P2] for the definitions.) Sequences (3.1) and (3.2) are related by the following commutative diagram:

$$(3.3) \quad \begin{array}{ccccccc} W_0^{-\lambda}(\mathbf{Q}G) & \longrightarrow & W_0^{-\lambda}(\mathbf{Q}G/\mathbf{Z}G) & \longrightarrow & W_1^{\lambda}(\mathbf{Z}G) & \longrightarrow & W_1^{\lambda}(\mathbf{Q}G) \\ \downarrow \cong & & \downarrow \cong & & \downarrow & & \downarrow \\ L_{2k+2}^h(\mathbf{Q}G, \omega) & \longrightarrow & L_{2k+1}^t(\mathbf{Z}G; \mathbf{Z} - \{0\}) & \longrightarrow & L_{2k+1}^h(G; \omega) & \longrightarrow & L_{2k+1}^h(\mathbf{Q}G; \omega) \end{array}$$

where the left two vertical arrows are identity maps (the groups are identical) and two right vertical arrows divide out by $w_1^{\lambda} = \begin{pmatrix} 0 & 1 \\ \lambda & 0 \end{pmatrix}$.

Let \mathcal{F} be the category whose objects are pairs $(G; \omega)$, where G is a finite group and $\omega: G \rightarrow \mathbf{Z}/2\mathbf{Z}$ is a homomorphism. A morphism $F: (G, \omega_G) \rightarrow (H, \omega_H)$ of \mathcal{F} is a homomorphism $f: G \rightarrow H$ such that $\omega_G = \omega_H \circ f$. Let Ab denote the category of abelian groups. The group rings RG ($R = \mathbf{Z}$ or \mathbf{Q}) are understood to be endowed with the involution $\sum n_g g = \sum n_g \omega(g)g^{-1}$.

PROPOSITION 3.4. *The correspondences*

$$(G, \omega) \rightarrow W_i^{\lambda}(RG), \quad i = 0 \text{ or } 1, \lambda = \pm 1,$$

and

$$(G, \omega) \mapsto W_0^\lambda(\mathbf{Q}G/\mathbf{Z}G)$$

give rise to functors from \mathfrak{F} to Ab so that sequence (3.2) (and thus (3.1)) is functorial.

PROOF. $W_1^\lambda(RG)$ are functors in the usual way. The only nonobvious point is to define $f_*: W_0^\lambda(\mathbf{Q}G/\mathbf{Z}G) \rightarrow W_0^\lambda(\mathbf{Q}H/\mathbf{Z}H)$ for an \mathfrak{F} -morphism $f: (G, \omega_G) \rightarrow (H, \omega_H)$. By definition of $W_0^\lambda(\mathbf{Q}G/\mathbf{Z}G)$ [P2, p. 10] it suffices to have f_* defined on classes in W_0^λ which are represented by a triple (M, φ, ψ) , where

- (1) M is a finite $\mathbf{Z}G$ -module admitting a short free $\mathbf{Z}G$ -resolution

$$0 \rightarrow F_1 \xrightarrow{\mu} F_0 \rightarrow M \rightarrow 0.$$

- (2) $\varphi: M \times M \rightarrow \mathbf{Q}G/\mathbf{Z}G$ is a nonsingular λ -hermitian form.

- (3) $\psi: M \rightarrow \mathbf{Q}G/S_\lambda(\mathbf{Z}G)$ is a function satisfying (i)–(iii) of [P2].

Let us define the image by f_* of a class (M, φ, ψ) to be the class of (M', φ', ψ') , where

- (1) $M' = M \otimes_{\mathbf{Z}G} \mathbf{Z}H$.

- (2) $\varphi': M' \times M' \rightarrow \mathbf{Q}H/\mathbf{Z}H$ is defined by $\varphi'(x \otimes a, y \otimes b) = \bar{a}f(\varphi(x, y))b$ (see [Ba, §6]).

- (3) $\psi': M' \rightarrow \mathbf{Q}H/S_\lambda(\mathbf{Z}H)$ is the unique extension of ψ such that $\psi'(x \otimes 1) = f(\psi(x))$ [Ba, 6.3].

Let us check that (M', φ', ψ') actually determines a class in $W_0^\lambda(\mathbf{Q}H/\mathbf{Z}H)$. M' is a finite $\mathbf{Z}H$ -module; it admits the following short free resolution:

$$0 \rightarrow F_1 \otimes \mathbf{Z}H \xrightarrow{\mu \otimes 1} F_0 \otimes \mathbf{Z}H \rightarrow M' \rightarrow 0$$

(the tensor product is always understood over $\mathbf{Z}G$). Indeed $F_i \otimes \mathbf{Z}H$ are $\mathbf{Z}H$ -free of same $\mathbf{Z}H$ -rank. Since H is a finite group, $F_1 \otimes \mathbf{Z}H$ and $F_2 \otimes \mathbf{Z}H$ have same finite \mathbf{Z} -rank. Thus $\mu \otimes 1$ is injective.

To prove that φ' is nonsingular, it suffices [Ba, p. 45] to check that the homomorphism

$$j_M: \text{Hom}_{\mathbf{Z}G}(M; \mathbf{Q}G/\mathbf{Z}G) \otimes \mathbf{Z}H \rightarrow \text{Hom}_{\mathbf{Z}H}(M'; \mathbf{Q}H/\mathbf{Z}H)$$

given by $j_M(h \otimes a)(x \otimes b) = \bar{a}f(h(x))b$ is an isomorphism (observe that $\mathbf{Q}G/\mathbf{Z}G \otimes \mathbf{Z}H \simeq \mathbf{Q}H/\mathbf{Z}H$). By [P2, Proposition 1.4], $\text{Hom}_{\mathbf{Z}G}(M; \mathbf{Q}G/\mathbf{Z}G)$ has the following free resolution:

$$0 \rightarrow \text{Hom}_{\mathbf{Z}G}(F_0; \mathbf{Z}G) \xrightarrow{\bar{\mu}} \text{Hom}_{\mathbf{Z}G}(F_1; \mathbf{Z}G) \rightarrow \text{Hom}_{\mathbf{Z}G}(M; \mathbf{Q}G/\mathbf{Z}G) \rightarrow 0.$$

As above, tensoring by $\mathbf{Z}H$ leaves this sequence exact. Hence, one has

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 \text{Hom}_{\mathbf{Z}G}(F_0; \mathbf{Z}G) \otimes \mathbf{Z}H & \xrightarrow{j_{F_0}} & \text{Hom}_{\mathbf{Z}H}(F_0 \otimes \mathbf{Z}H; \mathbf{Z}H) \\
 \downarrow \bar{\mu} \otimes 1 & & \downarrow \bar{\mu} \otimes 1 \\
 \text{Hom}_{\mathbf{Z}G}(F_1; \mathbf{Z}G) \otimes \mathbf{Z}H & \xrightarrow{j_{F_1}} & \text{Hom}_{\mathbf{Z}H}(F_1 \otimes \mathbf{Z}H; \mathbf{Z}H) \\
 \downarrow & & \downarrow \\
 \text{Hom}_{\mathbf{Z}G}(M; \mathbf{Q}G/\mathbf{Z}G) & \xrightarrow{j_M} & \text{Hom}_{\mathbf{Z}H}(M'; \mathbf{Q}H/\mathbf{Z}H) \\
 \downarrow & & \downarrow \\
 0 & & 0
 \end{array}$$

The commutativity of the lowest square comes from the definitions of j_{F_1} and j_M and of the identification of $\text{coker } \bar{\mu}$ with $\text{Hom}_{\mathbf{Z}G}(M; \mathbf{Q}G/\mathbf{Z}G)$ [P2, (1.5)]. Clearly, $j_{\mathbf{Z}G}$ is an isomorphism. By additivity, j_{F_0} and j_{F_1} are isomorphisms. Thus, j_M is an isomorphism.

Finally, one checks without difficulty that a kernel is mapped onto a kernel. Thus $f_*: W_0^\lambda(\mathbf{Q}G/\mathbf{Z}G) \rightarrow W_0^\lambda(\mathbf{Q}H/\mathbf{Z}H)$ is well defined.

The functoriality of sequence (3.2) then comes directly from the definitions of each arrow.

4. Proof of Theorem 1. Theorem 1 will be proven first when G is a finite 2-group, then when G is a 2-hyerelementary group and last in the general case using the induction theorem due to Dress. We shall make use of the following lemma, in which $r: L_{2k+1}^h(G; \Phi \circ \omega) \rightarrow L_{2k+1}^h(\mathbf{Q}G; \Phi \circ \omega)$ is the natural homomorphism, as in the Introduction.

LEMMA 4.1. *Let $1 \rightarrow N \rightarrow G \xrightarrow{\Phi} B \rightarrow 1$ be an extension of finite groups, with N a p -group. Let $\omega: B \rightarrow \mathbf{Z}/2\mathbf{Z}$ be a homomorphism. Then, for all $x \in \text{Ker}(L_{2k+1}^h(G; \omega \circ \Phi) \rightarrow L_{2k+1}^h(B; \omega))$, one has $r(x) = 0$.*

PROOF. Let M^{2k} be a manifold, $k \geq 3$, with $\pi_1(M) = G$ and orientation homomorphism $\omega \circ \Phi$. Accordingly [W2, Theorem 6.5] there exists a cobordism (W^{2k+1}, M, M') and a normal map of degree one $(f, b) f: (W, M, M') \rightarrow (M \times I, M \times \{0\}, M \times \{1\})$ such that $f|M = \text{id}_M$, the map $f|M'$ is a homotopy equivalence, and the surgery obstruction $\theta(f, b) \in L_{2k+1}(G; \omega \circ \Phi)$ is equal to x .

Let $\Gamma_{2k+1}^h(\mathbf{Z}G \xrightarrow{\Phi} \mathbf{Z}B)$ be the Cappell-Shaneson surgery obstruction group [CS]. This is a subgroup of $L_{2k+1}^h(B; \omega)$. Therefore, the assumption on x implies that x belongs to the kernel of the natural homomorphism

$$L_{2k+1}^h(G; \omega \circ \Phi) \rightarrow \Gamma_{2k+1}^h(\mathbf{Z}G \xrightarrow{\Phi} \mathbf{Z}B).$$

Hence, by [CS, Proposition 2.1], (f, b) is normally cobordant to (f', b') where f' is a k -connected \mathbf{ZB} -homology equivalence. By Proposition 2.1, f' is a rational homology equivalence and then, by [C, Theorem 4.10], $r(x) = 0$.

PROOF OF THEOREM 1.

Case 1. G is a finite 2-group. If ω is the trivial homomorphism, the result follows from Lemma 4.1 applied to $N = G$. If ω is not trivial, one uses Lemma 4.1 applied to the case $B = \mathbf{Z}/2\mathbf{Z}$ and $\Phi = \omega_G$ together with the fact that $L_{2k+1}^h(\mathbf{Z}/2\mathbf{Z}; \text{id}) = 0$ [W2, Theorem 13.A.1].

Case 2. G is a 2-hyerelementary group. This means that G has a decomposition $1 \rightarrow C \rightarrow G \rightarrow P \rightarrow 1$, where P is a 2-group and C is cyclic. The proof will be by induction on $d(G)$, the greatest odd divisor of $|G|$. If $d(G) = 1$, G is a 2-group and we can use Case 1. If $d > 1$, one may write $C = C_1 \times C_2$, where C_1 is a cyclic p -group with $p \neq 2$, and $(p, |C_2|) = 1$. Thus, one has a split exact sequence:

$$1 \longrightarrow C_1 \longrightarrow G \xrightarrow[\substack{\Phi \\ s}]{\substack{\Phi \\ s}} \bar{G} \longrightarrow 1$$

and $\omega_G: G \rightarrow \mathbf{Z}/2\mathbf{Z}$ factors through $\omega_{\bar{G}}: \bar{G} \rightarrow \mathbf{Z}/2\mathbf{Z}$. \bar{G} is a 2-hyerelementary group with $d(\bar{G}) < d(G)$. By induction hypothesis, $r: L_{2k+1}^h(\bar{G}; \omega_{\bar{G}}) \rightarrow L_{2k+1}^h(\mathbf{Q}\bar{G}; \omega_{\bar{G}})$ is zero. Using the functoriality of sequence (3.1), one gets a diagram:

$$\begin{array}{ccccc} \longrightarrow & L_{2k+1}^t(\mathbf{Z}G; \mathbf{Z} - \{0\}) & \xrightarrow{i} & L_{2k+1}^h(G; \omega_G) & \xrightarrow{r} & L_{2k+1}^h(\mathbf{Q}G; \omega_G) \\ & \downarrow \Phi^t \Big\} s^t & & \downarrow \Phi_* \Big\} s_* & & \downarrow \\ \longrightarrow & L_{2k+1}^t(\mathbf{Z}\bar{G}; \mathbf{Z} - \{0\}) & \xrightarrow{\bar{i}} & L_{2k+1}^h(\bar{G}; \omega_{\bar{G}}) & \xrightarrow{\bar{r}} & L_{2k+1}^h(\mathbf{Q}\bar{G}; \omega_{\bar{G}}) \\ & & & \searrow 0 & & \end{array}$$

Thus one can write $x \in L_{2k+1}^h(G; \omega_G)$ as $x = x_1 + x_2$ with $x_1 = i(y)$ and $x_2 \in \ker \Phi_*$. By Lemma 4.1, $r(x) = 0$.

Case 3. General case. Since $L_{\text{odd}}^h(G; \omega)$ is a 2-group [C], the Dress induction theorem [D] asserts that the product of restrictions $L_{2k+1}^h(G; \omega) \rightarrow \prod_{H \in \mathfrak{H}(G)} L_{2k+1}^h(H; \omega|_H)$ is injective, where $\mathfrak{H}(G)$ is the set of 2-hyerelementary subgroups of G . The same holds for $L_{2k+1}^h(\mathbf{Q}G, \omega)$. Theorem 1 then follows, using Case 2.

5. Proof of Theorem 3 and Corollaries 4 and 5. Let M^{2k} be a closed manifold with $\pi_1(M) = G$ and orientation character $\omega \circ \Phi$. Represent y as the surgery obstruction for a normal map of degree one:

$$f: (W^{2k+1}, M, M') \rightarrow (M \times I, M \times \{0\}, M \times \{1\}).$$

As in the proof of Lemma 4.1, the condition on y implies that f is normally cobordant to an f' that is a k -connected \mathbf{ZB} -homology equivalence. By Proposition 2.1, f' is a $\mathbf{Z}_{(2)}$ -homology equivalence. Therefore, using [PP, §5], one can consider y as an element of $J_k(G)$, the Grothendieck group based on

linking forms over finite ZG -modules without 2-torsion. $J_k(G)$ has exponent 4 [PP, Theorem 5.1]. Thus $4y = 0$.

Corollary 4 is a consequence of Theorem 4, applied to $N = G$ in the orientable case and to $\Phi = \omega$ otherwise (see the proof of Theorem 1, Case 1). For Corollary 5, one uses Theorem 3 for N equal to the normal 2-Sylow group of G , and the fact that $L_{\text{odd}}^h(H) = 0$ when H has odd order ([Bak] or [P1]).

REFERENCES

- [Bak] A. Bak, *Odd dimension surgery groups of odd torsion groups vanish*, *Topology* **14** (1975), 367–374.
- [Bak2] ———, *The computation of surgery groups of finite groups with abelian 2-hyperelementary*, *Algebraic K-theory, Lecture Notes in Math.*, Springer, Berlin and New York, vol. 551, pp. 384–409.
- [Ba] H. Bass, *Unitary algebraic K-theory*, Proc. Conf. Seattle Res. Center, Batelle Memorial Inst., 1972, *Lecture Notes in Math.*, vol. 343, Springer-Verlag, Berlin and New York, 1973. MR **51** #8211.
- [CS] S. Cappell and J. Shaneson, *The codimension two placement problem and homology equivalent manifolds*, *Ann. of Math. (2)* **99** (1974), 277–348. MR **49** #3978.
- [C] F. Connolly, *Linking numbers and surgery*, *Topology* **12** (1973), 389–409.
- [D] A. Dress, *Induction and structure theorem for orthogonal representations of finite groups*, *Ann. of Math. (2)* **102** (1975), 291–326.
- [G] R. M. Geist, *Semicharacteristic detection of obstructions to rational homotopy equivalence*, *Notices Amer. Math. Soc.* **21** (1974), A-451. Abstract #74T-G72.
- [H] M. Hall, *The theory of groups*, Macmillan, New York, 1959. MR **21** #1996.
- [KM] M. Kervaire and J. Milnor, *Groups of homotopy spheres. I*, *Ann. of Math. (2)* **77** (1963), 504–537. MR **26** #5584.
- [P1] W. Pardon, *Local surgery and application to the theory of quadratic forms*, *Bull. Amer. Math. Soc.* **82** (1976), 131–134.
- [P2] ———, *The exact sequence of a localization for Witt groups*, *Algebraic K-Theory, Lecture Notes in Math.*, vol. 551, Springer, Berlin and New York, pp. 303–310.
- [PP] D. Passman and T. Petrie, *Surgery with coefficients in a field*, *Ann. of Math. (2)* **95** (1972), 385–405. MR **46** #10011.
- [W1] C. T. C. Wall, *Surgery of non-simply-connected manifolds*, *Ann. of Math. (2)* **84** (1966), 217–276. MR **35** #3692.
- [W2] ———, *Surgery on compact manifolds*, Academic Press, New York, 1970.
- [W3] ———, *Classification of hermitian forms. VI. Group rings*, *Ann. of Math. (2)* **103** (1976), 1–80.

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NEW JERSEY 08540

Current address: Section de Mathématiques Université, 2-4 rue de Lièvre, 1211 Geneva 24, Switzerland