

SYMMETRY FOR FINITE DIMENSIONAL HOPF ALGEBRAS

J. E. HUMPHREYS¹

ABSTRACT. This note refines criteria given by R. G. Larson and M. E. Sweedler for a finite dimensional Hopf algebra to be a symmetric algebra, with applications to restricted universal enveloping algebras and to certain finite dimensional subalgebras of the hyperalgebra of a semisimple algebraic group in characteristic p .

Let A be a finite dimensional associative algebra over a field K . Then A is called *Frobenius* if there exists a nondegenerate bilinear form $f: A \times A \rightarrow K$ which is associative in the sense that $f(ab, c) = f(a, bc)$ for all $a, b, c \in A$ [3, Chapter IX]. A is called *symmetric* if there exists a symmetric form of this type [3, §66]. For example, semisimple algebras and group algebras of finite groups are symmetric. We investigate here the extent to which finite dimensional Hopf algebras (with antipode) are symmetric; they are always Frobenius, thanks to the main theorem of [8].

1. Hopf algebras. In this section H denotes a finite dimensional Hopf algebra over an arbitrary field K , with antipode s and augmentation $\varepsilon: H \rightarrow K$. According to the main theorem of [8], existence of the antipode implies (and is implied by) the existence of a (nonsingular) left *integral* $\Lambda \in H$, which is unique up to scalar multiples. By definition, Λ satisfies: $h\Lambda = \varepsilon(h)\Lambda$, for all $h \in H$. Equally well, H has a right integral Λ' , unique up to scalar multiples. If Λ' is proportional to Λ , H is called *unimodular*.

With a left integral Λ is associated a nondegenerate bilinear associative form b on H [8, §7]. As a result, H is a Frobenius algebra. From the second corollary of Proposition 8 in [8], applied to the dual Hopf algebra (whose antipode has the same order as s), we obtain immediately:

THEOREM 1. *With notation as above, b is symmetric if and only if H is unimodular and $s^2 = 1$. In particular, if the latter conditions hold, then H is symmetric.*

We can apply this to the algebras u_n ($n = 1, 2, \dots$) defined in [6, Appendix U], [7]. These are finite dimensional Hopf subalgebras of the hyperalgebra U_K of a simply connected, semisimple algebraic group G over an algebraically

Received by the editors February 28, 1977.

AMS (MOS) subject classifications (1970). Primary 16A24, 16A36, 17B50; Secondary 17B45.

Key words and phrases. Hopf algebra, Frobenius algebra, symmetric algebra, restricted Lie algebra, integral, unimodular, hyperalgebra of semisimple algebraic group.

¹Research partially supported by NSF Grant MCS72-05055 A04.

© American Mathematical Society 1978

closed field K of characteristic $p > 0$, with antipode of order 2.

COROLLARY. *The algebras u_n are symmetric.*

PROOF. G acts naturally on u_n as a group of Hopf algebra automorphisms (adjoint action) and therefore $g\Lambda$ is another left integral, if $g \in G$ and Λ is a left integral in u_n ; this must be of the form $\alpha(g)\Lambda$ for $\alpha(g) \in K^*$. Since $G = (G, G)$, it has no nontrivial homomorphisms into K^* , forcing $\alpha = 1$; therefore, Λ is G -invariant. In turn, Λ is invariant under the derived adjoint action of u_n , i.e., Λ lies in the center of u_n and is therefore also a right integral. Now u_n is unimodular, with antipode of order 2, so the theorem applies. Q.E.D.

REMARKS. (1) W. J. Haboush [4] has recently been able to construct explicitly an integral for each u_n , and has used these to obtain a new family of "central differential operators" which are capable of separating the various Steinberg modules for G . His work inspired the present note, in particular the proof of the corollary.

(2) It follows from the corollary that the *Cartan matrix* C of u_n , recording the composition factor multiplicities of principal indecomposable modules, is symmetric. Perhaps it is even of the form $C = 'D \cdot D$ for a suitable "decomposition matrix" D . This is the case when $n = 1$ [5], u_1 being the restricted universal enveloping algebra of the Lie algebra of G .

THEOREM 2. *If H is symmetric, then H is unimodular.*

PROOF. Let f be a nondegenerate symmetric, associative bilinear form on H , and let Λ (resp. Λ') be a left (resp. right) integral. If $h \in \text{Ker } \epsilon$, $f(h, \Lambda) = f(1, h\Lambda) = 0 = f(\Lambda'h, 1) = f(\Lambda', h)$. Since f is symmetric, both Λ and Λ' lie in the orthogonal complement of $\text{Ker } \epsilon$, which has dimension 1. So Λ is proportional to Λ' . Q.E.D.

If H is commutative or cocommutative, then automatically $s^2 = 1$ (cf. [8, p. 77]); in any event, s has finite order [9]. But examples are known for which s has higher (necessarily even) order. It is not clear whether such algebras can be unimodular without being symmetric.

2. Restricted enveloping algebras. In this section K is a field of characteristic $p > 0$, L a restricted Lie algebra (Lie p -algebra) over K , $u(L)$ its restricted universal enveloping algebra, which is a finite dimensional Hopf algebra with antipode of order 1 or 2. A. Berkson [1] showed directly that $u(L)$ is Frobenius by using a bilinear form f defined as follows. Fix an ordered basis x_1, \dots, x_n of L , so the monomials $x_1^{i_1} \cdots x_n^{i_n}$ ($0 \leq i_j < p$) form a basis of $u(L)$. Let φ_0 be the linear function taking value 1 at $u_0 = x_1^{p-1} \cdots x_n^{p-1}$ and value 0 at other monomials, and define $f(u, v) = \varphi_0(uv)$. In turn, J. R. Schue proved:

THEOREM 3 [10]. *The bilinear form f is symmetric if and only if $\text{Tr}(\text{ad } x) = 0$*

for all $x \in L$. In particular, if the latter condition is satisfied, $u(L)$ is symmetric.

Here $\text{ad } x(y) = [xy]$; the notation D_x is used in [10] and $D(x)$ in [8]. This criterion for symmetry was applied in [5] to show that the algebra u_1 discussed above is symmetric. On the other hand, an example given in [8, p. 85] to show that $u(L)$ need not be unimodular also fails (as it must, by Theorems 2 and 3) to meet Schue's criterion.

The third corollary of Proposition 8 in [8] states that $u(L)$ is unimodular if and only if $\text{Tr}(\text{ad } x) = 0$ for all $x \in L$. We offer here a different proof, based on the following lemma.

LEMMA. Let Λ be a left integral in $u(L)$. If x_1, \dots, x_n is any ordered basis of L , and Λ is written as a linear combination of the monomials $x_1^{i_1} \cdots x_n^{i_n}$, then $u_0 = x_1^{p-1} \cdots x_n^{p-1}$ must occur with nonzero coefficient.

PROOF. Suppose the contrary. Define the degree of $x_1^{i_1} \cdots x_n^{i_n}$ to be $\sum i_j$, and write $\Lambda = \Lambda_1 + \Lambda_2$, where Λ_1 contains all monomials of the highest degree d occurring (so $d < n(p - 1)$). If we change basis by permuting the x_i , the new expression for Λ will be of the form $\Lambda'_1 + \Lambda_3$, where Λ'_1 is obtained from Λ_1 by permuting the x_i and the monomials in Λ_3 are again of degree $< d$ (cf. the proof of Lemma 1 in [10]). We may therefore assume that not all monomials occurring in Λ_1 involve the factor x_1^{p-1} .

The augmentation for $u(L)$ maps all x_i to 0, so by definition of left integral, $0 = x_1\Lambda = x_1\Lambda_1 + x_1\Lambda_2$. Because of the p -structure, $x_1^p \in L$, so $x_1^p x_2^{i_2} \cdots x_n^{i_n}$ can be rewritten in $u(L)$ as a linear combination of monomials having degrees $< p + \sum i_j$ (sum over $j \neq 1$). In particular, $x_1\Lambda_2$ involves only monomials of degree $\leq d$, while $x_1\Lambda_1$ involves such monomials along with one or more linearly independent monomials of degree $d + 1$ (corresponding to monomials in Λ_1 not involving x_1^{p-1}). This is clearly impossible. Q.E.D.

We remark that the integral constructed by Haboush [4] for the algebra u_1 illustrates this lemma very nicely.

THEOREM 4. If $u(L)$ is a symmetric algebra, then $\text{Tr}(\text{ad } x) = 0$ for all $x \in L$.

PROOF. Since $u(L)$ is symmetric, it is unimodular (Theorem 2), so any integral Λ lies in the center of $u(L)$. Choose an ordered basis x_1, \dots, x_n of L , and define the form f as above, relative to the basis of $u(L)$ consisting of monomials. By the lemma, we may (after multiplying Λ by a nonzero scalar) write $\Lambda = u_0 + u_1$, where u_1 is a linear combination of monomials of degrees $< n(p - 1)$. With φ_0 as above, it follows from Lemma 1 of [10] (cf. proof of theorem) that $\varphi_0(u_1x) = \varphi_0(xu_1)$ for all $x \in L$. On the other hand, Lemma 3 of [10] says that $\varphi_0(u_0x) = \varphi_0(xu_0) + \text{Tr}(\text{ad } x)$ for $x \in L$. Since $\Lambda x = x\Lambda$, it follows at once that $\text{Tr}(\text{ad } x) = 0$. Q.E.D.

Theorems 3 and 4, combined with Theorems 1 and 2, show that $u(L)$ is unimodular if and only if $\text{Tr}(\text{ad } x) = 0$ for all $x \in L$, as stated in [8]. This is analogous to the classical criterion for a Lie group G to be unimodular:

$\det(\text{Ad } g) = 1$ for all $g \in G$ [2, Chapter III, 3, no. 16, Corollary to Proposition 55].

REFERENCES

1. A. Berkson, *The u -algebra of a restricted Lie algebra is Frobenius*, Proc. Amer. Math. Soc. **15** (1964), 14–15. MR **28** #2132.
2. N. Bourbaki, *Groupes et algèbres de Lie*, Chapters II–III, Hermann, Paris, 1972.
3. C. W. Curtis and I. Reiner, *Representation theory of finite groups and associative algebras*, Interscience, New York, 1962. MR **26** #2519.
4. W. J. Haboush, (manuscript).
5. J. E. Humphreys, *Modular representations of classical Lie algebras and semisimple groups*, J. Algebra **19** (1971), 51–79. MR **44** #271.
6. ———, *Ordinary and modular representations of Chevalley groups*, Lecture Notes in Math., vol. 528, Springer-Verlag, New York and Berlin, 1976.
7. ———, *On the hyperalgebra of a semisimple algebraic group*, Contributions to Algebra: A Collection of Papers Dedicated to Ellis Kolchin, Academic Press, New York, 1977.
8. R. G. Larson and M. E. Sweedler, *An associative orthogonal bilinear form for Hopf algebras*, Amer. J. Math. **91** (1969), 75–94. MR **39** #1523.
9. D. E. Radford, *The order of the antipode of a finite-dimensional Hopf algebra*, Amer. J. Math. **98** (1976), 333–355. MR **53** #10852.
10. J. R. Schue, *Symmetry for the enveloping algebra of a restricted Lie algebra*, Proc. Amer. Math. Soc. **16** (1965), 1123–1124. MR **32** #2515.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MASSACHUSETTS, AMHERST, MASSACHUSETTS 01003

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NEW JERSEY 08540

Current address: Department of Mathematics, University of Massachusetts, Amherst, Massachusetts 01003