

PENCILS OF HIGHER DERIVATIONS OF ARBITRARY FIELD EXTENSIONS

JAMES K. DEVENEY AND JOHN N. MORDESON

ABSTRACT. Let L be a field of characteristic $p \neq 0$. A subfield K of L is Galois if K is the field of constants of a group of pencils of higher derivations on L . Let $F \supset K$ be Galois subfields of L . Then the group of L over F is a normal subgroup of the group of L over K if and only if $F = K(L^{p^r})$ for some nonnegative integer r . If L/K splits as the tensor product of a purely inseparable extension and a separable extension, then the algebraic closure of K in L , \bar{K} , is also Galois in L . Given K , for every Galois extension L of K , \bar{K} is also Galois in L if and only if $[K : K^p] < \infty$.

0. Introduction. Throughout we assume L is a field of characteristic $p \neq 0$. A rank t higher derivation on L is a sequence $d = \{d_i | 0 \leq i < t + 1\}$ of additive maps of L into L such that

$$d_r(ab) = \sum \{d_i(a)d_j(b) | i + j = r\}$$

and d_0 is the identity map. The set of all rank t higher derivations forms a group with respect to the composition $d \circ e = f$ where $f_j = \sum \{d_m e_n | m + n = j\}$. Let $H(L/K)$ be the set of all higher derivations on L trivial on K and having rank some power of p . Given d in $H(L/K)$, $v(d) = f$ where $\text{rank } f = p(\text{rank } d)$, $f_{pi} = d_i$ and $f_j = 0$ if $p \nmid j$. Two higher derivations f and g are equivalent if $g = v^i(f)$ or $f = v^i(g)$ for some i . The equivalence class of d is \bar{d} and is called the pencil of d . The set of all pencils, $\bar{H}(L/K)$, can be given a group structure by defining $\bar{d}\bar{f}$ to be the pencil of $d'f'$ where $d' \in \bar{d}$, $f' \in \bar{f}$ and $\text{rank } d' = \text{rank } f'$ [3]. A subfield K of L will be called Galois if K is the field of constants of a group of pencils on L or equivalently if L/K is modular and $\bigcap_i K(L^{p^i}) = K$ [2, Proposition 1]. In §1 it is shown that if $F \supset K$ are Galois subfields of L , then $\bar{H}(L/F)$ is an invariant subgroup of $\bar{H}(L/K)$ if and only if $F = K(L^{p^r})$ for some nonnegative integer r . This generalizes the result given in [2, Theorem 8] for the bounded exponent finite transcendence degree case.

Let \bar{K} denote the algebraic closure of K in L . L/K is said to split when $L = J \otimes_K D$ where J/K is purely inseparable and D/K is separable. §2 examines the question of when \bar{K} is Galois in L , given L/K is Galois. Sufficient conditions are shown to be the splitting of L/K . Moreover, for every Galois extension L of K , \bar{K} is also Galois in L if and only if

Presented to the Society, August 10, 1978 under the title *On pencil Galois theory*; received by the editors March 8, 1978.

AMS (MOS) subject classifications (1970). Primary 12F15.

Key words and phrases. Modular field extension, pencils of higher derivations.

© 1979 American Mathematical Society
 0002-9939/79/0000-0200/\$02.75

$[K : K^p] < \infty$ (and in this case L/K splits). In view of these results it appeared that \bar{K} being Galois in L was related to L/K splitting. However, an example is constructed with L/K and L/\bar{K} both Galois and yet L/K does not split.

Pencils of higher derivations were originally constructed by Heerema to incorporate into a single theory the Galois theories of finite and infinite rank higher derivations. Basically the infinite higher derivations would be the group of L/\bar{K} (L/\bar{K} being separable). However, in the proof of Theorem 2.2, an example of a Galois extension is constructed with L/\bar{K} being relatively perfect, and hence having no infinite rank higher derivations. Thus in this most general setting some different fields of constants are obtained.

1. Invariant subgroups. Let $F \supset K$ be Galois subfields of L . This section develops necessary and sufficient conditions for $\bar{H}(L/F)$ to be $\bar{H}(L/K)$ -invariant.

(1.1) LEMMA. *Suppose L/K is purely inseparable Galois. Let F^* be an intermediate field of L/K such that L/F^* is modular and F^*/K has exponent ≤ 1 . If for every maximal pure independent set M of L/K every element of M has the same exponent over F^* that it has over K , then $F^* = K$.*

PROOF. Suppose some c in L has c^{p^i} in F^* but not in $K(K^{p^{-1}} \cap L^{p^{i+1}})$. By modularity,

$$K(K^{p^{-1}} \cap L^{p^{i+1}}) = K(L^{p^{i+1}}) \cap K^{p^{-1}},$$

and hence c^{p^i} is not in $K(L^{p^{i+1}})$. For $j \leq i$, c^{p^j} cannot be in $K(L^{p^{j+1}})$. Thus c is pure independent [9] and is part of a maximal pure independent set of L/K . But c has exponent $i + 1$ over K and exponent i over F^* , contrary to the hypothesis. Hence

$$F^* \cap L^{p^i} \subseteq K(K^{p^{-1}} \cap L^{p^{i+1}}), \quad i = 0, 1, \dots$$

In an entirely similar manner as in the proof of [7, Lemma 2, p. 339] we obtain $F^* = K(F^* \cap L^p) = \dots = K(F^* \cap L^{p^i}) = \dots$. Hence

$$K \subseteq F^* = \bigcap_i K(F^* \cap L^{p^i}) \subseteq \bigcap_i K(L^{p^i}) = K,$$

i.e., $F^* = K$.

(1.2) LEMMA. *Suppose L/K is purely inseparable Galois. Let F be an intermediate field of L/K such that L/F is modular and $F \cap L^{p^n} \subseteq K$ for some nonnegative integer n . If for every maximal pure independent set M of L/K every element of M has the same exponent over F that it has over K , then $F = K$.*

PROOF. The proof is exactly the same as the proof of [7, Lemma 3, p. 340] with “maximal pure independent set” replacing “modular base” there.

(1.3) THEOREM. *Suppose $p \neq 2$. Let $K \subset F$ be Galois subfields of L . Then $\bar{H}(L/F)$ is $\bar{H}(L/K)$ -invariant if and only if $F = K(L^{p^r})$ for some nonnegative integer r .*

PROOF. If $F = K(L^{p^r})$ for some r , then $\bar{H}(L/K)$ leaves F invariant. Hence clearly $\bar{H}(L/F)$ is $\bar{H}(L/K)$ -invariant. Conversely, suppose $\bar{H}(L/F)$ is $\bar{H}(L/K)$ -invariant. We prove the theorem first for the case $p > 3$. Suppose $\bigcap_i K(F \cap L^{p^i})(L^{p^i}) = F$ for all nonnegative integers j . Then

$$K = \bigcap_j \bigcap_i K(F \cap L^{p^i})(L^{p^i}) = F,$$

a contradiction. Let j be such that $\bigcap_i K(F \cap L^{p^i})(L^{p^i}) \subset F$ and set

$$K_j = \bigcap_i K(F \cap L^{p^i})(L^{p^i}).$$

Then $\bigcap_i K_j(L^{p^i}) = K_j$ and L/K_j is modular [7, Lemma 1, p. 339], [9, Proposition 1.2(b), p. 40]. Thus K_j is Galois in L and $\bar{H}(L/F)$ is invariant in the smaller group $\bar{H}(L/K_j)$. Now F/K_j is purely inseparable of bounded exponent. By [8, Lemma 1.61(c), p. 56], \bar{F}/K_j is modular. Also $F \cap L^{p^n} \subseteq K_j$ for some n , namely $n = j$. Hence $F \cap \bar{F}^{p^n} \subseteq K_j$. By Lemma 1.2, there exists a maximal pure independent set X of \bar{F}/K_j with $x \in X$ such that the exponent t of x over F is less than the exponent s of x over K_j . Let Y be a maximal pure independent set of \bar{F}/F . Suppose \bar{F}/F is of unbounded exponent. If $F(Y)/F$ is of bounded exponent, then $\bar{F} = J \otimes_F F(Y)$ for some intermediate field J of \bar{F}/F [9, Proposition 2.6, p. 43]. Since Y is necessarily a relative p -basis of \bar{F}/F , J/F is relatively perfect. Hence $\bigcap_i F(\bar{F}^{p^i}) = J \supset F$, a contradiction. Thus $F(Y)/F$ is of unbounded exponent. Hence there exists $y \in Y$ such that $u > s$ where u is the exponent of y over F . Hence $u > s > t$. Let e be any positive integer such that $e > u$. Since L/\bar{F} is modular, L/\bar{F} is separable and thus preserves p -independence. It follows that there exists $d = \{d_0, d_1, \dots, d_{p^e}\} \in H(L/K_j)$ and $d' = \{d'_0, d'_1, \dots, d'_{p^e}\} \in H(L/F)$ with first nonzero maps of positive subscript being q and q' respectively, such that $d_q(x) = y, d'_q(y) \neq 0, q = p^{e-s} + 1, q' = p^{e-u} + 1$.

Since $\bar{H}(L/F)$ is $\bar{H}(L/K_j)$ -invariant, $d^{-1}d'd$ restricted to F must be the identity higher derivation, i.e. $d'd = d$ when restricted to F . Suppose $(q + q')p^t \leq p^e$. Then

$$\begin{aligned} (d'd)_{(q+q')p^t}(x^{p^t}) &= \sum \{d'_i d_{(q+q')p^t-i}(x^{p^t}) \mid 0 \leq i \leq (q + q')p^t\} \\ &= \sum \{d'_j (d_{q+q'-j}(x))^{p^t} \mid 0 \leq j \leq q + q'\} \\ &= d_{(q+q')p^t}(x^{p^t}) + d'_{q'}(y)^{p^t} \\ &\neq d_{(q+q')p^t}(x^{p^t}), \text{ a contradiction.} \end{aligned}$$

Thus $(q + q')p^t > p^e$, so $p^{e-s} + p^{e-u} + 2 > p^{e-t}$. Hence $p^{-s} + p^{-u} + 2p^{-e} > p^{-t}$. Since we can take e as large as we wish, we have $p^{-s} + p^{-u} \geq p^{-t}$ so $p^{t-s} + p^{t-u} \geq 1$. Since $s - t \geq 1$ and $u - t \geq 2$, we have $p^{-1} + p^{-1} > p^{t-s} + p^{t-u}$, i.e., $2 > p$, a contradiction. Thus \bar{F}/F has bounded exponent so L/K_j has finite inseparability exponent. Suppose $\bar{F} \subset L$. Then as in the proof of [2, Theorem 8], we obtain $F = K_j$ a contradiction. Hence $\bar{F} = L$.

Thus L/F has bounded exponent so $L \supseteq F \supseteq K(L^{p^n})$ for some n . Now $\bar{H}(L/F)$ is $\bar{H}(L/K(L^{p^n}))$ -invariant. Hence $F = K(L^{p^r})$ for some r by [2, Theorem 8].

The proof for the case $p = 3$ is exactly the same, once it is noted that [2, Theorem 8] is true for $p = 3$. This follows from [1, Theorem, p. 277] and in particular [1, Lemma, p. 278]. Here, for large e the key inequality becomes $2p^{e-1} + 2p^{e-1} > p^e$. Since t_1 is fixed, large e force $p = 2$.

2. Galois subfields. Let L be a Galois extension of K , i.e., L/K is modular and $\cap_i K(L^{p^i}) = K$. Then certainly $\cap_i K(\bar{K}^{p^i}) = K$ and since \bar{K} is modular over K , \bar{K} is a Galois extension of K . Moreover L/\bar{K} is separable (hence modular) so L/\bar{K} will be Galois if and only if $\cap_i \bar{K}(L^{p^i}) = \bar{K}$. We now investigate conditions which will guarantee L/\bar{K} is Galois.

(2.1) PROPOSITION. *Suppose K is a Galois subfield of L . If L/K splits, then K is Galois in L .*

PROOF. $L = S \otimes_K \bar{K}$ where S is an intermediate field of L/K which is separable over K . As noted L/\bar{K} is separable, so it suffices to show $\cap_i \bar{K}(L^{p^i}) = \bar{K}$. Now $\cap_i \bar{K}(L^{p^i}) = \cap_i (K(S^{p^i}) \otimes_K \bar{K}) = \bar{K}$.

(2.2) THEOREM. *Let K be a field. Then $[K : K^p] < \infty$ if and only if for every field extension L/K such that K is Galois in L , \bar{K} is Galois in L .*

PROOF. Suppose $[K : K^p] < \infty$. Let L/K be Galois. Then $\cap_i K(\bar{K}^{p^i}) = K$ and since any relative p -basis of \bar{K}/K is finite, we have \bar{K}/K has bounded exponent. By [5, Theorem 4, p. 1178], L/K splits and so Proposition 2.1 applies.

Conversely, suppose $[K : K^p] = \infty$. Let $x_1, x_3, \dots, x_{2n-1}, \dots$ be p -independent in K . Let

$$L = K(z, z^{p^{-1}} + x_1^{p^{-2}}, \dots, z^{p^{-n}} + x_1^{p^{-n-1}} + x_3^{p^{-n-2}} + \dots + x_{2n-1}^{p^{-2n}}, \dots)$$

where z is transcendental over K . Then


$$\bar{K} = K(x_1^{p^{-1}}, x_3^{p^{-3}}, \dots, x_{2n-1}^{p^{-2n+1}}, \dots).$$

Since L/\bar{K} is a union of ascending chain of separable extensions of \bar{K} , L/\bar{K} is separable. Now $\bar{K}(L^p) = L$ so \bar{K} is not Galois in L . Clearly \bar{K}/K is purely inseparable modular so L/K is modular [5, Theorem 1, p. 1117]. Hence in order to show K is Galois in L it suffices to show $\cap_i K(L^{p^i}) = K$. Now $\{z^{p^{-n}} + x_1^{p^{-n-1}} + \dots + x_{2n-1}^{p^{-2n}} | n = 1, 2, \dots\}$ is a subbasis of $L/K(z)$. Hence $\cap_i K(z)(L^{p^i}) = K(z)$. Let $K^* = \cap_i K(L^{p^i})$. Since $\cap_i K^*(L^{p^i}) = K^*$, K^* is separably algebraically closed in L . Clearly $K^* \subseteq K(z)$. Suppose $K^* \neq K$. Then $K(z)/K^*$ is algebraic and thus $K^* = K(z^{p^e})$ for some nonnegative integer e . Now $z^{p^e} \in K(L^{p^{2e+1}})$. Therefore

$$x_{2e+1}^{p^{-1}} \in K(L^{p^{2e+1}}) \cap \bar{K}.$$

By the separability of L/\bar{K} , the modularity of \bar{K}/K , [4, Lemma p. 162], and

the following diagram,




we have $x_{2e+1}^{p^{-1}} \in K(L^{p^{2e+1}}) \cap \bar{K} = K(\bar{K}^{p^{2e+1}})$ which is clearly impossible. Hence $K^* = K$ and K is a Galois subfield of L .

Consider the example constructed in the proof of Theorem 2.2. Heerema [3] originally developed pencils of higher derivations in order to incorporate both the finite and infinite rank higher derivation Galois theories into 1 unified theory. He considered finitely generated modular extensions L/K . In this case \bar{K} would be the field of constants of the group of infinite rank higher derivations (pencils with infinite extended rank in the new theory). However, in the example above, L/\bar{K} is relatively perfect and hence has no infinite higher derivations and yet L/K is Galois. Thus in the nonfinitely generated case a different type of field of constants can occur.

In Proposition 2.1 and Theorem 2.2 the sufficient condition given for \bar{K} to be Galois in L also imply L/K splits. We now develop an example to show that L/K and L/\bar{K} being Galois does not imply L/K splits.

(2.3) PROPOSITION. *Let F be an intermediate field of L/K such that L/F is separable Galois and F/K is Galois. Then L/K is Galois.*

PROOF. Since L/F is separable and $F/K(F^{p^i})$ is modular, $L/K(F^{p^i})$ is modular, $i = 0, 1, \dots$. Hence $L/\cap_i K(F^{p^i})$ is modular, i.e. L/K is modular [9, Proposition 1.2(b), p. 40]. Thus we have linear disjointness in the following diagram



Hence $F \cap K(L^{p^i}) = K(F^{p^i})$. Clearly $\cap_i K(L^{p^i}) \subseteq F$. Thus

$$\bigcap_i K(L^{p^i}) = \bigcap_i K(L^{p^i}) \cap F = \bigcap_i K(F^{p^i}) = K.$$

(2.4) COROLLARY. *If L/\bar{K} and \bar{K}/K are Galois, then L/K is Galois.*

PROOF. L/\bar{K} is separable since L/\bar{K} is modular and \bar{K} is algebraically closed in L .

(2.5) EXAMPLE. L/K and L/\bar{K} are Galois, yet L/K does not split: Let $K = P(z_1^p, \dots, z_j^p, \dots)$ and

$$L = K(z_1, \dots, z_j^{p^{-j+1}}, \dots)(y, u_0, \dots, u_n^{p^{-n}}, \dots)$$

where P is a perfect field, $y, u_0, z_1, \dots, z_j, \dots$ are algebraically independent indeterminants over P and $u_n^{p^{-n}} = y^{p^{-1}} + z_n^{p^{-n}} u_{n-1}^{p^{-n}}, n = 1, 2, \dots$. Then $\bar{K} = K(z_1, \dots, z_j^{p^{-j+1}}, \dots)$. Now $\bar{K}(y, u_n^{p^{-n}}), n = 0, 1, \dots$, is an ascending chain of separable extensions of \bar{K} whose union is L . Thus L/\bar{K} is separable and L/K is modular. In order to show L/K and L/\bar{K} are Galois, it suffices to show L/\bar{K} is Galois by Corollary (2.4) since \bar{K}/K is obviously Galois. Let $Z = \{z_i | i = 1, 2, \dots\}$ and $Z^{p^{-\infty}} = \{z_i^{p^{-j}} | i, j = 1, 2, \dots\}$. Let $M = L(Z^{p^{-\infty}})$. Then $P(Z^{p^{-\infty}})$ is the maximal perfect subfield of M [6, Lemma 11, p. 392]. Hence

$$\bigcap_i \bar{K}(L^{p^i}) \subseteq \bigcap_i P(Z^{p^{-\infty}})(M^{p^i}) = P(Z^{p^{-\infty}}).$$

Thus $\bigcap_i K(L^{p^i})$ is algebraic over \bar{K} and so is equal to \bar{K} . We now show L/K does not split. We first show $\bigcap_i K(y)(L^{p^i}) = K(y, u_0^p)$. Clearly $K(y, u_0^p) \subseteq \bigcap_i K(y)(L^{p^i})$. Now $\{u_0, u_1^{p^{-1}}, \dots, u_n^{p^{-n}}, \dots\}$ is a subbasis of $L/K(y, u_0^p)$. Hence

$$\bigcap_i K(y, u_0^p)(L^{p^i}) = K(y, u_0^p).$$

Thus $\bigcap_i K(y)(L^{p^i}) = K(y, u_0^p)$. Suppose L/K does split, say $L = S \otimes_K \bar{K}$, where S is an intermediate field with S/K separable. Let y have exponent t over S . Now $y^{p^t} \notin K = \bigcap_i K(S^{p^i})$. Hence there is a nonnegative integer s such that $y^{p^t} \in K(S^{p^s}), y^{p^t} \notin K(S^{p^{s+1}})$. Suppose $y^{p^{t-1}} \in \bar{K}(S^{p^s})$. Then

$$y^{p^t} \in (K(S^{p^{s+1}}) \otimes_K K(\bar{K}^p)) \cap (K(S^{p^s}) \otimes_K 1) = K(S^{p^{s+1}}),$$

a contradiction. Hence y has exponent t over $\bar{K}(S^{p^s})$. Thus $K(S^{p^s})(y)$ and $\bar{K}(S^{p^s})$ are linearly disjoint over $K(S^{p^s})$. Since $K(S^{p^s})$ and \bar{K} are linearly disjoint over $K, K(S^{p^s})(y)$ and \bar{K} are linearly disjoint over K . Since $\bar{K} \supset K^{p^{-1}}, K(S^{p^s})(y)$ is separable over K . Let $S' = K(S^{p^s})(y)$. Then $L = K(L^{p^s})(y) = S' \otimes_K \bar{K}$ and $y \in S'$. Since $\{y\}$ must be a relative p -basis of S'/K and S'/K is separable, $S'/K(y)$ is separable. Hence $L = S' \otimes_{K(y)} \bar{K}(y)$. Now

$$K(y, u_0^p) = \bigcap_i K(y)(L^{p^i}) = \bigcap_i (K(y)(S'^{p^i}) \otimes_{K(y)} K(y)(\bar{K}^{p^i})) = S'.$$

Hence $L = \bar{K}(y, u_0^p)$, a contradiction. Thus L/K does not split.

REFERENCES

1. J. Deveney and J. Mordeson, *Invariant subgroups of groups of higher derivations*, Proc. Amer. Math. Soc. **68**, (1978), 277-280.

2. _____, *On Galois theory using pencils of higher derivations*, Proc. Amer. Math. Soc. **72** (1978), 233–238.
3. N. Heerema, *Higher derivation Galois theory of fields*, Trans. Amer. Math. Soc. (to appear).
4. N. Jacobson, *Lectures in abstract algebra*. Vol. III: *Theory of fields and Galois theory*, Van Nostrand, Princeton, N. J., 1964. MR **30** #3087.
5. H. Kreimer and N. Heerema, *Modularity vs. separability for field extensions*, Canad. J. Math. **27** (1975), 1176–1182.
6. S. Mac Lane, *Modular fields*. I: *Separating transcendence bases*, Duke Math. J. **5** (1939), 372–393.
7. J. Mordeson, *On a Galois theory for inseparable field extensions*, Trans. Amer. Math. Soc. **214** (1975), 337–347. MR **52** #5635.
8. J. Mordeson and B. Vinograd, *Structure of arbitrary purely inseparable field extensions*, Lecture Notes in Math., vol. 173, Springer-Verlag, Berlin and New York, 1970. MR **43** #1952.
9. W. Waterhouse, *The structure of inseparable field extensions*, Trans. Amer. Math. Soc. **211** (1975), 39–56. MR **52** #359.

DEPARTMENT OF MATHEMATICAL SCIENCES, VIRGINIA COMMONWEALTH UNIVERSITY, RICHMOND, VIRGINIA 23284

DEPARTMENT OF MATHEMATICS, CREIGHTON UNIVERSITY, OMAHA, NEBRASKA 68134