

CONVEXITY OF THE DOMINANT EIGENVALUE OF AN ESSENTIALLY NONNEGATIVE MATRIX

JOEL E. COHEN¹

ABSTRACT. The dominant eigenvalue of a real $n \times n$ matrix A with nonnegative elements off the main diagonal is a convex function of the diagonal of A . We give a short proof using Trotter's product formula and a theorem on log-convexity due to Kingman.

An $n \times n$ real matrix A with nonnegative elements a_{ij} ($i \neq j$) off the main diagonal is called essentially nonnegative. Such an A has an eigenvalue $r(A)$, called the dominant eigenvalue, that is real and greater than or equal to the real part of any other eigenvalue of A . $r(A)$ need not be the eigenvalue of A with the largest modulus, but when every element of A is nonnegative, $r(A)$ is indeed the Perron-Frobenius root or spectral radius of A . $r(A)$ was shown to be a convex function of a single diagonal element of nonnegative A , when all other elements of A are held constant, by use of determinants [1]. $r(A)$ was shown to be a convex function of all diagonal elements jointly of essentially nonnegative A by a Feynman-Kac formula from the theory of random evolutions [2] and by a variational formula of Donsker and Varadhan [3]. Here we give a short proof that $r(A)$ is a convex function of the diagonal of essentially nonnegative A using the Trotter product formula [6] and a theorem on log-convexity due to Kingman [4] as extended by Seneta [5].

The Trotter product formula for any two real $n \times n$ matrices S and T asserts

$$e^{S+T} = \lim_{k \rightarrow \infty} (e^{S/k} e^{T/k})^k. \quad (1)$$

A positive function f on a convex domain is log-convex if $\log f$ is convex on that domain. Kingman showed that the set of log-convex functions is closed under positive linear combinations, multiplication, positive powers and $\lim \sup$. Let $T(u)$ be a nonnegative (elementwise) $n \times n$ matrix in which each element is either identically 0 or is a log-convex function of a parameter u in some convex domain in R^m , $m > 1$. Then, provided that $r(T(u))$ is positive in this domain, $r(T(u))$ is log-convex there [5, p. 83]. (Kingman [4] assumed every element of $T(u)$ positive.)

Let D be a diagonal real $n \times n$ matrix $\text{diag}(d_1, \dots, d_n)$. Let A be essentially nonnegative.

THEOREM. $r(A + D)$ is a convex function of D .

Received by the editors December 12, 1979 and, in revised form, July 30, 1980.

AMS (MOS) subject classifications (1970). Primary 15A42, 15A48.

Key words and phrases. Perron-Frobenius root, convexity, log-convexity, Trotter product formula, spectral radius.

¹Supported in part by U. S. National Science Foundation grant DEB80-11026.

© 1981 American Mathematical Society
0002-9939/81/0000-0183/\$01.50

PROOF. We have $r(A + D) = \log e^{r(A+D)} = \log r(e^{A+D})$. Since r is a continuous function, (1) gives

$$r(e^{A+D}) = \lim_{k \rightarrow \infty} r[(e^{A/k} e^{D/k})^k],$$

hence

$$r(A + D) = \lim_{k \rightarrow \infty} \log r[(e^{A/k} e^{D/k})^k]. \quad (2)$$

Now for every positive integer k , every element of $e^{A/k}$ is nonnegative. Moreover $r(e^{A/k}) = e^{r(A/k)} > 0$, and multiplying $e^{A/k}$ by a nonnegative nonsingular matrix will yield a nonnegative matrix whose spectral radius is also positive. In particular, $T_k(D) = (e^{A/k} e^{D/k})^k$ is nonnegative with $r(T_k(D)) > 0$. For $i = 1, 2, \dots, n$, the i th diagonal element of $e^{D/k}$, which is just $e^{d_i/k}$, is a log-convex function of D . Hence every element of $T_k(D)$ is either 0 or a log-convex function of D . The theorem of Kingman as extended by Seneta implies that $\log r(T_k(D))$ is a convex function of D . By (2), $r(A + D)$ is also a convex function of D .

Some applications of this theorem are discussed in [1].

I thank E. Seneta and a referee for corrections and comments.

REFERENCES

1. J. E. Cohen, *Derivatives of the spectral radius as a function of non-negative matrix elements*, Math. Proc. Cambridge Philos. Soc. **83** (1978), 183–190.
2. _____, *Random evolutions and the spectral radius of a non-negative matrix*, Math. Proc. Cambridge Philos. Soc. **86** (1979), 345–350.
3. S. Friedland, *Convex spectral functions*, Linear and Multilinear Algebra (to appear).
4. J. F. C. Kingman, *A convexity property of positive matrices*, Quart. J. Math. Oxford Ser. (2) **12** (1961), 283–284.
5. E. Seneta, *Non-negative matrices: an introduction to theory and applications*, George Allen and Unwin, London, 1973.
6. H. F. Trotter, *On the product of semigroups of operators*, Proc. Amer. Math. Soc. **10** (1959), 545–551.

DEPARTMENT OF POPULATIONS, ROCKEFELLER UNIVERSITY, NEW YORK, NEW YORK 10021