

A GROUP THEORETICAL EQUIVALENT OF THE ZERO DIVISOR PROBLEM

A. I. LICHTMAN¹

ABSTRACT. We prove that the zero divisor problem is equivalent to some problem about subgroups of a free group.

1. Introduction. Let G be an arbitrary torsion free group. The problem of whether the group ring KG contains zero divisor is open, although positive solution has been obtained for some important special classes of groups, in particular, for polycyclic-by-finite-groups (Farkas and Snider [1], Cliff [2]) and for one-relator groups (J. Lewin and T. Lewin [3]). In this note we show that in the cases when K is the ring of integers Z or the field Z_p of residue classes modulo p the zero divisor problem for KG is equivalent to a problem about subgroups of a free group.

Let $G = F/N$ be a presentation of G as a quotient group of a free group F ; as usual $\gamma_k(N)$ ($k = 1, 2, \dots$) denotes the k th term of the lower central series of N . If $1 \neq x \in N$ then there exists a maximal number k such that $x \in \gamma_k(N)$; we can say too that $x \in \gamma_k(N) \setminus \gamma_{k+1}(N)$.

THEOREM 1. *The ring ZG contains no zero divisors if for every system of elements $x, f_1, f_2, \dots, f_n \in F$, where $x \in \gamma_k(N) \setminus \gamma_{k+1}(N)$ and is not a power in $\gamma_k(N)$ (and hence not in N) and f_i ($i = 1, 2, \dots, n$) are taken from different cosets of G/N , the subgroup N_1 , generated by the elements*

$$(1) \quad x_i = f_i^{-1} x f_i \quad (i = 1, 2, \dots, n)$$

satisfies the condition

$$(2) \quad N_1 \cap \gamma_{k+1}(N) = N'_1.$$

The inverse assertion is true without the restriction that x is not a power in $\gamma_k(N) \setminus \gamma_{k+1}(N)$.

PROPOSITION 1. *Assume that ZG contains no zero divisors. Then for every element $x \in \gamma_k(N) \setminus \gamma_{k+1}(N)$ and f_1, f_2, \dots, f_n taken from different cosets of F/N , the elements (1) are free generators of the subgroup N_1 and (2) holds.*

REMARK. Since $x \in \gamma_k(N)$ we have $N_1 \subseteq \gamma_k(N)$ and thus $N'_1 \subseteq \gamma_{k+1}(N)$, and hence

$$N_1 \cap \gamma_{k+1}(N) \supseteq N'_1.$$

Thus, the real sense of condition (2) is that $N_1 \cap \gamma_{k+1}(N)$ does not properly include N'_1 . \square

When the element x in Theorem 1 does not belong to N' , i.e., $k = 1$, we immediately obtain the following corollary.

Received by the editors April 25, 1985 and, in revised form, June 11, 1985.

1980 *Mathematics Subject Classification.* Primary 16A27, 20C07.

¹Partially supported by NSF Grant No. D14S-8400445.

COROLLARY. *The ring ZG contains no zero divisors iff*

$$(3) \quad N_1 \cap N' = N'_1,$$

for all choices of x, f_1, f_2, \dots, f_n as described.

It is worth remarking that our group theoretical equivalent of the zero divisor problem is obtained for group rings over Z or Z_p ; the connection between the zero divisor problem over Z or Z_p and over an arbitrary field K is unknown.

2. Let F be a free group, $N \triangleleft F$ and $G = F/N$. It is easy to see that the free abelian group N/N' becomes a ZG -module via the conjugation in F ; this is, the so-called relation module which has been studied intensively during the recent years. We refer the reader to K. Gruenberg's lecture notes [4] for the detailed definition and properties of these modules.

For $k > 1$ the factors $\gamma_k(N)/\gamma_{k+1}(N)$ become ZG -modules in the same way. We need the following property of these higher relation modules:

LEMMA 1. *For every natural k the ZG -module $\gamma_k(N)/\gamma_{k+1}(N)$ can be embedded in a free module.*

PROOF. The result is known and is proven even for a more general type of module: the higher relative relation module (see [5]). We give, therefore, only the outline of the proof.

When $k = 1$, the embedding of the relation module N/N' into a free module M is described in [4, lecture 2]. It is based on the existence of the exact sequence of ZG -modules

$$(4) \quad 0 \rightarrow N/N' \rightarrow M \rightarrow \omega(ZG) \rightarrow 0,$$

where M is a free ZG -module and $\omega(ZG)$ is the augmentation ideal of ZG . When $k > 1$ the assertion follows from the observation that $\gamma_k(N)/\gamma_{k+1}(N)$ is embedded into the k th tensor power

$$M^{(k)} = \underbrace{M \otimes_Z M \otimes \cdots \otimes_Z M}_k.$$

Clearly, $M^{(k)}$ is a free ZG -module.

LEMMA 2. *Let R be a prime ring, and let M be a left nonzero R -module which can be embedded into a free module. Then R contains no zero divisors iff for any elements $r \in R$, $m \in M$, the condition*

$$(5) \quad rm = 0$$

implies necessarily that $r = 0$ or $m = 0$.

PROOF. Let $0 \neq r \in R$. If (5) holds, then r must annihilate all the coordinates of m and hence, in the case when R contains no zero divisors, we obtain that $m = 0$. Conversely, let x, y be two nonzero elements in R such that $xy = 0$.

Consider the subset $yM \subseteq M$. Clearly, $x(yM) = 0$ and it remains to prove that $yM \neq 0$. But if $yM = 0$ we obtain a nonzero two sided ideal $A = (y)$ such that

$$(6) \quad AM = 0.$$

Now let M be embedded into a free R -module with a basis e_i ($i \in I$). The i th coordinates of elements from M form a left ideal $B_i \subseteq R$ ($i \in I$) and we obtain from (6) that

$$(7) \quad AB_i = 0 \quad (i \in I).$$

Since $M \neq 0$ at least one of these left ideals B_i is not zero and we conclude easily via the assumption that R is prime that (7) is impossible, i.e., $yM \neq 0$ and the assertion follows.

3. Proofs of Theorem 1 and Proposition 1. Since G is torsion free, ZG must be prime (see [6]). Lemma 2 implies that ZG contains zero divisors iff there exist nonzero

$$r = \sum_{i=1}^n \alpha_i g_i \in ZG \quad \text{and} \quad \bar{x} \in M = \gamma_k(N)/\gamma_{k+1}(N)$$

such that

$$(8) \quad \sum_{i=1}^n \alpha_i (g_i \bar{x}) = 0,$$

or, in other words, ZG contains zero divisors iff there exist elements $g_i \in G$ ($i = 1, \dots, n$) and $0 \neq \bar{x} \in M$ such that elements $g_i \bar{x}$ are linearly dependent over Z . Pick an arbitrary coset representative f_i in the different cosets $\bar{f}_i = f_i N$ ($i = 1, 2, \dots, n$) and x in $\bar{x} = x\gamma_{k+1}(N)$. We can formulate the last condition:

(*) *The elements $f_i^{-1} x f_i$ ($i = 1, 2, \dots, n$) of $\gamma_k(N)$ are linearly dependent modulo $\gamma_{k+1}(N)$.*

We now prove Proposition 1. Since ZG is a domain the condition (*) implies that the elements $f_i^{-1} x f_i$ ($i = 1, 2, \dots, n$), which generate N_1 , are linearly independent modulo $N_1 \cap \gamma_{k+1}(N)$ and hence are linearly independent modulo N'_1 . We therefore conclude that these elements freely generate N_1 . It is easy to see now that their linear independence modulo $N_1 \cap \gamma_{k+1}(N) \supseteq N'_1$ implies (2) and the proof of Proposition 1 is completed.

We prove now Theorem 1. If ZG contains zero divisors then (*) holds for some $x \in \gamma_k(N)$. Furthermore, we can assume that x is not a power in $\gamma_k(N)$. Indeed, this is obvious if \bar{x} is not a power in the free abelian group $\bar{M} = \gamma_k(N)/\gamma_{k+1}(N)$; but if there exists $\bar{y} \in M$ such that $x = n\bar{y}$ then we can replace \bar{x} by \bar{y} in (8) and then replace x by y in (*).

We now observe that the quotient group $F/\gamma_k(N)$ is torsion free: when $k = 1$ it coincides with the torsion free group G ; when $k > 1$ this fact follows for instance from [7]. We can therefore conclude that x is not a power in F .

Now let T be the normal subgroup of F , generated by the element x . Clearly, $T \subseteq N$; hence the elements f_i ($i = 1, 2, \dots, n$) belong to different cosets of G/T . The quotient group G/T is a torsion-free one-relator group and we obtain from Lyndon's Identity Theorem (see [8]) that the elements (1) are linearly independent in T/T' . But the subgroup N_1 , generated by these elements, belongs to T ; this implies easily that the elements (1) are linearly independent in $N_1/N_1 \cap T'$ and hence in N_1/N'_1 . Once again, condition (*) implies that zero divisors in ZG exist iff the elements (1) are linearly dependent in $N_1/N_1 \cap \gamma_{k+1}(N)$, i.e., iff $N_1 \cap \gamma_{k+1}(N)$ properly contains N'_1 and the assertion follows.

4. To obtain an analog of Theorem 1 for the ring Z_pG we have to consider instead of the lower central series the p -series $M_j(N)$ ($j = 1, 2, \dots$) of the dimension subgroups of Z_pN (see [6, Chapter 11]). We remind the reader that $M_1(N) = N$, $M_2(N) = N'N^p$ (the product of the commutator subgroup and the subgroup N^p , generated by the p th powers of all the element of N), that all the factors $M_j(N)/M_{j+1}(N)$ are abelian groups of exponent p , and that

$$(9) \quad \bigcap_{j=1}^{\infty} M_j(N) = 1.$$

Once again, the quotient group $M_j(N)/M_{j+1}(N)$ becomes a Z_pG -module via the conjugation in F .

LEMMA 3. For every given j the Z_pG -module $M_j(N)/M_{j+1}(N)$ can be embedded into a free module.

PROOF. The module $M_1(N)/M_2(N) = N/N'N^p$ is isomorphic to the Z_pG -module $\bar{N}/(p\bar{N})$, where \bar{N} is the relation module N/N' . This implies via (4) the existence of the exact sequence of Z_pG -modules

$$(4') \quad 0 \rightarrow M_1(N)/M_2(N) \rightarrow M/pM \rightarrow \omega(Z_pG) \rightarrow 0.$$

Once again, $M_j(N)/M_{j+1}(N)$ is embedded into the j th tensor power of the free Z_pG -module M/pM . \square

THEOREM 2. The ring Z_pG contains no zero divisors iff for every system of elements $x, f_1, f_2, \dots, f_n \in F$, where $x \in M_j(N) \setminus M_{j+1}(N)$ and is not a power in N and f_i ($i = 1, 2, \dots, n$) are taken from different cosets of G/N , the subgroup N_1 , generated by the elements (1), satisfies the condition

$$(2') \quad N_1 \cap M_{j+1}(N) = M_2(N_1).$$

REMARK 1. It is worth remarking that the inclusion $N_1 \cap M_{j+1}(N) \supseteq M_2(N_1)$ follows immediately from the condition $x \in M_j(N)$.

REMARK 2. If ZG contains no zero divisors then (2') holds without the assumption that x is not a power in N . The proof of this fact is similar to the proof of Proposition 1 and we omit it.

LEMMA 4. Let H be a free group, and let U be a normal subgroup of finite index. Then every coset hU ($h \in H$) contains an element which is not a proper power in H .

PROOF. Let h_i ($i \in I$) be a free system of generators for H and let

$$h = h_{i_1}^{\alpha_1} h_{i_2}^{\alpha_2} \dots h_{i_k}^{\alpha_k};$$

pick n such that $h_{i_1}^n \in H$ and $\alpha_1 + n > \alpha_i$ ($i = 1, 2, \dots, n-1$). It is easy to verify that the element

$$h_{i_1}^{(n+\alpha_1)} h_{i_2}^{\alpha_2} \dots h_{i_k}^{\alpha_k} = h_{i_1}^n h$$

is not a power in H .

PROOF OF THEOREM 2. We only give the outline of the proof since the argument is essentially the same as in Theorem 1.

First, find $0 \neq \bar{x} \in M_j(N)/M_{j+1}(N)$ such that (8) holds (with coefficients $\alpha_i \in Z_p$). Then find x in the coset \bar{x} such that x is not a proper power in N , and hence in F . Then, similarly to condition (*), obtain that the elements (1) of $M_j(N)$ are linearly dependent modulo $M_{j+1}(N)$.

On the other hand, the elements (1) are linearly independent in $T/T'T^p$ and hence in $N_1/N_1'N^p$; thus, there exist zero divisors in Z_pG iff $N_1 \cap M_{j+1}(N)$ properly contains $M_2(N_1)$.

COROLLARY. *Let $x \in N \setminus N'N^p$ and f_1, f_2, \dots, f_k be as in Theorem 2. Then Z_pG contains no zero divisors iff $N_1 \cap N'N^p = N_1'N_1^p$.*

ACKNOWLEDGMENT. I am grateful to the referee whose remarks helped me to improve the exposition of this paper.

REFERENCES

1. D. Farkas and R. L. Snider, *K_0 and Noetherian group rings*, J. Algebra **42** (1976), 192–198.
2. G. H. Cliff, *Zero divisors and idempotents in group rings*, Canad. J. Math. **32** (1980), 596–602.
3. J. Lewin and T. Lewin, *An embedding of the group algebra of a torsion free 1-relator group in a field*, J. Algebra **52** (1978), 39–77.
4. K. W. Gruenberg, *Relation modules of finite groups*, CBMS Regional Conf. Ser. in Math., no. 25, Amer. Math. Soc., Providence, R.I., 1976.
5. B. Hartley and A. I. Lichtman, *Relative higher relation modules*, J. Pure Appl. Algebra **22** (1981), 75–89.
6. D. S. Passman, *The algebraic structure of group rings*, Wiley, New York, 1977.
7. G. Baumslag, *Wreath products and extensions*, Math. Z. **81** (1963), 286–299.
8. K. W. Gruenberg, *Cohomological topics in group theory*, Lecture Notes in Math., vol. 143, Springer-Verlag, Berlin and New York, 1970.

DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA, ISRAEL

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-PARKSIDE, KENOSHA, WISCONSIN 53141 (Current address)