

SUBSPACES OF SMALL CODIMENSION OF FINITE-DIMENSIONAL BANACH SPACES

ALAIN PAJOR AND NICOLE TOMCZAK-JAEGERMANN¹

ABSTRACT. Given a finite-dimensional Banach space E and a Euclidean norm on E , we study relations between the norm and the Euclidean norm on subspaces of E of small codimension. Then for an operator taking values in a Hilbert space, we deduce an inequality for entropy numbers of the operator and its dual.

In this note we study the following problem: given an n -dimensional Banach space E and a Euclidean norm $\|\cdot\|_2$ on E and $0 < \lambda < 1$, find a subspace F of E with $\dim F \geq \lambda n$ such that

$$(*) \quad \|x\|_2 \leq M_* f(1 - \lambda) \|x\| \quad \text{for } x \in F.$$

Here M_* denotes the Levy mean of the dual norm of E (see the notation below).

This problem was considered by V. Milman, who proved in [18] that estimate (*) holds for a certain exponential function f . The estimate was improved later in [10] to $f(1 - \lambda) \leq K/(1 - \lambda)$, where K is a universal constant. The latter result turned out to be important for various applications (cf. [1, 15, 11, 19]).

The main result of this note proves (*) with the function $f(1 - \lambda) \leq K/\sqrt{1 - \lambda}$. This estimate, besides being optimal (up to a logarithmic factor), can be used to compare entropy numbers of an operator and its dual for operators taking values in a Hilbert space.

Let us recall some notation.

Let E be an n -dimensional Banach space; i.e., $E = (R^n, \|\cdot\|)$. Let $[\cdot, \cdot]$ be an inner product on R^n , and let $\|\cdot\|$ be the associated Euclidean norm on R^n defined by $\|x\| = [x, x]^{1/2}$, for $x \in R^n$.

Let B_E be the closed unit ball in E . Set

$$(1) \quad \|x\|_* = \sup\{|[x, y]| \mid y \in B_E\} \quad \text{for } x \in R^n.$$

Clearly, $(R^n, \|\cdot\|_*)$ can be identified with the dual space E^* . Let $S = \{x \in R^n \mid \|x\| = 1\}$, and let μ be the normalized rotation invariant measure on S . Define the Levy means M and M_* by

$$M = \left(\int_S \|x\|^2 d\mu \right)^{1/2}, \quad M_* = \left(\int_S \|x\|_*^2 d\mu \right)^{1/2}.$$

We shall employ a similar notation in a context of symmetric convex bodies. For a closed symmetric convex body $V \subset R^n$, by V^* we denote the dual body defined by $V^* = \{x \in R^n \mid |[x, y]| \leq 1 \text{ for all } y \in V\}$.

Received by the editors May 29, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 46B20; Secondary 47B10.

¹Research partially supported by NSERC Grant A8854.

Note that if $\|\cdot\|_V$ is the norm associated to V (i.e., $V = \{x \in \mathbb{R}^n \mid \|x\|_V \leq 1\}$), then the dual norm $\|\cdot\|_{V^*} = (\|\cdot\|_V)^*$, as defined by (1), is associated to V^* . We set $E_V = (\mathbb{R}^n, \|\cdot\|_V)$ and $E_{V^*} = (\mathbb{R}^n, \|\cdot\|_{V^*})$, and we denote by M_V and M_{V^*} the corresponding Levy means.

The main result of this note is

THEOREM 1. *Let $E = (\mathbb{R}^n, \|\cdot\|)$ and let $\|\cdot\|$ be a Euclidean norm on \mathbb{R}^n . For every $0 < \lambda < 1$ there exists a subspace $F \subset \mathbb{R}^n$ with $\dim F > \lambda n$ such that*

$$\|\|x\|\| \leq KM_*(1-\lambda)^{-1/2}\|x\| \quad \text{for } x \in F,$$

where K is a universal constant.

In particular, we have a corollary for finite-dimensional subspaces of a Banach space of cotype 2. It improves earlier results from [4, 10, 18] (cf. also [11]).

COROLLARY 1. *Let X be a Banach space of cotype 2 and let $E \subset X$ be an n -dimensional subspace. For every $0 < \lambda < 1$ there exists $F \subset E$ with $m = \dim F > \lambda n$ such that*

$$d(F, l_2^m) \leq KC_2(X)(1-\lambda)^{-1/2} \log[C_2(X)(1-\lambda)^{-1/2} + 1],$$

where $C_2(X)$ is the Gaussian cotype 2 constant of X and K is a universal constant.

SKETCH OF THE PROOF. It can be shown, using results of [15 and 16] (cf. e.g. [1, 4, 10, 11, 14]), that the Euclidean norm $\|\cdot\|$ induced on E by the ellipsoid of maximal volume contained in B_E satisfies $\|x\| \leq \|\|x\|\|$ for $x \in E$ and $M_* \leq K'C_2(X) \log[d(E, l_2^n) + 1]$. Combining this fact either with Milman's iteration procedure (cf., e.g., [1, 4, 10]) or with an argument from [10] (proof of Theorem 1 and Remark after Theorem 8), yields the result. \square

The estimate obtained in Theorem 1 and Corollary 1 is, up to a logarithmic factor, the best possible. If $F \subset l_1^n$ is an m -dimensional subspace (with $m > \lambda n$), considering a projection $P: l_1^n \rightarrow l_1^n$ with $\ker P = F$ and $\gamma_2(P) \leq \sqrt{n-m}$, one can easily show that $d(F, l_2^m) \geq c(1-\lambda)^{-1/2}$, where $c > 0$ is an absolute constant.

Theorem 1 can be used to obtain some new estimates between various s -numbers of operators from a Banach space X into l_2 . For this let us recall some more notation (cf., e.g., [12]).

For any compact metric space (T, d) , we denote by $N(T, d, \varepsilon)$ the smallest number of open balls of radius ε which cover T . If T is the unit ball in an n -dimensional Banach space then $N(T, \|\cdot\|, \varepsilon) \leq (1+2/\varepsilon)^n$ (cf., e.g., [6, Lemma 2.4]). Let X and Y be Banach spaces and let B_X denote the closed unit ball in X . For a compact operator $u: X \rightarrow Y$ between Banach spaces we define the k th entropy number of n by

$$e_k(u) = \inf\{\varepsilon > 0 \mid N(u(B_X), \|\cdot\|_Y, \varepsilon) \leq 2^k\}.$$

Moreover, we define the k th Gelfand number of u by

$$c_k(u) = \inf\{\|u|_Z\| \mid Z \subset X, \text{codim } Z < k\}$$

and the k th Kolmogorov number by

$$d_k(u) = \inf_{\substack{L \subset Y \\ \dim L < K}} \sup_{x \in B_X} \inf_{y \in L} \|ux - y\|.$$

It is well known (cf., e.g., [12]) that

$$(2) \quad c_k(u) = d_k(u^*).$$

For the operator $u: l_2^n \rightarrow X$, we define

$$l(u) = \left(\int_{R^n} \|ux\|^2 d\gamma_n(x) \right)^{1/2},$$

where γ_n denotes the canonical (normalized) Gaussian measure on R^n (cf., e.g., [6]). For any bounded operator $u: l_2 \rightarrow X$ we set

$$l(u) = \sup\{l(uv) | v: l_2^n \rightarrow l_2, v \leq 1, n \in \mathbf{N}\}.$$

If $\dim E = n$ and $u: l_2^n \rightarrow E$ is one-to-one, then $l(u) = \sqrt{n}M_V$, where $V = u^{-1}(B_E)$.

Now we have

PROPOSITION 1. *Let X be a Banach space and let $u: X \rightarrow l_2$ be a compact operator. Then*

$$\sup_k \sqrt{k}c_k(u) \leq Kl(u^*),$$

where K is a universal constant.

PROOF. Approximating a given operator by a finite-rank operator and replacing u by $\bar{u}: X/\ker u \rightarrow u(X)$, we may assume that $u: X \rightarrow l_2^n$ is one-to-one. Define $\|x\| = \|u^{-1}x\|_X$ for $x \in R^n$. The conclusion follows from Theorem 1 applied for $E = (R^n, \|\cdot\|)$, $\|\|\cdot\|\|$ the l_2 -norm on R^n and $\lambda = 1 - k/n$. \square

The next theorem compares entropy numbers of operators u and u^* for $u: X \rightarrow l_2$. Let us recall that it is still an open problem whether for every compact operator $u: X \rightarrow Y$, $e_k(u)$ and $e_k(u^*)$ are asymptotically of the same order, as $k \rightarrow \infty$. For more details on this and related problems and results, cf. [2, 3, 8].

THEOREM 2. *Let X be a Banach space and let $u: X \rightarrow l_2$ be a compact operator. Then*

$$\sup_k \sqrt{k}e_k(u^*) \leq K \sum_{k=1}^{\infty} \frac{e_k(u)}{\sqrt{k}},$$

where K is a universal constant.

The proof of the theorem is based upon two lemmas on entropy numbers. Lemma 1 is an operator version of results on Gaussian processes: the majorization theorem is due to Dudley [5], the minorization theorem to Sudakov [13]. A direct proof of Lemma 1 can be also found in [9].

LEMMA 1. *Let X be a Banach space and let $u: X \rightarrow l_2$ be a compact operator. Then*

$$(i) \quad l(u^*) \leq K \sum_{k=1}^{\infty} e_k(u)/\sqrt{k},$$

$$(ii) \quad \sup_k \sqrt{k}e_k(u) \leq Kl(u^*),$$

where $K \geq 1$ is a universal constant.

The next result is due to Carl [2].

LEMMA 2. Let $0 < p < \infty$. Let X and Y be Banach spaces and let $v: X \rightarrow Y$ be an operator. Then

$$\sup_{k \leq n} k^{1/p} e_k(v) \leq b_p \sup_{k \leq n} k^{1/p} d_k(v) \quad \text{for } n = 1, 2, \dots,$$

where b_p depends only on p .

Now Theorem 2 follows immediately from Lemma 2, formula (2), Proposition 1 and Lemma 1(i).

Our proof of Theorem 1 is based upon a similar idea to [7]. It will be a consequence of the following technical result.

PROPOSITION 2. There exists $0 < d < 2^{-7/2}$ such that the following is true. Let $||| \cdot |||$ be a Euclidean norm on R^n and let B_2 be the closed unit ball. Let $1 \leq j \leq n$. Let V be a closed symmetric convex body such that

$$(3) \quad V \subset \bigcup_{z \in \Lambda} (z + d|||z|||W) \cup cB_2,$$

with a set $\Lambda \subset R^n \setminus \{0\}$ with $|\Lambda| \leq e^{d^2 j}$ and some closed symmetric convex body W such that $W \subset B_2$ and $M_{W^*} \leq \sqrt{j/n}$, and for some constant c . Then there exists a subspace $F \subset R^n$ with $\dim F \geq n - j$ such that

$$(4) \quad |||x||| \leq c\|x\|_V \quad \text{for } x \in F.$$

PROOF. Applying an affine transformation, if necessary, we may assume that $|||x||| = (\sum_{i=1}^n x_i^2)^{1/2}$ for $x \in R^n$. Denote by M_j the median of the function $\psi_j(x) = (\sum_{i=1}^j x_i^2)^{1/2}$ on S . It is well known (cf., e.g., [6]) that $M_j \geq a\sqrt{j/n}$, where $0 < a < \sqrt{2}$ is a universal constant. Set $d = 2^{-4}a$. Let $\{e_i\}$ be the standard unit vector basis in R^n . Let $P_j: R^n \rightarrow R^n$ denote the orthogonal projection onto the subspace spanned by the first j unit vectors.

We need two lemmas. In their formulation $O(n)$ denotes the group of isometries of l_2^n and \mathcal{P} is the Haar measure on $O(n)$. For an operator $T: R^n \rightarrow R^n$, $\|T: E \rightarrow \bar{F}\|$ denotes the norm of T as an operator from E into F .

LEMMA 3. Let

$$A_1 = \{U \in O(n) \mid \|P_j U: E_W \rightarrow l_2^n\| > 8M_j/a\}.$$

Then $\mathcal{P}(A_1) < \frac{1}{2}$.

LEMMA 4. Let

$$A_2 = \{U \in O(n) \mid \exists z \in \Lambda \mid |||P_j U z||| < \frac{3}{4}M_j|||z|||\}.$$

Then $\mathcal{P}(A_2) < \frac{1}{2}$.

Assuming the truth of Lemmas 3 and 4 we complete the proof as follows. Pick an isometry $U_0: l_2^n \rightarrow l_2^n$ such that

$$\|P_j U_0: E_W \rightarrow l_2^n\| \leq 8M_j/a, \quad |||P_j U_0 z||| \geq (3M_j/4)|||z||| \quad \text{for all } z \in \Lambda.$$

Then

$$(5) \quad (\ker P_j U_0) \cap (z + d|||z|||W) = \emptyset \quad \text{for all } z \in \Lambda.$$

Indeed, if $y \in (\ker P_j U_0) \cap (z + d\|z\|W)$, then

$$\begin{aligned} 0 &= \|P_j U_0 y\| \geq \|P_j U_0 z\| - \|P_j U_0(y - z)\| \\ &\geq (3M_j/4)\|z\| - (8M_j/a)\|y - z\|_W \geq (3M_j/4)\|z\| - (8M_j/a)d\|z\| > 0, \end{aligned}$$

giving a contradiction. Set $F = \ker P_j U_0$. From (5) and (3) it follows that $F \cap V \subset cB_2$, which is equivalent to (4). \square

PROOF OF LEMMA 3. Let $\tilde{\Lambda}$ be a $\frac{1}{2}$ -net in $B_2 \cap [e_1, \dots, e_j]$ with minimal cardinality. Then $|\tilde{\Lambda}| = N(B_2 \cap [e_1, \dots, e_j], \|\cdot\|, \frac{1}{2}) \leq 5^j$. Thus

$$\begin{aligned} \mathcal{P}(A_1) &= \mathcal{P}\{U \in O(n) \mid \|U: l_2^j \rightarrow E_{W^*}\| > 8M_j/a\} \\ &\leq \mathcal{P}\{U \in O(n) \mid \exists z \in \tilde{\Lambda} \mid \|Uz\|_{W^*} > 4M_j/a\} \\ &\leq 5^j \mu\{y \in S \mid \|y\|_{W^*} > 4M_j/a\}. \end{aligned}$$

By the isoperimetric inequality (cf. [6, (2.6)]), the measure of the latter set is smaller than or equal to

$$4 \exp(-n(4M_j/a - M_{W^*})^2/2) \leq 4 \exp(-9j/2).$$

Therefore

$$\mathcal{P}(A_1) \leq 4 \cdot 5^j \exp(-9j/2) < 1/2. \quad \square$$

Lemma 4 follows immediately from the isoperimetric inequality and the definition of d .

Now we are ready to prove Theorem 1.

PROOF OF THEOREM 1. Fix $0 < \lambda < 1$. Let j be the smaller integer larger than or equal to $(1 - \lambda)n$. Set $V = (d_1 \sqrt{1 - \lambda}/M_*)B_E$, where a constant $0 < d_1 < 1$ will be defined later, $W = (\sqrt{1 - \lambda}/M_*)B_E \cap B_2$. Since $W \subset (\sqrt{1 - \lambda}/M_*)B_E$, the dual bodies satisfy the opposite inclusion and $(M_*/\sqrt{1 - \lambda})B_{E^*} \subset W^*$. Thus $M_{W^*} \leq (\sqrt{1 - \lambda}/M_*)M_* \leq \sqrt{j/n}$. Notice also that $M_{V^*} = (d_1 \sqrt{1 - \lambda}/M_*)M_* = d_1 \sqrt{1 - \lambda}$.

Consider the compact metric space $(V, \|\cdot\|)$. Let Λ be a $1/2$ -net in V with minimal cardinality. It follows from Lemma 1(ii), applied to the formal identity operator $u: (R^n, V) \rightarrow (R^n, \|\cdot\|)$ that

$$|\Lambda| = N(V, \|\cdot\|, \varepsilon) \leq 2^{4K^2 n M_{V^*}^2} \leq \exp(d^2 j),$$

where $K \geq 1$ is the constant from Lemma 1 and $d_1 = d/2K\sqrt{\log 2}$.

To show (3) fix $x \in V$. Pick $z \in \Lambda$ such that $\|x - z\| < 1/2$. If $\|x\| \geq 1/K\sqrt{\log 2} + 1/2d + 1/2$, then $\|z\| \geq 1/K\sqrt{\log 2} + 1/2d$. Therefore,

$$\begin{aligned} x - z &\in 2V = (2d_1 \sqrt{1 - \lambda}/M_*)B_E \subset d\|z\|(\sqrt{1 - \lambda}/M_*)B_E, \\ x - z &\in \frac{1}{2}B_2 \subset d\|z\|B_2. \end{aligned}$$

Hence $x - z \in d\|z\|W$. This shows (3) with

$$c = 1/K\sqrt{\log 2} + 1/2d + 1/2.$$

Now Proposition 2 implies that there exists $F \subset R^n$ with $\dim F \geq n - j > \lambda n - 1$ such that

$$\|x\| \leq c\|x\|_V = K'M_*(1 - \lambda)^{-1/2}\|x\| \quad \text{for } x \in F,$$

where $K' = 2cK\sqrt{\log 2}/d$. \square

REFERENCES

1. J. Bourgain and V. D. Milman, *Sections euclidiennes et volume des corps convexes symétriques*, C.R. Acad. Sci. Paris Sér. A I Math. **300** (1985), 435–438.
2. B. Carl, *Entropy numbers, s -numbers and eigenvalue problem*, J. Funct. Anal. **41** (1981), 290–306.
3. ———, *On Gelfand, Kolmogonov and entropy numbers of operators acting between special Banach spaces* (to appear).
4. S. Dilworth and S. Szarek, *The cotype constant and almost Euclidean decomposition of finite dimensional normed spaces* (to appear).
5. R. J. Dudley, *The size of compact subsets of Hilbert space and continuity of Gaussian processes*, J. Funct. Anal. **1** (1967), 290–330.
6. T. Figiel, J. Lindenstrauss and V. D. Milman, *The dimensions of almost spherical sections of convex bodies*, Acta Math. **139** (1977), 53–94.
7. E. D. Gluskin, *Norms of random matrices and diameters of finite dimensional sets*, Mat. Sb. **120** (1983), 180–189. (Russian)
8. Y. Gordon, H. König and C. Schütt, *Geometric and probabilistic estimates for entropy and approximation numbers of operators*, J. Approx. Theory (to appear).
9. T. Kühn, *γ -radonifying operators and entropy ideals*, Math. Nachr. **107**(1982), 53–58.
10. V. D. Milman, *Random subspaces of proportional dimension of finite dimensional normed spaces; approach through the isoperimetric inequality*, Séminaire d'Analyse Fonctionnelle 84/85, Université Paris VI et VII, Paris.
11. V. D. Milman and G. Pisier, *Banach spaces with a weak cotype 2 property*, Israel J. Math. (to appear).
12. A. Pietsch, *Operator ideals*, North-Holland, Berlin, 1978.
13. V. N. Sudakov, *Gaussian random processes and measures of solid angles in Hilbert space*, Soviet Math. Dokl. **12** (1971), 412–415.
14. A. Pajor and N. Tomczak-Jaegermann, *Nombres de Gelfand et sections euclidiennes de grande dimension*, Séminaire d'Analyse Fonctionnelle 84/85, Université Paris VI et VII, Paris.
15. J. Bourgain and V. D. Milman, *On Mahler's conjecture on the volume of a convex symmetric body and its polar*, Preprint, I.H.E.S.
16. W. J. Davis, V. D. Milman and N. Tomczak-Jaegermann, *The distance between certain n -dimensional spaces*, Israel J. Math. **39** (1981), 1–15.
17. T. Figiel and N. Tomczak-Jaegermann, *Projections onto Hilbertian subspaces of Banach spaces*, Israel J. Math. **33** (1979), 155–171.
18. V. D. Milman, *Volume approach and iteration procedures in local theory of normed spaces*, Missouri Conf. 1984.
19. V. D. Milman and G. Pisier, *Gaussian processes and mixed volumes*, Ann. Probab. (to appear).
20. A. Yu. Garnaev and E. D. Gluskin, *On diameters of the Euclidean sphere*, Dokl. Akad. Nauk SSSR **277** (1984), 1048–1052.

UNIVERSITÉ DES SCIENCES ET TECHNIQUES DE LILLE, U.E.R. DE MATHÉMATIQUES,
59655 VILLENEUVE D'ASCQ CEDEX, FRANCE

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA
T6J 2G1, CANADA