

U-EMBEDDED SUBSETS OF NORMED LINEAR SPACES

RONNIE LEVY AND M. D. RICE

ABSTRACT. A subset S of a metric space X is U -embedded in X if every uniformly continuous function $f: S \rightarrow R$ extends to a uniformly continuous function $F: X \rightarrow R$. Thus U -embedding is the uniform analogue of C -embedding. The Tietze extension theorem tells us exactly which subsets of metric spaces are C -embedded. The uniform analogue would tell us exactly which subsets of metric spaces are U -embedded. In this paper, a characterization of U -embedded subsets of the Euclidean plane (or any normed linear space) is given.

A subset S of a uniform space X is U -embedded in X if every real-valued uniformly continuous function $f: S \rightarrow R$ extends to a uniformly continuous function $F: X \rightarrow R$. Thus, U -embedding is the uniform analogue of C -embedding in topological spaces. One consequence of the Tietze extension theorem is that a subset of a metric space is C -embedded if and only if it is closed. Unlike the topological situation, a characterization of U -embedded subsets of metric spaces seems quite complicated. In this paper, we characterize those subsets of normed linear spaces which are U -embedded. As is usual in such situations, it is the convexity which will help us.

1. Preliminary definitions and results. Suppose that (X, d) is a metric space. If $a, b \in X$, and $\varepsilon > 0$, then we say that a and b are ε -linked by n links in X if there exists $a = x_0, x_1, \dots, x_n = b \in X$ such that $d(x_{k-1}, x_k) \leq \varepsilon$ for $k = 1, 2, \dots, n$. The finite sequence $a = x_0, \dots, x_n = b$ is called an ε -chain from a to b . If there exists an n such that a and b are ε -linked by n links in X , we say that a and b are ε -linked in X . A metric space is *uniformly connected* if it is not the union of two nonempty subsets which are a positive distance apart. Clearly, every connected metric space is uniformly connected. It is not difficult to see that X is uniformly connected if and only if for each $\varepsilon > 0$ and for each a and b in X , a and b are ε -linked in X .

Suppose (X, d) is a metric space and S is a subset such that every two elements of S are ε -linked in S , where ε is a positive number. Define

$$d_\varepsilon^S(a, b) = \inf \left\{ \sum_{i=1}^m d(z_{i-1}, z_i) : a = z_0, \dots, z_m = b \text{ is an } \varepsilon\text{-chain in } S \right\}$$

and let

$$m_\varepsilon^S(a, b) = \min\{m : \text{there exists an } \varepsilon\text{-chain in } S \text{ from } a \text{ to } b \text{ having } m \text{ links}\}.$$

Then $d_\varepsilon^S(a, b)$ measures the shortest distance one has to travel between a and b given that each step taken is in S and each step is at most ε units long. On the

Received by the editors March 15, 1985 and, in revised form, July 22, 1985.

1980 *Mathematics Subject Classification*. Primary 54C20.

Key words and phrases. U -embedding, Lipschitz for large distances.

other hand, $m_\varepsilon^S(a, b)$ gives the fewest number of steps of length at most ε one must take to get from a to b provided that each step is in S . Fairly simple examples show that it is possible to have the sum of the distances along every ε -chain from a to b having $m_\varepsilon^S(a, b)$ links be bounded away from $d_\varepsilon^S(a, b)$. Now let

$$\hat{r}_\varepsilon^S = \sup\{d_\varepsilon^S(a, b)/d(a, b) : a, b \in S, a \neq b\}$$

and let

$$r_\varepsilon^S = \sup\{m_\varepsilon^S(a, b)/m_\varepsilon^X(a, b) : a, b \in S, a \neq b\}.$$

When the subspace S is clear from the context, we omit the superscript S in \hat{r}_ε^S and r_ε^S .

1.1. PROPOSITION. *Suppose (X, d) is a metric space and S is a subspace of X . Then $r_\varepsilon \leq 4\hat{r}_\varepsilon + 2$. Therefore, if \hat{r}_ε is finite, then r_ε is finite.*

PROOF. Suppose a and b are in S . Given $\delta > 0$, choose an ε -chain $a = z_0, z_1, \dots, z_N = b$ from a to b such that

$$A = \sum_{i=1}^N d(z_{i-1}, z_i) \leq d_\varepsilon^S(a, b) + \delta$$

where N is the smallest integer for which there exists such an ε -chain. Then

$$(*) \quad d(z_{2k-2}, z_{2k}) > \varepsilon \quad \text{for } k = 1, \dots, [N/2],$$

because if this inequality did not hold, we could get an ε -chain having fewer links and the corresponding ε -chain would give a sum of distances not exceeding A . From $(*)$, we get $A \geq [N/2]\varepsilon/2$, so

$$[N/2]\varepsilon/2 \leq d_\varepsilon^S(a, b) + \delta.$$

Set $m = m_\varepsilon^S(a, b)$. By the choice of N and the definition of $m_\varepsilon^S(a, b)$, $m \leq N$. Repeated applications of the triangle inequality give $d(a, b) \leq \varepsilon m_\varepsilon^X(a, b)$. Then we get

$$\begin{aligned} m_\varepsilon^S(a, b)/m_\varepsilon^X(a, b) &= \varepsilon m_\varepsilon^S(a, b)/\varepsilon m_\varepsilon^X(a, b) \\ &\leq \varepsilon m_\varepsilon^S(a, b)/d(a, b) = \varepsilon m/d(a, b) \leq N\varepsilon/d(a, b), \end{aligned}$$

so

$$\begin{aligned} m_\varepsilon^S(a, b)/m_\varepsilon^X(a, b) &\leq N\varepsilon/d(a, b) \leq [4(d_\varepsilon^S(a, b) + \delta) + 2\varepsilon]/d(a, b) \\ &= 4d_\varepsilon^S(a, b)/d(a, b) + (4\delta + 2\varepsilon)/d(a, b). \end{aligned}$$

If $d(a, b) \geq \varepsilon$, then $m_\varepsilon^S(a, b)/m_\varepsilon^X(a, b) \leq 4\hat{r}_\varepsilon + (4\delta + 2\varepsilon)/\varepsilon = 4\hat{r}_\varepsilon + 2 + (4\delta/\varepsilon)$. On the other hand, if $d(a, b) < \varepsilon$, then $m_\varepsilon^S(a, b) = m_\varepsilon^X(a, b) = 1$ and $d_\varepsilon^S(a, b) = d(a, b)$, so $m_\varepsilon^S(a, b)/m_\varepsilon^X(a, b) = 1 = d_\varepsilon^S(a, b)/d(a, b)$. Therefore, $r_\varepsilon \leq 4\hat{r}_\varepsilon + 2 + 4\delta/\varepsilon$ for each $\delta > 0$, so $r_\varepsilon \leq 4\hat{r}_\varepsilon + 2$.

If (X, d) is a metric space and $f: X \rightarrow R$ is a function, then f is *Lipschitz for large distances* if for each $\varepsilon > 0$, there exists a constant K (which will in general depend upon ε) such that $d(x, y) \geq \varepsilon$ implies that $|f(x) - f(y)| \leq Kd(x, y)$. If F is a family of functions from X to R , then F is said to be *jointly Lipschitz for large distances* if for each $\varepsilon > 0$ there exists a constant K (depending upon ε) such that if $d(x, y) \geq \varepsilon$ and $f \in F$, then $|f(x) - f(y)| \leq Kd(x, y)$. The phrase “ (X, d) is a normed linear space” will be used to mean that d is the metric induced by the norm on the normed linear space X .

1.2. LEMMA [LR₁]. *If (X, d) is a normed linear space and S is a subset of X , then S is U -embedded in X if and only if each uniformly continuous function $f: S \rightarrow R$ is Lipschitz for large distances.*

1.3. LEMMA [LR₂]. *If (X, d) is a normed linear space and S is a uniformly connected subset of X , then S is U -embedded in X if and only if each equi-uniformly-continuous family F of functions from S to R is jointly Lipschitz for large distances.*

2. The uniformly connected case. In this section, we give a characterization of those uniformly connected subsets of normed linear spaces which are U -embedded. In the next section we show how to modify the characterization for the case where the subset S is not assumed to be uniformly connected.

2.1. PROPOSITION. *Suppose (X, d) is a normed linear space, and suppose that S is a uniformly connected U -embedded subset of X . Then \hat{r}_ϵ is finite for each $\epsilon > 0$.*

PROOF. Assume $\hat{r}_\epsilon = +\infty$ for some $\epsilon > 0$. Then there exist sequences (x_k) and (y_k) of points of S such that $x_k \neq y_k$ and $d_\epsilon^S(x_k, y_k) \geq kd(x_k, y_k)$ for $k = 1, 2, \dots$. Since $d(x, y) \leq \epsilon$ implies that $d_\epsilon^S(x, y) = d(x, y)$, the choice of the x_k 's and y_k 's gives us that $d(x_k, y_k) > \epsilon$ for $k \geq 2$. For $k = 1, 2, \dots$, define $g_k: S \rightarrow R$ by $g_k(x) = d_\epsilon^S(x, y_k)$. We claim that the family $\{g_k: k = 1, 2, \dots\}$ is equi-uniformly continuous. Choose $\eta > 0$ and let $\delta = \min\{\eta, \epsilon\}$. Suppose $x, y \in S$ and $d(x, y) < \delta$. Then $d(x, y) \leq \epsilon$. Given $\rho > 0$, choose ϵ -chains $y = y_0, y_1, \dots, y_{L(k)} = y_k$ and $x = x_0, x_1, \dots, x_{M(k)} = y_k$ such that

$$\sum_{i=1}^{L(k)} d(y_{i-1}, y_i) < d_\epsilon^S(y, y_k) + \rho$$

and

$$\sum_{j=1}^{M(k)} d(x_{j-1}, x_j) < d_\epsilon^S(x, y_k) + \rho.$$

Then

$$d_\epsilon^S(x, y_k) \leq d(x, y) + \sum_{i=1}^{L(k)} d(y_{i-1}, y_i) < d(x, y) + d_\epsilon^S(y, y_k) + \rho.$$

Therefore,

$$d_\epsilon^S(x, y_k) - d_\epsilon^S(y, y_k) < d(x, y) < \eta.$$

Similarly, one shows that

$$d_\epsilon^S(y, y_k) - d_\epsilon^S(x, y_k) < d(x, y) < \eta.$$

Therefore, $|g_k(x) - g_k(y)| \leq \eta$. This proves the claim. However, $|g_k(x_k) - g_k(y_k)| = d_\epsilon^S(x_k, y_k) \geq kd(x_k, y_k)$, $k = 1, 2, \dots$. Since $d(x_k, y_k) > \epsilon$ for $k \geq 2$, this violates 1.3.

2.2. PROPOSITION. *Suppose that S is a uniformly connected subset of a normed linear space (X, d) . If r_ϵ is finite for each positive ϵ , then S is U -embedded in X .*

PROOF. Suppose $f: S \rightarrow R$ is uniformly continuous. Suppose $\epsilon > 0$. We must find a constant K_ϵ such that $d(x, y) \geq \epsilon$ implies that $|f(x) - f(y)| < K_\epsilon d(x, y)$.

Choose $\delta > 0$ such that $d(x, y) < \delta$ implies that $|f(x) - f(y)| < 1$. Assume $d(x, y) \geq \varepsilon$. Let $m = m_\delta^S(x, y)$ and let $x = x_0, x_1, \dots, x_m = y$ be a δ -chain in S . Then if we let $K_\varepsilon = r_\delta[(\varepsilon + \delta)/\varepsilon\delta]$, we get

$$\begin{aligned} |f(x) - f(y)| &\leq \sum_{i=1}^m |f(x_{i-1}) - f(x_i)| \\ &\leq m = m_\delta^X(x, y)(m/m_\delta^X(x, y)) \\ &\leq r_\delta[(d(x, y)/\delta) + 1] \quad (\text{because } m_\delta^X(x, y) = [(d(x, y)/\delta) + 1]) \\ &\leq r_\delta\{d(x, y) + (\delta/\varepsilon)\varepsilon\}/\delta \\ &\leq r_\delta\{(\varepsilon d(x, y) + \delta d(x, y))/\varepsilon\delta\} \quad (\text{since } \varepsilon \leq d(x, y)) \\ &= K_\varepsilon d(x, y). \end{aligned}$$

Therefore, f is Lipschitz for large distances.

Combining 1.1 with the preceding two propositions gives the following theorem:

2.3. THEOREM. *Suppose that S is a uniformly connected subset of the normed linear space X . Then the following are equivalent:*

- (i) \hat{r}_ε is finite for each $\varepsilon > 0$.
- (ii) r_ε is finite for each $\varepsilon > 0$.
- (iii) S is U -embedded in X .

3. The general case. In this section, we give a characterization of those subsets of normed linear spaces which are U -embedded. The characterization will use the characterization given in §2 for the case where the subspace is uniformly connected.

3.1. PROPOSITION. *Suppose S is a U -embedded subset of the normed linear space (X, d) . Then for each $\varepsilon > 0$ there exists a compact subset F of X , which may be taken to be the union of finitely many line segments, such that*

- (i) every two elements of $S \cup F$ are ε -linked in $S \cup F$,
- (ii) $\hat{r}_\varepsilon^{S \cup F}$, and therefore $r_\varepsilon^{S \cup F}$, is finite.

PROOF. For each $p \in S$, define $C_p = \{x \in S : x \text{ and } p \text{ are } \varepsilon/3\text{-linked in } S\}$. If $C_p \neq C_q$, then $d(C_p, C_q) \geq \varepsilon/3$, so $\mathcal{C} = \{C_p : p \in S\}$ is a uniformly discrete family of nonempty subsets of X whose union is the U -embedded set S . Therefore, \mathcal{C} is finite. (See [LR₁].) Write $\mathcal{C} = \{D_0, D_1, \dots, D_N\}$ and for each $k = 0, 1, \dots, N$ choose $p_k \in D_k$. For $k = 1, 2, \dots, N$, let F_k be the line segment from p_{k-1} to p_k and let $F = \bigcup_{k=1}^N F_k$. With this definition, condition (i) is clearly satisfied. Now let $\hat{S} = \{p \in X : d(p, S \cup F) \leq \varepsilon/3\}$. It is easy to see that \hat{S} is uniformly connected. We claim that every uniformly continuous function $g : \hat{S} \rightarrow R$ which is identically zero on S is bounded. It will then follow from [LR₁] that \hat{S} is U -embedded in X . So assume $g : \hat{S} \rightarrow R$ is uniformly continuous and identically zero on S . Since F is compact, it follows that the restriction of g to $S \cup F$ is bounded. Let K be an upper bound for the absolute value of this restriction. Now let $\delta > 0$ be such that $x, y \in S \cup F$, $d(x, y) < \delta$ imply $|g(x) - g(y)| \leq 1$. Given a point x of \hat{S} , there exists a point p of $S \cup F$ such that the segment from x to p is contained in \hat{S} and has length at most $\varepsilon/3$. Then $|g(x)| \leq B + K$, where $B = [(\varepsilon/3\delta) + 1]$. Therefore, \hat{S} is U -embedded in X .

It follows from 2.3 that $\hat{r}_{\varepsilon/3}^S$ is finite. We will show that this implies that $\hat{r}_{\varepsilon}^{S \cup F}$ is finite. Suppose $a, b \in S \cup F$. Given $\rho > 0$, choose an $\varepsilon/3$ -chain $a = a_0, a_1, \dots, a_M = b$ in \hat{S} such that

$$(\#) \quad A = \sum_{i=1}^M d(a_{i-1}, a_i) < d_{\varepsilon/3}^S(a, b) + \rho.$$

We may assume without loss of generality that this $\varepsilon/3$ chain is minimal in the sense that $d(a_0, a_2) > \varepsilon/3, d(a_2, a_4) > \varepsilon/3, \dots$ (If this is not the case, inductively choose the elements of the chain so that the resulting chain is minimal and (#) will still hold.) Then at least half of the distances $d(a_{i-1}, a_i)$ are at least $\varepsilon/6$, so

$$(*) \quad A \geq (\varepsilon/6)[M/2], \quad \text{that is,} \quad M \leq (12A/\varepsilon) + 2.$$

For each $k = 1, 2, \dots, M - 1$, choose $x_k \in S \cup F$ such that $d(x_k, a_k) < \varepsilon/3$. Then if $x_0 = a$ and $x_M = b$, one gets that $d(x_{k-1}, x_k) \leq d(x_{k-1}, a_{k-1}) + d(a_{k-1}, a_k) + d(a_k, x_k) \leq d(a_{k-1}, a_k) + 2\varepsilon/3 \leq \varepsilon/3 + 2\varepsilon/3 = \varepsilon$, so $a = x_0, x_1, \dots, x_M = b$ is an ε -chain in $S \cup F$. Furthermore,

$$\sum_{i=1}^M d(x_{i-1}, x_i) \leq \sum_{i=1}^M d(a_{i-1}, a_i) + 2M\varepsilon/3.$$

Therefore, using (*) we get $\sum_{i=1}^M d(x_{i-1}, x_i) \leq A + (2\varepsilon/3)[(12A/\varepsilon) + 2] = 9A + (4\varepsilon/3) \leq 9d_{\varepsilon/3}^S(a, b) + [9\rho + (4\varepsilon/3)]$. Therefore, $d_{\varepsilon}^{S \cup F}(a, b) \leq 9d_{\varepsilon/3}^S(a, b) + (4\varepsilon/3)$. Hence, $\hat{r}_{\varepsilon}^{S \cup F} \leq 9\hat{r}_{\varepsilon/3}^S + (4\varepsilon/3)$.

3.2. PROPOSITION. *Suppose S is a subset of a normed linear space X . Suppose that for each $\varepsilon > 0$ there exists a compact set F such that*


- (i) *every two elements of $S \cup F$ are ε -linked in $S \cup F$,*
- (ii) *$r_{\varepsilon}^{S \cup F}$ is finite.*

Then S is U -embedded in X .

PROOF. Suppose $f: S \rightarrow R$ is uniformly continuous. By a theorem of Isbell [I], there exists an $\varepsilon > 0$ such that f can be extended to a uniformly continuous function $f_1: S_{\varepsilon} \rightarrow R$, where $S_{\varepsilon} = \{x \in X: d(x, S) \leq \varepsilon\}$. Choose a positive $\delta < \varepsilon$ such that $x, y \in S_{\varepsilon}$ and $d(x, y) < \delta$ imply that $|f_1(x) - f_1(y)| < 1$. Let F be the compact set given by the hypothesis corresponding to δ . By [LR₁], f_1 can be extended to a uniformly continuous function $\hat{f}: S_{\varepsilon} \cup F \rightarrow R$. We claim that there exists a constant C such that if x and y are elements of $S \cup F$ satisfying $d(x, y) < \delta$, then $|\hat{f}(x) - \hat{f}(y)| \leq C$. Let M be a constant such that $|f(x)| < M$ for all x in F . Suppose $x, y \in S \cup F$ and $d(x, y) < \delta$. If $\{x, y\} \subseteq F$, then $|f(x) - f(y)| \leq 2M$. If $\{x, y\} \subseteq S$, then $|f(x) - f(y)| < 1$. If $x \in S, y \in F$, then $y \in S_{\varepsilon}$ (because $d(x, y) < \varepsilon$) so again we have $|f(x) - f(y)| < 1$. Therefore, we may choose $C = 2M + 1$.

Now assume $d(x, y) \geq \varepsilon$, where x and y are points of $S \cup F$. Let $m = m_{\delta}^{S \cup F}(x, y)$ and let $x = x_0, x_1, \dots, x_m = y$ be a δ -chain in $S \cup F$. Then

$$\begin{aligned} |f(x) - f(y)| &\leq \sum_{i=1}^m |f(x_{i-1}) - f(x_i)| \leq mC \\ &\leq Cr_{\delta}^{S \cup F}[(\varepsilon + \delta)/\varepsilon\delta]d(x, y) = K_{\varepsilon}d(x, y), \end{aligned}$$



Diagram

where $K_\varepsilon = Cr_\delta^{S \cup F}(\varepsilon + \delta)/\varepsilon\delta$. Therefore, f is Lipschitz for large distances.

Combining 3.1 and 3.2 gives the following theorem:

3.3. THEOREM. *Suppose S is a subset of the normed linear space (X, d) . Then the following are equivalent:*

- (i) S is U -embedded in X ,
- (ii) for each $\varepsilon > 0$, there exists a compact set F (which may be taken to be a union of finitely many line segments) such that any two elements of $S \cup F$ are ε -linked in $S \cup F$ and such that $r_\varepsilon^{S \cup F}$ is finite.

REMARKS. 1. If (X, d) is any uniformly connected metric space, then by embedding X isometrically in a normed linear space and appealing to 1.2 and 2.3 one sees that if every uniformly continuous function $f: X \rightarrow R$ is Lipschitz for large distances, then r_ε^X is finite for each positive ε . In fact, the converse of this statement is true as well: If r_ε^X is finite for each positive ε , then every uniformly continuous function $f: X \rightarrow R$ is Lipschitz for large distances. Since there are easy examples of metric spaces where not every uniformly continuous real-valued function is Lipschitz for large distances, this means that the finiteness of r_ε^S for each positive ε is not in general equivalent to the U -embedding of a subspace S of a space X .

2. We do not know an example of a metric space (X, d) and a non- U -embedded uniformly connected subset S such that r_ε^S is finite for each positive ε . However, the following example shows that a uniformly connected, U -embedded subset of a metric space can have infinite r_ε^S for all sufficiently small $\varepsilon > 0$.

3.4. EXAMPLE. There exists a (uniformly) connected metric space $X \subset R^2$ and a U -embedded (uniformly) connected subset S such that $r_\varepsilon = +\infty$ for all positive $\varepsilon < 1/2$. Rather than give a description of X and S , we will draw pictures. Any uniformly continuous $f: S \rightarrow R$ can be extended to a uniformly continuous function $F: X \rightarrow R$ by making F constant on the small vertical whiskers growing along the y -axis. By restricting our attention to those horizontal bands of S of height $1/n$, one can show that $r_{1/(2n)} = \infty$ for $n = 1, 2, \dots$. For an arbitrary positive $\varepsilon \leq 1/2$, find an n satisfying $1/(2n) < \varepsilon < 1/n$ (that is, $1/2 < n\varepsilon \leq 1$), and by restricting our attention to the horizontal bands of height $1/n$, we can again show that $r_\varepsilon = \infty$.

REMARKS. 1. By modifying the construction in 3.4, it is possible to find a connected metric space X and a connected U -embedded subset S such that $r_\varepsilon = \infty$ for all $\varepsilon > 0$.

2. By appealing to the results in [LR₃] the results in this paper also give results about Hilbert spaces. For example, it is shown in [LR₃] that a subset S of a Hilbert space H is U -embedded in H if and only if every uniformly continuous function $f: S \rightarrow H$ extends to a uniformly continuous function $F: H \rightarrow H$. Thus, we also have characterized those subsets of Hilbert space for which every uniformly continuous function into a Hilbert space extends to a uniformly continuous function.

REFERENCES

- [I] John Isbell, *Uniform spaces*, Math. Surveys, no. 12, Amer. Math. Soc., Providence, R. I., 1964.
- [LR₁] Ronnie Levy and M. D. Rice, *Techniques and examples in U-embedding*, Topology Appl. (to appear).
- [LR₂] —, *The extension of equi-uniformly continuous families of mappings*, Pacific J. Math. **12** (1985), 149–161.
- [LR₃] —, *The approximation and extension of uniformly continuous Banach space valued mappings*, Comment. Math. Univ. Carolin. **24** (1983), 251–265.

DEPARTMENT OF MATHEMATICAL SCIENCES, GEORGE MASON UNIVERSITY, FAIRFAX, VIRGINIA 22030

DEPARTMENT OF COMPUTER SCIENCE, GEORGE MASON UNIVERSITY, FAIRFAX, VIRGINIA 22030