

SPACES EVERY QUOTIENT OF WHICH IS METRIZABLE

A. J. JAYANTHAN AND V. KANNAN

(Communicated by Dennis Burke)

ABSTRACT. We characterise those topological spaces for which every quotient image is metrizable. This supplements the earlier known results in this direction, in a fairly complete manner.

0. Introduction. Metrizable of quotients of metric spaces has been studied by many mathematicians [1, 2, 6, 7]. In an elementary course in general topology we learn that every continuous image in Hausdorff space of a compact metric space is metrizable [8]. In [7], Willard proved that every closed continuous image of a metric space X in Hausdorff space is metrizable if and only if the set of all limit points of X is compact. We consider the following problem. What are all topological spaces every Hausdorff quotient space of which is metrizable? To our surprise, the condition on the space turns out to be the same as that of Willard. We prove that every quotient of a metric space X in Hausdorff spaces is metrizable if and only if the set of all limit points of X is compact. In the process we give plenty of equivalent conditions for this.

1. The main theorem. We first fix some notation and terminology. Let X be a topological space. The set of all limit points of X is denoted by X^1 . By the term map we always mean a continuous map. Let Y be any other topological space and $f: X \rightarrow Y$ be any map onto Y . We say that f is compact-to-one (respectively finite-to-one, n -to-one) if $f^{-1}(y)$ is a compact (respectively a finite, having at most n points) subset of X for all points y in Y , and in these cases Y is called a compact-to-one (respectively finite-to-one, n -to-one) image of X . For $f: X \rightarrow Y$ and $A \subset X$, we say that A is saturated with respect to f , if $f^{-1}(f(a)) \subset A$ for all a in A . All spaces considered in this article are assumed to be Hausdorff spaces and all maps are assumed to be continuous from Hausdorff spaces to Hausdorff spaces. A quotient space which is also a Hausdorff space is called a Hausdorff quotient space or simply a Hausdorff quotient. ω denotes the first infinite ordinal number and \mathbf{N} denotes the set of all natural numbers.

EXAMPLE. We now see a simple example of a metric space which has a non-metrizable quotient. Let $A = \{0, 1, \frac{1}{2}, \frac{1}{3}, \dots\}$. Define $X = \mathbf{N} \cup \{0\} \times A \subset \mathbf{R}^2$ and $Y = \{(x, y) \in A \times A : x = 0 \text{ implies } y = 0\}$. Define a map $q: X \rightarrow Y$ as follows:

$$q(x) = \begin{cases} (1/n, 0) & \text{if } x \text{ is } (0, 1/n) \text{ or } (n, 0), \\ (1/m, 1/n) & \text{if } x \text{ is } (m, 1/n), m \geq 1, \\ (0, 0) & \text{if } x \text{ is } (0, 0). \end{cases}$$

Received by the editors February 12, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 54B15, 54E35.

Key words and phrases. Quotient spaces, metrizable.

The first author was supported by the Council of Scientific and Industrial Research, India.

Give the usual topology for X and the quotient topology induced by q for Y . One can verify that Y is not even first countable. Y with this topology is denoted by S_2 throughout this article (see also [3]). We exploit properties of S_2 directly and indirectly at many places below.

A metric d on a metric space X is called normal if $d(A, B) = \inf\{d(a, b) : a \in A, b \in B\} > 0$ for every pair A, B of disjoint closed subsets of X . The following theorem will be useful in the proof of our main theorem.

THEOREM 1.1 [4]. *Suppose X is a metric space. Then there exists an equivalent normal metric on X if and only if X^1 is compact.*

Now we prove the first set of equivalent forms to the metrizability of every Hausdorff quotient of a metric space. Hereafter we write simply "quotient" for Hausdorff quotient space.

THEOREM 1.2. *Let X be a metric space. Then the following conditions are equivalent:*

- (1) *Every quotient of X is metrizable.*
- (2) *X is the union of a compact subspace and a discrete subspace.*
- (3) *X^1 is compact.*
- (4) *ω^2 cannot be embedded as a closed subspace in X .*

PROOF. (1) \Rightarrow (2). Suppose X cannot be written as a union of a compact subspace and a discrete subspace. We know that $X \setminus X^1$ is a discrete subspace and hence X^1 is not compact. Choose a sequence $\{x_n\} \subset X^1$ such that $x_m \neq x_n$ for every $m \neq n$ and $\{x_n : n \in \mathbb{N}\}$ does not have a limit point in X . So $\{x_n : n \in \mathbb{N}\}$ is a closed subspace of X . Since x_1 is in X^1 , there exists a sequence $\{y_n\} \subset X$ such that $y_m \neq y_n$ for every $m \neq n$ and $y_n \rightarrow x_1$. Define a new space Y by identifying x_n with y_n for every $n > 1$ and giving the corresponding quotient topology to Y . Then one can verify that Y contains a copy of S_2 (see the proof of Theorem 2.2) which is not metrizable. So Y is not metrizable. This proves that (1) \Rightarrow (2).

(2) \Rightarrow (3). Suppose $X = A \cup B$ where A is a discrete subspace and B is a compact subspace. We may assume that $A \cap B = \emptyset$.

Claim. B contains X^1 .

On the contrary, if $A \cap X^1 \neq \emptyset$, say, x is in $A \cap X^1$, there exists an open subset $U \subset X$ such that $U \cap A = \{x\}$. Clearly, $\overline{U} \setminus \{x\}$ is a closed subset of B which fails to be compact since x is a limit point of \overline{U} . This contradiction proves our claim.

So X^1 is compact, being a closed subspace of B .

(3) \Rightarrow (1). Suppose X^1 is compact and $f : X \rightarrow Y$ is a quotient map onto Y . Denote the set $f(X^1)$ by Y_1 . We use, for convenience, the same symbol Y_1 to denote the same space as subspace of different spaces. This does not create confusion because the subspace topology is the same in all these cases. D_1 denotes the set $\{x \in X \setminus X^1 : f(x) \in Y_1\}$ and D_2 denotes $(X \setminus X^1) \setminus D_1$. Note that $X = X^1 \cup D_1 \cup D_2$ where \cup denotes the disjoint set union. We denote the set $Y_1 \cup D_1 \cup D_2$ by X_1 and define a function $g_1 : X \rightarrow X_1$ as follows:

$$g_1(x) = \begin{cases} f(x) & \text{if } x \text{ is in } X^1, \\ x & \text{otherwise.} \end{cases}$$

Provide X_1 with the quotient topology induced by g_1 . Then we claim that g_1 is a closed map. For, let $K \subset X$ be a closed subset. Then, $K = K_1 \cup K_2$ where

$K_1 \subset X^1$ and $K_2 \cap X^1 = \emptyset$. $g_1^{-1}(g_1(K)) = g_1^{-1}(g_1(K_1)) \cup K_2 = g_1^{-1}(g_1(K_1)) \cup K$ is closed in X because $g_1(K_1)$ is closed in X_1 . Therefore, g_1 is a closed map. Now by Willard's theorem [7], X_1 is metrizable. Since $X_1^1 \subset Y^1$, X_1^1 is compact and hence by Theorem 1.1, there exists a normal metric d on X_1 . Now, let X_2 be the subspace $Y_1 \cup D_2$ of X_1 . Define $g_2: X_1 \rightarrow X_2$ as follows:

$$g_2(x) = \begin{cases} f(x) & \text{if } x \text{ is in } D_1, \\ x & \text{otherwise.} \end{cases}$$

It is not difficult to see that the quotient topology induced by g_2 and the subspace topology on X_2 coincide. For every point y in $f(D_2)$, $f^{-1}(y) \subset D_2$. We use the same notation $f^{-1}(y)$ as a subset of X_2 also. Since d is a normal metric, for every y in $f(D_2)$ there exists x_y in $f^{-1}(y)$ such that $d(x_y, Y_1) < 2d(f^{-1}(y), Y_1)$. Let $D_3 = \{x_y: y \in f(D_2)\}$ and X_3 be the subspace $Y_1 \cup D_3$ of X_2 . We define $g_3: X_2 \rightarrow Y$ as follows:

$$g_3(x) = \begin{cases} f(x) & \text{if } x \text{ is in } D_2, \\ x & \text{otherwise.} \end{cases}$$

For any point x in X_3 , $[x]$ denotes the subset $g_3^{-1}(g_3(x))$ and for any subset $A \subset X_3$, $[A] = g_3^{-1}(g_3(A))$. We claim that a subset $F \subset X_3$ is closed in X_3 if and only if $[F]$ is closed in X_2 . Since $[F] \cap X_3 = F$, F is closed in X_3 whenever $[F]$ is closed in X_2 . Now, suppose F is closed in X_3 and x is in $\overline{[F]}$. To show that x is in $[F]$. Note that x is in $[F]$ if x is an isolated point in $\overline{[F]}$. So we assume that x is not an isolated point in $\overline{[F]}$. Observe that x is in Y_1 . If every neighborhood of x meets $[F] \cap Y_1 = F \cap Y_1$ at infinitely many points, then x is in $\overline{F \cap Y_1} = F \cap Y_1$ (because Y_1 is compact and F is closed) $\subset [F]$. Suppose x has a neighborhood U_x such that $U_x \cap Y_1 = \{x\}$. For a given $\varepsilon > 0$, let $B_\varepsilon(x)$ denote the ε -ball around x in X_2 . Choose x' in $B_\varepsilon(x) \cap U_x$ (which is infinite) such that $x \neq x'$ and $d(x', x) < \varepsilon/2$. Then by the choice of D_3 , $d(x_{f(x')}, x) < \varepsilon$ and therefore $x_{f(x')}$ is in $B_\varepsilon(x)$. Hence $x_{f(x')}$ is in $B_\varepsilon(x) \cap F$, showing that $B_\varepsilon(x) \cap F$ is nonempty for all ε . So x is in $\overline{F} = F \subset [F]$. This completes the proof that $[F]$ is also closed whenever F is closed.

Now, let h denote the restriction of g_3 to X_3 . We claim that $h: X_3 \rightarrow Y$ is a homeomorphism. h is one-one and onto. Let F be closed in Y . Then $f^{-1}(F)$ is closed in X and saturated with respect to $g_2 \circ g_1$. So $g_2(g_1(f^{-1}(F)))$ is closed in X_2 and hence $h^{-1}(F) = g_2(g_1(f^{-1}(F))) \cap X_3$ is closed in X_3 . Thus h is continuous. If $[F]$ is closed in X_2 , then $g_1^{-1}(g_2^{-1}([F]))$ is closed in X . Since $g_1^{-1}(g_2^{-1}([F]))$ is saturated with respect to f , $f(g_1^{-1}(g_2^{-1}([F]))) = h(F)$ is closed in Y . Hence h is a homeomorphism. This proves that Y is metrizable because X_3 is.

(3) \Rightarrow (4). Suppose X^1 is compact. If ω^2 is embedded in X as a closed subspace, then $\{\omega.n: n \in \mathbb{N}\}$ is an infinite discrete closed subspace of X^1 . This contradicts the hypothesis that X^1 is compact and thus (3) \Rightarrow (4) is proved.

(4) \Rightarrow (3) can be proved by a reverse argument of the proof of (3) \Rightarrow (4).

2. Some more equivalent forms. In this section, we give many more equivalent forms of the statements of Theorem 1.2. A topological space Y is called a Fréchet space if for any subset $A \subset Y$, y is in \overline{A} if and only if there exists a sequence in A converging to y . We exploit properties of S_2 to obtain many more equivalent forms of the property that "every quotient of the space is metrizable".

THEOREM 2.1. *For a metric space X , the following conditions are equivalent:*

- (a) *Every quotient of X is metrizable.*
- (b) *Every quotient of X is first countable.*
- (c) *Every quotient of X is a Fréchet space.*

Statement (a) is equivalent to any of the following:

- (d) *every compact-to-one quotient of X is P ,*
- (e) *every finite-to-one quotient of X is P ,*
- (f) *every n -to-one quotient of X is P for all n in \mathbb{N} ,*

where P stands for the property of being either 'metrizable' or 'first countable' or 'a Fréchet space'.

Every statement obtained by replacing 'quotient' by 'open continuous image' in statements (b) to (f) is again equivalent to (a).

PROOF. The proof follows from the following facts:

- (i) S_2 does not satisfy P (where P is as in the statement) [3].
- (ii) We can find an open, two-to-one, continuous image of X containing S_2 whenever X^1 is not compact (see the space Y in the proof of (3) \Rightarrow (2) of Theorem 2.2).
- (iii) Metrizable \Rightarrow first countable \Rightarrow a Fréchet space [3].

A quotient map $q: X \rightarrow Y$ onto Y is called hereditarily quotient if the restriction $q|_A$ of q to $A \subset X$ is again a quotient map for every saturated subspace A of X . q is called pseudo-open if y is in $\text{Int } q(U)$ for every open set U containing $q^{-1}(y)$.

Arkhangel'skii [1] has proved that a quotient map is hereditarily quotient if and only if it is pseudo-open. See also [4].

THEOREM 2.2. *On a metric space X the following conditions are equivalent:*

- (1) *Every quotient of X is metrizable.*
- (2) *Every surjective quotient map from X is pseudo-open.*
- (3) *Every surjective quotient map from X is hereditarily quotient.*

PROOF. (1) \Rightarrow (2). By Theorem 1.2, (1) implies that X^1 is compact. Let $f: X \rightarrow Y$ be a surjective quotient map and $D = X \setminus f^{-1}(f(X^1))$. Denote by X_1 the set $D \cup f(X^1)$ and define $g: X \rightarrow X_1$ as follows:

$$g(x) = \begin{cases} x & \text{if } x \text{ is in } D, \\ f(x) & \text{otherwise.} \end{cases}$$

Providing X_1 with the quotient topology induced by g , one can prove that g is a closed map by a similar argument as in the proof of Theorem 1.2. Define $h: X_1 \rightarrow Y$ as follows:

$$h(x) = \begin{cases} f(x) & \text{if } x \text{ is in } D, \\ x & \text{otherwise.} \end{cases}$$

Then h is an open map onto Y and $f = h \circ g$. Since both open maps and closed maps are pseudo-open, we can prove that f is pseudo-open.

(2) \Rightarrow (3) is known [1].

(3) \Rightarrow (1). If X does not satisfy (1), X^1 is not compact by Theorem 1.2. So we can choose an infinite discrete closed subset $\{x_m: m = 0, 1, 2, \dots\}$ of X^1 . Let $\{U_m: m = 0, 1, 2, \dots\}$ be a sequence of pairwise disjoint open subsets of X such

that x_m is in U_m for all m . For each $m = 0, 1, 2, \dots$, we can choose a sequence $\{x_{m,n} : n \in \mathbf{N}\} \subset U_m$ such that $x_{m,n}$ converges to x_m . Let

$$Y = X \setminus \{x_n : n = 1, 2, \dots\} \quad \text{and} \quad f: X \rightarrow Y$$

be defined as follows:

$$f(x) = \begin{cases} x_{0,n} & \text{if } x = x_n, \\ x & \text{otherwise.} \end{cases}$$

Give quotient topology for Y (induced by f).

Claim. f is not hereditarily quotient.

For, consider $A = \{x_{m,n} : m, n = 1, 2, \dots\} \cup \{x_0\}$. A is saturated. If $f|_A$ were a quotient map onto its image $f(A)$, then $f(x_0)$ would have been isolated in $f(A)$. But $f(x_0)$ is not isolated in $f(A)$. Thus the claim is proved. This proves (3) \Rightarrow (1).

REMARKS. 1. The statements (2) and (3) of Theorem 2.2 have some more equivalent forms similar to those of Theorem 2.1. Each statement obtained by replacing 'quotient map' in (2) and (3) by 'compact-to-one quotient map', 'finite-to-one quotient map' etc. is equivalent to statement (1). Further the statements obtained by replacing 'quotient maps' by 'open maps' in all the above cases are also equivalent to (1).

2. In [4], the author has obtained a necessary and sufficient condition for every quotient of a space to be hereditarily quotient. The reader shall note that our conditions are entirely different from what is obtained in [4] because we consider quotients in only Hausdorff spaces in the present article.

3. Arkhangel'skii discusses three general problems in his survey article [1]. The first among them is the following: Under what circumstances can each space of a given class A be mapped onto a space of a given class B by means of a mapping belonging to a given class L ? When this is possible he writes $A L B$. Here we have found a maximal subclass A of the class of all metric spaces in connection with this problem for different classes L of maps such as open maps, quotient maps, finite-to-one quotient maps, hereditarily quotient maps etc., and for classes B such as metric spaces, first countable spaces and Fréchet spaces and showed that $A L B$ holds.

REFERENCES

1. A. V. Arkhangel'skii, *Mappings and spaces*, Russian Math. Surveys **21** (1966), 115–162.
2. —, *A condition for preservation of metrizability*, Soviet Math. Dokl. **6** (1965), 1127–1130.
3. A. V. Arkhangel'skii and S. P. Franklin, *Ordinal invariants for topological spaces*, Michigan Math. J. **15** (1968), 313–320.
4. V. Kannan, *Ordinal invariants in topology*, Mem. Amer. Math. Soc. No. 245 (1981).
5. S. G. Mrowka, *Normal metrics*, Amer. Math. Monthly **72** (1965), 998–1001.
6. A. H. Stone, *Metrizability of decomposition spaces*, Proc. Amer. Math. Soc. **7** (1956), 690–700.
7. S. Willard, *Metric spaces all of whose decompositions are metric*, Proc. Amer. Math. Soc. **21** (1969), 126–128.
8. —, *General topology*, Addison-Wesley, 1970.

SCHOOL OF MATHEMATICS & COMPUTER/INFORMATION SCIENCES, UNIVERSITY OF HYDERABAD, HYDERABAD 500 134, INDIA