

## A DYNAMICAL SYSTEM ON $R^3$ WITH UNIFORMLY BOUNDED TRAJECTORIES AND NO COMPACT TRAJECTORIES

K. M. KUPERBERG AND COKE S. REED

(Communicated by Kenneth R. Meyer)

**ABSTRACT.** This paper contains an example of a rest point free dynamical system on  $R^3$  with uniformly bounded trajectories, and with no circular trajectories. The construction is based on an example of a dynamical system described by P. A. Schweitzer, and on an example of a dynamical system on  $R^3$  constructed previously by the authors.

### INTRODUCTION

In order to answer S. Ulam's question posed in the Scottish Book (see Problem 110 in [5 or 3]) we constructed in [2] a rest point free dynamical system  $\Phi$  on  $R^3$  with all trajectories uniformly bounded by a given  $\varepsilon > 0$ . The function  $f(x) = \Phi(1, x)$  provided a counterexample to Ulam's problem. The dynamical system  $\Phi$  uses elements similar to Fuller's flow described in [1] (see also Wilson [6]), and hence  $\Phi$  contains circular trajectories. The purpose of this paper is to eliminate all circular trajectories by modifying the example described in [2]. This is done with the help of Schweitzer's flow, constructed in [4] in order to solve the Seifert conjecture on closed integral curves of a nonvanishing vector field on  $S^3$ .

### THE EXAMPLE

P. A. Schweitzer, see [4, pp. 393–394], based the construction of a “plug” with no circular trajectories on Denjoy's vector field on a surface of a torus. A  $C^1$  Denjoy vector field is included in the Appendix of [4]. We will now define a Schweitzer flow in such a way that the plug “stops” an open set of trajectories.

Assume the following notation:  $R^n =$  Euclidean  $n$ -space,  $S^1 =$  one-dimensional sphere,  $T = S^1 \times S^1$ ,  $d =$  distance function on  $T$ ,  $Z = C^1$  Denjoy's vector field on  $T$ ,  $A =$  exceptional minimal set of  $Z$ , and  $\Phi(t, x) =$  trajectory of a dynamical system  $\Phi$  passing through  $x$ .

Let  $B$  be a compact proper subset of  $T$ , invariant under the dynamical system generated by  $Z$ , and such that  $A \subset \text{Int } B$ . Let  $p_0 \in T - B$  be a point.

---

Received by the editors June 19, 1988 and, in revised form, October 26, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 58F25; Secondary 54H20.

©1989 American Mathematical Society  
0002-9939/89 \$1.00 + \$.25 per page

For  $i = 0, 1, 2$ , let  $D_i \subset T$  be a disc centered at  $p_0$  with radius  $r_i$  such that  $r_0 < r_1 < r_2 \leq d(p_0, B)$ . Let  $f: T \rightarrow R^1$  be a  $C^\infty$  mapping such that  $f|_{D_1} \equiv 0$ ,  $f|_{T - D_2} \equiv 1$ , and if  $q_1, q_2 \in D_2 - D_1$  are such that  $d(q_1, p_0) < d(q_2, p_0)$ , then  $f(q_1) < f(q_2)$ . Let  $g: R^1 \rightarrow R^1$  be an increasing  $C^\infty$  function such that  $g|_{(-\infty, 0]} \equiv 0$  and  $g|_{[1/6, \infty)} \equiv 1$ . Define  $\alpha, \beta: R^1 \rightarrow R^1$  by

$$\alpha(s) = \begin{cases} 0 & \text{if } s \leq \frac{1}{6}, \\ g(s - \frac{1}{6}) & \text{if } \frac{1}{6} \leq s \leq \frac{1}{3}, \\ g(\frac{1}{2} - s) & \text{if } \frac{1}{3} \leq s \leq \frac{1}{2}, \\ -\alpha(1 - s) & \text{if } s \geq \frac{1}{2}, \end{cases}$$

$$\beta(s) = \begin{cases} \alpha(s) & \text{if } s \leq \frac{1}{2}, \\ -\alpha(s) & \text{if } s \geq \frac{1}{2}. \end{cases}$$

Set  $M = T - D_0$ . Let  $V$  be a vector field defined on the manifold  $M \times R^1$  by setting

$$V((p, s)) = (f(p)Z(p)\alpha(s), 1 - f(p)\beta(s)).$$

Notice that if  $V((p, s)) = (v_1, v_2)$ , then  $V((p, 1 - s)) = (-v_1, v_2)$ ; and  $V((p, s)) = (0, 1)$  if either  $p \in D_1 - D_0$ ,  $s \leq 1/6$ , or  $s \geq 5/6$ . The dynamical system  $\Phi$  generated by  $V$  has the following properties:

- (i) for every point  $p \in B$  the trajectory containing the point  $(p, 1)$  has its  $\alpha$ -limit points in the interior of the set  $M \times [0, 1]$ , and the trajectory containing the point  $(p, 0)$  has its  $\omega$ -limit points in the interior of the set  $M \times [0, 1]$ ,
- (ii) for any number  $t > 0$  and any point  $(p, s) \in [M \times (R^1 - \{1/3, 2/3\})] \cup \text{Int}[(D_2 - D_0) \times R^1]$  if  $\Phi(t, (p, s)) = (q, r)$ , then  $r > s$ ,
- (iii) if  $(p, 0)$  and  $(q, 1)$  belong to the same trajectory, then  $p = q$ ,
- (iv)  $\Phi$  contains no circular trajectories.

There exists a  $C^\infty$  embedding  $\mu: M \times [0, 1] \rightarrow R^3$  such that for any  $p \in M$ , the set  $\{p\} \times [0, 1]$  is contained in a line parallel to the  $z$ -axis, and for any  $(p, s) \in M \times [0, 1]$ , there exists a neighborhood  $N$  of  $p$  in  $M$  such that the projection onto the  $xy$ -plane restricted to the set  $\mu(N \times \{s\})$  is an embedding, see [2]. We may also assume that  $\mu(M \times [0, 1])$  is a subset of the cube  $\{(x, y, z) \in R^3: 0 \leq x, y, z \leq 1\}$ .

There exists a  $C^1$  vector field  $U$  defined on  $R^3$  such that  $U$  and the dynamical system  $\psi$  generated by  $U$  satisfy the following conditions:

- (i)  $(x, y, z) \notin \mu(M \times [0, 1])$ , then  $U((x, y, z)) = (0, 0, 1)$ ,
- (ii) for any trajectory  $\psi(t, (x, y, z))$ , the set  $\psi(t, (x, y, z)) \cap \mu(M \times [0, 1])$  is contained in the set  $\mu(\Phi(t, (p, s)))$  for some  $(p, s) \in M \times [0, 1]$ ,
- (iii) for any trajectory  $\psi(t, (x, y, z))$ , the set  $\psi(t, (x, y, z)) \cap (R^3 - \mu(M \times [0, 1]))$  is contained in a line parallel to the  $z$ -axis.

*Remark.* No trajectory of  $\psi$  is compact, and if a trajectory of  $\psi$  passes through the points  $(x_1, y_1, 0)$  and  $(x_2, y_2, 1)$ , then  $(x_1, y_1) = (x_2, y_2)$ .

There exists a nonempty subset of the plane  $Q = \{(x, y) \in R^2: a \leq x \leq a + \delta, b \leq y \leq b + \delta\}$  such that if  $(x, y) \in Q$ , then the  $\alpha$ -limit points of  $\psi(t, (x, y, 1))$  and the  $\omega$ -limit points of  $\psi(t, (x, y, 0))$  are in the interior of the cube  $\{(x, y, z) \in R^3: 0 \leq x, y, z \leq 1\}$ .

Let  $[x]$  denote the greatest integer less than or equal to  $x$ . Let  $N_0$  be an integer greater than  $1/\delta$ . We define a vector field  $W$  on  $R^3$  in the following way: if  $iN_0 + j \leq z < iN_0 + j + 1$ , where  $i$  and  $j$  are integers, and  $0 \leq j < N_0$ , then put

$$W((x, y, z)) = U((x - i/N_0 - [x - i/N_0], y - j/N_0 - [y - j/N_0], z - [z])).$$

Clearly,  $W$  is a  $C^1$  vector field defined on  $R^3$ . Similarly as in [1], the trajectories of the dynamical system generated by  $W$  are uniformly bounded; every trajectory is contained in a box of dimensions 3, 3, and  $N_0^2$ , and hence it is bounded by  $\sqrt{18 + N_0^4}$ .

*Remark 1.* By rescaling the example, the uniform bound for the diameter of the trajectories can be changed to an arbitrarily pre-assigned  $\varepsilon > 0$ .

*Remark 2.* The vector field constructed in this paper is of class  $C^1$ . The following question remains unanswered: Does there exist a  $C^\infty$  dynamical system on  $R^3$  with uniformly bounded trajectories and no compact trajectories?

## REFERENCES

1. F. B. Fuller, *An index of fixed point type for periodic orbits*, Amer. J. Math. **89** (1967), 133–148.
2. K. Kuperberg and C. Reed, *A rest point free dynamical system on  $R^3$  with uniformly bounded trajectories*, Fund. Math. **114** (1981), 229–234.
3. R. D. Mauldin (ed.), *The Scottish book: Mathematics from the Scottish Café*, Birkhauser, Boston, 1981.
4. P. A. Schweitzer, *Counterexamples to the Seifert conjecture and opening closed leaves of foliations*, Ann. of Math. (2) **100** (1974), 386–400.
5. S. M. Ulam, *The Scottish book*, Los Alamos Scientific Monograph LA-6832.
6. F. W. Wilson, Jr., *On the minimal sets of non-singular vector fields*, Ann. of Math. (2) **84** (1966), 529–536.

DEPARTMENT OF MATHEMATICS, AUBURN UNIVERSITY, AUBURN, ALABAMA 36849

SUPERCOMPUTING RESEARCH CENTER, LANHAM, MARYLAND 20706