

ON WEAK CONVERGENCE IN $H^1(\mathbf{R}^d)$

PETER W. JONES AND JEAN-LIN JOURNÉ

(Communicated by J. Marshall Ash)

ABSTRACT. We prove that the a.e. convergence of a sequence of functions bounded in $H^1(\mathbf{R}^d)$ to a function in $L^1(\mathbf{R}^d)$ implies weak convergence.

Let $H^1(\mathbf{R}^d)$ denote the Hardy space of functions on \mathbf{R}^d whose Poisson maximal function lies in L^1 , and let $\text{BMO}(\mathbf{R}^d)$ denote the dual space [2] of functions of bounded mean oscillation. The space $\text{VMO}(\mathbf{R}^d)$, the closure of the Schwartz space \mathcal{S} in $\text{BMO}(\mathbf{R}^d)$, is the predual of $H^1(\mathbf{R}^d)$. We answer a question of Lions and Meyer by proving the following result. The theorem is also valid for Martingale H^1 , and the proof we give carries over directly to that setting.

Theorem. *Suppose $\{f_n\}$ is a sequence of $H^1(\mathbf{R}^d)$ functions such that $\|f_n\|_{H^1} \leq 1$ for all n and such that $f_n(x) \rightarrow f(x)$ for almost every $x \in \mathbf{R}^d$. Then $f \in H^1(\mathbf{R}^d)$, $\|f\|_{H^1} \leq 1$, and*

$$\int_{\mathbf{R}^d} f_n \varphi \, dx \rightarrow \int_{\mathbf{R}^d} f \varphi \, dx$$

for all $\varphi \in \text{VMO}(\mathbf{R}^d)$.

Proof. We may suppose that $\|\varphi\|_{L^1}, \|\varphi\|_{L^\infty}, \|\partial\varphi/\partial x_j\|_{L^\infty} \leq 1$ and that support φ is compact. Fix $\delta > 0$ and pick $\eta > 0$ so that $\eta \exp\{\delta^{-1}\} \leq \delta^{1+d}$ and, whenever $|E| \leq C\eta \exp\{\delta^{-1}\}$, $\int_E |f| \, dx \leq \delta$. This can be done because by Fatou's lemma $\|f\|_{L^1} \leq 1$. Now pick n large enough so that

$$|E_n| = |\{x \in \text{support } \varphi : |f_n(x) - f(x)| > \eta\}| \leq \eta.$$

Define

$$\tau(x) = \max\{0, 1 + \delta \log M\chi_{E_n}(x)\},$$

where $M(\cdot)$ denotes the Hardy-Littlewood maximal function. Then $0 \leq \tau \leq 1$, $\tau = 1$ a.e. dx on E_n , and by a result due to Coifman and Rochberg [1], $\|\tau\|_{\text{BMO}} \leq C\delta$. By the weak $(1, 1)$ estimate for the maximal function [3],

$$|\text{support } \tau| \leq C|E_n| \exp\{\delta^{-1}\} \leq C\eta \exp\{\delta^{-1}\}.$$

Received by the editors October 13, 1989 and, in revised form, April 15, 1992.

1991 *Mathematics Subject Classification.* Primary 42B30; Secondary 30D55.

Key words and phrases. H^1 , BMO, VMO.

The first author was supported by NSF Grant DMS-86-02500. The second author was supported by NSF Grant DMS-88-15651.

Consequently,

$$\int_{\text{support } \tau} |f| dx \leq \delta.$$

We now write

$$\begin{aligned} \left| \int_{\mathbf{R}^d} (f - f_n) \varphi dx \right| &\leq \left| \int_{\mathbf{R}^d} (f - f_n) \varphi (1 - \tau) dx \right| + \left| \int_{\mathbf{R}^d} (f - f_n) \varphi \tau dx \right| \\ &\leq \eta \|\varphi\|_{L^1} + \int_{\text{support } \tau} |f \varphi| dx + \left| \int_{\mathbf{R}^d} f_n \varphi \tau dx \right| \\ &\leq \delta + \delta + C \|f_n\|_{H^1} \|\varphi \tau\|_{\text{BMO}}. \end{aligned}$$

The proof of the theorem will therefore be established as soon as we verify

$$\|\varphi \tau\|_{\text{BMO}} \leq C \delta.$$

Since $\|\varphi \tau\|_{L^1} \leq C \eta \exp\{\delta^{-1}\} \leq \delta^{1+d}$, it is sufficient to test the BMO norm over cubes of sidelength $\leq \delta$. Let τ_Q and φ_Q denote the mean values of τ , φ over such a cube Q , and note that $|\varphi - \varphi_Q| \leq d\delta$ on Q . Therefore,

$$\begin{aligned} \frac{1}{|Q|} \int_Q |\varphi \tau - \varphi_Q \tau_Q| dx &\leq \frac{1}{|Q|} \int_Q |\varphi \tau - \varphi_Q \tau| dx + \frac{|\varphi_Q|}{|Q|} \int_Q |\tau - \tau_Q| dx \\ &\leq d\delta + \|\tau\|_{\text{BMO}} \leq C \delta. \end{aligned}$$

Standard limiting arguments show $\|f\|_{H^1} \leq 1$.

We remark that if f_n lie in the analytic Hardy space $H_a^1(\mathbf{R}^1)$, a proof can be given by using only H^∞ theory. By taking subsequences and looking at $\log|f_n|$, one can find $\varphi_m \in H^\infty$ such that $\|\varphi_m\|_{H^\infty} \leq 1$, $\|f_n \varphi_m (1 + |x|^2)\|_{L^\infty} \leq m$, and $\varphi_m(x) \rightarrow 1$ a.e. dx . Taking limits in n , $\|f \varphi_m\|_{H^1} \leq 1$, and then taking limits in m , $\|f\|_{H^1} \leq 1$. Returning to the original sequence, one applies Jensen's inequality to $f_n - f$ to establish that $f_n(z) - f(z) \rightarrow 0$. This suffices to prove $f_n \rightarrow f$ weakly.

Remark. The theorem is not abstract. In other words, if one attempts to use only the facts $\|f_n\|_{L^1} \leq 1$, $f_n(x) \rightarrow f(x)$ a.e. dx , and \mathcal{S} is dense in VMO, one will not arrive at a correct proof.

REFERENCES

1. R. R. Coifman and R. Rochberg, *Another characterization of BMO*, Proc. Amer. Math. Soc. **79** (1980), 249–254.
2. C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math. **129** (1972), 137–193.
3. E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, NJ, 1970.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CONNECTICUT 06520

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08544