

LIMITS OF DIFFERENTIABLE FUNCTIONS

UDAYAN B. DARJI

(Communicated by J. Marshall Ash)

ABSTRACT. Suppose that $\{f_n\}$ is a sequence of differentiable functions defined on $[0,1]$ which converges uniformly to some differentiable function f , and $\{f'_n\}$ converges pointwise to some function g . Let $M = \{x : f'(x) \neq g(x)\}$. In this paper we characterize such sets M under various hypotheses. It follows from one of our characterizations that M can be the entire interval $[0,1]$.

1. INTRODUCTION

We say that a sequence of differentiable functions $\{f_n\}$ defined on the interval $[0,1]$ is *proper* if $\{f_n\}$ converges uniformly to some differentiable function f and $\{f'_n\}$ converges pointwise to some function g . For such proper $\{f_n\}$, we let $\Delta(\{f_n\}) = \{x : f'(x) \neq g(x)\}$. It is a standard theorem in elementary analysis texts [6] that if $\{f_n\}$ is proper and $\{f'_n\}$ converges uniformly to some function g , then $\Delta(\{f_n\}) = \emptyset$. It is rather easy to construct an example of a proper $\{f_n\}$ where $\Delta(\{f_n\}) \neq \emptyset$. In this paper we investigate the following questions:

Question 1. *Is there a proper $\{f_n\}$ such that $\Delta(\{f_n\}) = [0, 1]$?*

Question 2. *Can $\{\Delta(\{f_n\}) : \{f_n\} \text{ is proper}\}$ be characterized?*

Theorems 1, 2, 3 and 5 answer Question 2 under various hypotheses. It will follow from Theorem 2 that Question 1 has an affirmative answer. However, Theorem 4 implies that in order to make $\Delta(\{f_n\}) = [0, 1]$, the derivatives have to be complicated in some sense.

We now state some definitions and background theorems. Recall that the *density topology* \mathcal{D} on \mathbb{R} is

$$\{M \subset \mathbb{R} : M \text{ is Lebesgue measurable and has density 1 at each of its points}\}.$$

Sets in \mathcal{D} are said to be open in density topology. Whenever we say that a set is open, closed, G_δ , F_σ , etc., we mean that it is open, closed, G_δ , F_σ , etc. in the ordinary topology on \mathbb{R} . Whenever we want a set to be open or closed in the density topology, we will specifically so state. A function $f : [0, 1] \rightarrow \mathbb{R}$ is *approximately*

Received by the editors April 20, 1994 and, in revised form, July 6, 1994; this paper was presented at the Special Session on Real Analysis at the 1992 AMS meeting in Baltimore.

1991 *Mathematics Subject Classification.* Primary 26A24, 26A21; Secondary 40A30.

Key words and phrases. F_σ , $G_\delta\sigma$, density topology, approximate continuity, nowhere measure dense.

This is the core part of the author's dissertation which was directed by Professor Jack B. Brown.

continuous iff the preimage of every open set under f is open in the density topology. A set $M \subset [0, 1]$ is *nowhere measure dense in the interval J* iff the interior of $M \cap J$ in the density topology is nowhere dense in J in the ordinary topology. We will freely use the following facts about density topology, approximate continuity and Lebesgue integration theory throughout the paper. Their proofs may be found in [1], [3], [2].

Fact 1. *If $M \subset \mathbb{R}$ is measurable, then there exists an F_σ set $N \subset M$ such that N is open in the density topology and $\lambda(M \setminus N)$, the Lebesgue measure of $M \setminus N$, is zero.*

Fact 2. *Every bounded approximately continuous function is a derivative.*

Fact 3 (Zahorski's Theorem [7]). *If G_0 and G_1 are two disjoint G_δ sets which are closed in the density topology, then there exists an approximately continuous function $f : [0, 1] \rightarrow [0, 1]$ such that $f^{-1}(i) = G_i$ for $i = 0, 1$.*

Fact 4. *If g is an integrable derivative defined on $[0, 1]$, then $f(x) = \int_0^x g$ is differentiable everywhere and $f'(x) = g(x)$ for all $x \in [0, 1]$.*

Fact 5. *Suppose $\{g_n\}$ is a sequence of integrable functions defined on the interval $[0, 1]$ such that $\{g_n\}$ is dominated by an L^1 function and $\{g_n\}$ converges pointwise to g . Let $f_n(x) = \int_0^x g_n$ and $f(x) = \int_0^x g$ for each $x \in [0, 1]$. Then $\{f_n\}$ converges uniformly to f .*

2. MAIN RESULTS

Lemma 1. *If $\{f_n\}$ is proper and $\{f'_n\}$ is dominated by an L^1 function, then $\Delta(\{f_n\})$ has measure zero.*

Proof. Without loss of generality assume that $f_n(0) = 0$ for all n . Let g be the pointwise limit of $\{f'_n\}$ and $h(x) = \int_0^x g$. Note that $h'(x) = g(x)$ for almost all $x \in [0, 1]$. By the Lebesgue Dominated Convergence Theorem we have that

$$f(x) = \lim_{n \rightarrow \infty} f_n(x) = h(x) \quad \text{for all } x \in [0, 1].$$

Thus $f' = g$ a.e. and $\Delta(\{f_n\})$ has measure zero. \square

Lemma 2. *Suppose $\{f_n\}$ is proper. Then $\Delta(\{f_n\})$ is $G_{\delta\sigma}$.*

Proof. Let g be the pointwise limit of $\{f'_n\}$. Since derivatives are of Baire class 1, g is in Baire class 2 and hence $\Delta(\{f_n\}) = (f' - g)^{-1}(R \setminus \{0\})$ is $G_{\delta\sigma}$ [4]. \square

Theorem 1 (General Dominated Case). *A set $M \subset [0, 1]$ is $G_{\delta\sigma}$ and of measure zero iff $M = \Delta(\{f_n\})$ for some proper $\{f_n\}$ where $\{f'_n\}$ is dominated by an L^1 function.*

Proof. (\Leftarrow) This direction follows from Lemmas 1 and 2.

(\Rightarrow) The proof of this direction has a flavor similar to a result of Preiss [5]. Let $M = \bigcup G_k$ be a $G_{\delta\sigma}$ set of measure zero where each G_k is G_δ . Now for each positive integer k , let $\{U_{k,n}\}_{n=1}^\infty$ be a decreasing sequence of open sets such that $G_k = \bigcap_{n=1}^\infty U_{k,n}$. For each n and k we may obtain by Fact 3 an approximately continuous function $h_{k,n} : [0, 1] \rightarrow [0, 1]$ such that $h_{k,n}^{-1}(1) = G_k$ and $h_{k,n}^{-1}(0) = (U_{k,n})^c$, the complement of $U_{k,n}$. Note that for each k , $\{h_{k,n}\}_{n=1}^\infty$ converges pointwise to χ_{G_k} ,

the characteristic function of G_k . Now set

$$g_n = \sum_{k=1}^{\infty} \frac{1}{2^k} h_{k,n} \quad \text{and} \quad g = \sum_{k=1}^{\infty} \frac{1}{2^k} \chi_{G_k}.$$

It follows that g_n is a bounded derivative because it is the uniform sum of a sequence of bounded approximately continuous functions. Also note that $\{g_n\}$ converges pointwise to g .

Now let $f_n(x) = \int_0^x g_n$. That $f'_n(x) = g_n(x)$ for all $x \in [0, 1]$ follows from Fact 4. We also have that $\int_0^x g = 0$ for all $x \in [0, 1]$ as g is nonzero only on the measure zero set M . Since $\{g_n\}$ is a uniformly bounded sequence of integrable functions which converges pointwise to g , by Fact 5 it follows that $\{f_n\}$ converges uniformly to the zero function. But $\{f'_n\}$ converges pointwise to the function g which is nonzero precisely on set M . Therefore, $\Delta(\{f_n\}) = M$. \square

Lemma 3. *Let $M \subset [0, 1]$ be a nonempty F_σ set which is open in the density topology, and let $A > 0$. Then, there exists a bounded approximately continuous function $f : [0, 1] \rightarrow \mathbb{R}$ such that*

- (1) $\int_0^1 f = A$,
- (2) $f^{-1}(0) = M^c$, and
- (3) $f \geq 0$.

Proof. By Fact 3 there is a bounded approximately continuous function h which satisfies conditions (2) and (3). Let $f = c \cdot h$, where $c = \frac{A}{\int_0^1 h}$. This f satisfies all three required conditions. \square

Lemma 4. *If M is F_σ , open in the density topology and M^c is dense in $[0, 1]$, then there exists proper $\{f_n\}$ such that $M = \Delta(\{f_n\})$.*

Proof. Suppose M is F_σ , open in the density topology, and M^c is dense in $[0, 1]$. Let g be an approximately continuous function such that $0 \leq g(x) \leq 1$ for all x and $g^{-1}(0) = M^c$. Write $M = \bigcup_{i=1}^{\infty} F_i$ where each F_i is a nowhere dense closed subset of $[0, 1]$. Since each F_i is a nowhere dense closed subset of $[0, 1]$ and M is open in the density topology, it follows that for each interval J for which $M \cap J \neq \emptyset$ we have $(M \cap J) \setminus (\bigcup_{i=1}^n F_i) \neq \emptyset$ for all n .

We now construct a sequence of derivatives $\{g_n\}$ in the following fashion. Fix n . Let $N_n = M^c \cup F_1 \cup \dots \cup F_n$. Observe that N_n is a G_δ set and is also closed in the density topology. Let $\{x(n, k)\}_{k=1}^{m(n)+1}$ be a partition of $[0, 1]$ such that $x(n, i) \in M^c$ and $|x(n, i) - x(n, i + 1)| < \frac{1}{2^n}$ for each $1 \leq i \leq m(n)$. Fix $1 \leq i \leq m(n)$. Observe that $[x(n, i), x(n, i + 1)] \setminus N_n$ is either a nonempty F_σ set which is open in the density topology or $[x(n, i), x(n, i + 1)] \subset M^c$. If the latter is the case, let $h_{n,i}$ be the zero function on $[x(n, i), x(n, i + 1)]$ and we have that $\int_{x(n,i)}^{x(n,i+1)} h_{n,i} = \int_{x(n,i)}^{x(n,i+1)} g$. Otherwise using Lemma 3, obtain a bounded, nonnegative approximately continuous function $h_{n,i}$ defined on $[x(n, i), x(n, i + 1)]$ such that $h_{n,i}^{-1}(0) = N_n \cap [x(n, i), x(n, i + 1)]$ and $\int_{x(n,i)}^{x(n,i+1)} h_{n,i} = \int_{x(n,i)}^{x(n,i+1)} g$. Now, let h_n be the union of $h_{n,1}, \dots, h_{n,m(n)}$. Then, h_n is bounded and approximately continuous. Let $g_n = g - h_n$. As $\{h_n\}$ converges pointwise to the zero function, $\{g_n\}$ converges pointwise to g .

Let $f_n(x) = \int_0^x g_n$. Since g_n is bounded and approximately continuous, $f'_n = g_n$. Let us next show that $\|f_n\|$, the sup norm of f_n , is less than 2^{-n+1} . Let $x \in [0, 1]$

and i be such that $x \in [x(n, i), x(n, i + 1)]$. Then

$$\begin{aligned} |f_n(x)| &= \left| \int_0^x g_n \right| \leq \left| \int_0^{x(n,i)} g - h_n \right| + \left| \int_{x(n,i)}^x g - h_n \right| \\ &\leq 0 + \left| \int_{x(n,i)}^x g \right| + \left| \int_{x(n,i)}^x h_n \right| < 2^{-n} + 2^{-n} = 2^{-n+1}. \end{aligned}$$

The above estimate on $\|f_n\|$ implies that $\{f_n\}$ converges uniformly to the zero function. We also know that $\{g_n\}$ converges pointwise to g and $g^{-1}(0) = M^c$. Therefore, $M = \Delta(\{f_n\})$. □

Theorem 2 (General Nondominated Case). *A set $M \subset [0, 1]$ is $G_{\delta\sigma}$ iff there exists proper $\{f_n\}$ such that $M = \Delta(\{f_n\})$.*

Proof. (\Leftarrow) This direction follows from Lemma 2.

(\Rightarrow) Let M be $G_{\delta\sigma}$. From Fact 1 obtain two disjoint sets M_1 and M_2 such that $M_1 \cup M_2 = M$, M_1 is $G_{\delta\sigma}$ set of measure zero and M_2 is an F_σ set which is open in the density topology and M_2^c is dense in $[0,1]$. By Theorem 1 and Lemma 4, obtain proper sequences $\{f_n\}$ and $\{h_n\}$ such that $M_1 = \Delta(\{f_n\})$ and $M_2 = \Delta(\{h_n\})$. Then $\{f_n + h_n\}$ is proper and $M = \Delta(\{f_n + h_n\})$. □

Lemma 5. *Suppose $\{f_n\}$ is proper and for all n , $f_n \in C^1$, i.e. f'_n is continuous. Then $\Delta(\{f_n\})$ is F_σ .*

Proof. Let g be the pointwise limit of $\{f'_n\}$. Then g and f' are of Baire class 1. Therefore, $\Delta(\{f_n\}) = (f' - g)^{-1}(\mathbb{R} \setminus \{0\})$ is F_σ [4]. □

Theorem 3 (Dominated C^1 Case). *A set $M \subset [0, 1]$ is F_σ and of measure zero iff $M = \Delta(\{f_n\})$ for some proper $\{f_n\}$ where $f_n \in C^1$ for all n and $\{f'_n\}$ is dominated by an L^1 function.*

Proof. (\Leftarrow) This direction follows from Lemmas 1 and 5.

(\Rightarrow) Suppose M is F_σ and of measure zero. Let $M = \bigcup_{k=1}^\infty M_k$ where each M_k is closed. Let $\{G_{k,n}\}$ be such that each $G_{k,n}$ is a finite collection of closed intervals and

- (1) $\bigcup G_{k,n} \subset \bigcup G_{k,n+1}$, and
- (2) $\bigcup_{n=1}^\infty G_{k,n} = M_k^c$.

Now let $h_{k,n}$ be a continuous function defined on $[0,1]$ such that $0 \leq h_{k,n}(x) \leq 1$, $h_{k,n}(M_k) = 1$, and $h_{k,n}(\bigcup G_{k,n}) = 0$. Note that $\{h_{k,n}\}_{n=1}^\infty$ converges pointwise to χ_{M_k} , the characteristic function of M_k . Now set $g_n = \sum_{k=1}^\infty 2^{-k} \cdot h_{k,n}$ and $g = \sum_{k=1}^\infty 2^{-k} \cdot \chi_{M_k}$. Observe that g_n is a continuous function for all n , $0 \leq g_n(x) \leq 1$ and $\{g_n\}$ converges pointwise to g .

Setting $f_n(x) = \int_0^x g_n$, we have that $f'_n(x) = g_n(x)$ for all $x \in [0, 1]$. We also have that $\int_0^x g = 0$ for all $x \in [0, 1]$ as g is nonzero only on the measure zero set M . Since $\{g_n\}$ is a uniformly bounded sequence of continuous functions which converges pointwise to g , by Fact 5 it follows that $\{f_n\}$ converges uniformly to the zero function. But $\{f'_n\}$ converges pointwise to function g which is nonzero precisely on set M . Therefore, $\Delta(\{f_n\}) = M$. □

Lemma 6. *Suppose $M \subset [0, 1]$ is F_σ and nowhere measure dense. Then M is the union of two disjoint F_σ sets, one of which is of measure zero and the other nowhere dense.*

Proof. Let B_1, B_2, \dots be a countable basis for $[0, 1]$. Let $U = \bigcup\{B_i : \lambda(B_i \cap M) = 0\}$. Since M is nowhere measure dense, U is a dense open subset of $[0, 1]$. Let $M_1 = M \cap U$ and $M_2 = M \setminus U$. Then M_1 and M_2 are the desired sets. \square

Lemma 7. *Let $\{g_n\}$ be a sequence of a.e. continuous functions whose domain is interval I . Suppose $\{g_n\}$ converges pointwise to some function g on I . Then there exists an interval $J \subset I$ and $K > 0$, such that for all n , $|g_n(x)| < K$ for a.e. $x \in J$.*

Proof. We will prove this lemma by contradiction. Assume the hypothesis and that there is no $J \subset I$ such that $\{g_n\}$ is bounded a.e. on J . This implies that for every $K > 0$ and interval $J \subset I$, there are infinitely many integers n such that $|g_n| > K$ on a positive measure set contained in J . Using this observation and the fact that $\{g_n\}$ is a sequence of a.e. continuous functions, we may obtain a subinterval I_1 of I and a positive integer $n(1)$ such that $|g_{n(1)}| > 1$ on I_1 . Proceeding in a similar fashion we may obtain a decreasing sequence of closed intervals $\{I_k\}$ and an increasing sequence of positive integers $\{n(k)\}$ such that for each k , $|g_{n(k)}| > k$ on I_k . Now let $p \in \bigcap_{i=1}^\infty I_i$. Then $\{g_{n(k)}(p)\}_{k=1}^\infty$ does not converge, contradicting the hypothesis and concluding the proof of the lemma. \square

Theorem 4. *Suppose $\{f_n\}$ is proper, f'_n is integrable and the set of discontinuities of f'_n has measure zero for all n . Then $\Delta(\{f_n\})$ is nowhere measure dense.*

Proof. Let I be a subinterval of $[0, 1]$. Using Lemma 7 obtain an interval $J \subset I$ such that $\{f'_n\}$ is bounded a.e. on J . By Lemma 1 we have that $\Delta(\{f_n\}) \cap J$ has measure zero. This implies that the interior of $\Delta(\{f_n\})$ in the density topology is nowhere dense in $[0, 1]$. \square

Lemma 8. *Suppose M is an F_σ nowhere dense subset of $[0, 1]$. Then $M = \Delta(\{f_n\})$ for some proper $\{f_n\}$ where $f_n \in C^1$ for all n .*

Proof. Let $U = (\text{cl}(M))^c$ and F_1, F_2, \dots be a pairwise disjoint decomposition of M into closed sets. Let g be the function which is zero on M^c and 2^{-i} on F_i . Note that g is of Baire class 1. Let $\{G_n\}$ be such that each G_n is a finite collection of closed intervals and

- $\bigcup G_n \subset \bigcup G_{n+1}$, and
- $\bigcup_{n=1}^\infty (\bigcup G_n) = U$.

Using the fact that g is of Baire class 1, obtain a sequence of continuous functions $\{h_n\}$ such that $\{h_n\}$ converges pointwise to g , and for all n , $h_n(\bigcup G_n) = 0$ and $0 \leq h_n(x) \leq 1$.

We now construct g_n in the following manner. First, let $\{x(n, k)\}_{k=1}^{m(n)+1}$ be a partition of $[0, 1]$ such that $|x(n, k) - x(n, k + 1)| < 2^{-n}$ for $k = 1, 2, \dots, m(n)$. Now for each $1 \leq k \leq m(n)$, let $a_{n,k}$ be a continuous nonnegative function defined on $[x(n, k), x(n, k + 1)]$ such that

- $\int_{x(n,k)}^{x(n,k+1)} a_{n,k} = \int_{x(n,k)}^{x(n,k+1)} h_n$,
- $a_{n,k} = 0$ on $\bigcup G_n \cap [x(n, k), x(n, k + 1)]$,
- $a_{n,k}^{-1}(\mathbb{R} \setminus \{0\}) \subset U$, and
- $a_{n,k}(x(n, k)) = a_{n,k}(x(n, k + 1)) = 0$.

Let a_n be the union of $a_{n,1}, a_{n,2}, \dots, a_{n,m(n)}$ and $g_n = h_n - a_n$. As $\{a_n\}$ is a sequence of continuous functions which converges pointwise to the zero function, $\{g_n\}$ is a sequence of continuous functions which converges pointwise to g .

For each n , let $f_n(x) = \int_0^x g_n$. Observe that for $1 \leq k \leq m(n) + 1$

$$f_n(x(n, k)) = \int_0^{x(n, k)} g_n = \sum_{i=1}^{k-1} \int_{x(n, i)}^{x(n, i+1)} h_n - a_n = 0.$$

Using this observation we obtain an estimate on $\|f_n\|$. Let $x \in [0, 1]$. Let k be a positive integer such that $x \in [x(n, k), x(n, k + 1)]$.

$$\begin{aligned} |f_n(x)| &\leq |f_n(x(n, k))| + |f_n(x) - f_n(x(n, k))| \\ &= 0 + \left| \int_{x(n, k)}^x g_n \right| = \left| \int_{x(n, k)}^x h_n - a_n \right| \\ &\leq \left| \int_{x(n, k)}^x h_n \right| + \left| \int_{x(n, k)}^x a_n \right| \\ &< 2^{-n} + 2^{-n} = 2^{-n+1}. \end{aligned}$$

From above we have that $\{f_n\}$ converges uniformly to the zero function. As $\{f'_n\}$ converges pointwise to g and $g^{-1}(0) = M^c$, we have that $M = \Delta(\{f_n\})$. \square

Theorem 5 (Nondominated C^1 Case). *A set $M \subset [0, 1]$ is F_σ and nowhere measure dense iff $M = \Delta(\{f_n\})$ for some proper $\{f_n\}$ where $f_n \in C^1$ for all n .*

Proof. (\Leftarrow) This direction follows from Theorem 4 and Lemma 5.

(\Rightarrow) Suppose M is F_σ and nowhere measure dense. Then by Lemma 6, $M = M_1 \cup M_2$ where M_1 and M_2 are disjoint F_σ sets, one of which is nowhere dense and the other of measure zero. By Theorem 3 and Lemma 8, there are proper sequences $\{f_n\}$ and $\{h_n\}$ such that $\Delta(\{f_n\}) = M_1$ and $\Delta(\{h_n\}) = M_2$. Then $\{f_n + h_n\}$ is proper and $M = \Delta(\{f_n + h_n\})$. \square

The author thanks the referee for making helpful suggestions which improved the exposition of the paper.

REFERENCES

1. A. M. Bruckner, *Differentiation of real functions*, Lecture Notes in Math., vol. 659, Springer-Verlag, New York, 1978. MR **80h**:26002
2. K. Ciesielski, L. Larson, and K. Ostaszewski, *\mathcal{I} -Density continuous functions*, Mem. Amer. Math. Soc., no. 515, vol. 107, Amer. Math. Soc., Providence, RI, 1994. MR **94f**:54035
3. R. Jeffery, *The theory of function of a real variable*, Mathematical Exposition No. 6, University of Toronto Press, Ontario, 1951. MR **13**:2166
4. K. Kuratowski, *Topology*, Vol. I, Academic Press, New York, 1966. MR **36**:840
5. D. Preiss, *Limits of approximately continuous functions*, Czechoslovak Math. J. **21** (1971), 371–372. MR **44**:4154
6. W. Rudin, *Principles of mathematical analysis*, third edition, McGraw-Hill Inc., New York, 1976. MR **52**:5893
7. Z. Zahorski, *Sur la premiere derivee*, Trans. Amer. Math. Soc. **69** (1950), 1–54. MR **12**:247c

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LOUISVILLE, LOUISVILLE, KENTUCKY 40292
E-mail address: ubdarj01@homer.louisville.edu