

## POLYNOMIAL RINGS OVER GOLDIE-KERR COMMUTATIVE RINGS II

CARL FAITH

(Communicated by Wolmer V. Vasconcelos)

*In memory of Pere Menal*

ABSTRACT. An overlooked corollary to the main result of the stated paper (Proc. Amer. Math. Soc. **120** (1994), 989–993) is that any Goldie ring  $R$  of Goldie dimension 1 has Artinian classical quotient ring  $Q$ , hence is a Kerr ring in the sense that the polynomial ring  $R[X]$  satisfies the *acc* on annihilators ( $= acc\perp$ ).

More generally, we show that a Goldie ring  $R$  has Artinian  $Q$  when every zero divisor of  $R$  has essential annihilator (in this case  $Q$  is a local ring; see Theorem 1').

A corollary to the proof is Theorem 2: A commutative ring  $R$  has Artinian  $Q$  iff  $R$  is a Goldie ring in which each element of the Jacobson radical of  $Q$  has essential annihilator.

Applying a theorem of Beck we show that any  $acc\perp$  ring  $R$  that has Noetherian local ring  $R_P$  for each associated prime  $P$  is a Kerr ring and has Kerr polynomial ring  $R[X]$  (Theorem 5).

### INTRODUCTION

Throughout,  $R$  denotes a commutative ring.

It is convenient to state the corollary in generalized form as follows:

1. **Theorem.** *If  $R$  is a Goldie ring in which each zero divisor  $x$  has essential annihilator  $x^\perp$ , then  $R$  has Artinian quotient ring  $Q$ .*

In this case, it follows from Small's theorem [S] that  $R[X]$  has Artinian quotient ring.

A ring  $R$  has *finite Goldie* (or uniform) *dimension*  $n$  if  $n$  is the maximal number of nonzero ideals in a direct sum contained in  $R$ . Furthermore,  $R$  is *Goldie* if  $R$  has  $acc\perp$  and finite Goldie dimension. *Uniform ring* is another term for a ring with Goldie dimension 1, equivalently, 0 is an irreducible ideal. The *singular ideal* of  $R$  is the set

$$Z(R) = \{x \in R \mid x^\perp \text{ is essential}\}.$$

Then  $Z(R)$  is contained in the set  $z(R)$  of zero divisors. Obviously, when  $R$  is uniform, we have that  $z(R) = Z(R)$ , and then  $Z(Q)$  is the set of non-units of  $Q$ .

We can sharpen Theorem 1 in this terminology:

---

Received by the editors April 25, 1994 and, in revised form, August 5, 1994.

1991 *Mathematics Subject Classification*. Primary 13B25, 13CO5, 13EO5, 13H99, 13J10; Secondary 16D90, 16P60, 16S50.

1'. **Theorem.** *If  $R$  is a Goldie ring, and if  $z(R) \subseteq Z(R)$ , then  $Q$  is an Artinian local ring (hence  $R$  is Kerr). Conversely.<sup>1</sup>*

*Proof.*  $Q$  also is a Goldie ring (see [F3]), and  $Z(Q)$  is a nil ideal in any  $acc\perp$  ring (see, e.g., [F1]). Moreover, every element  $x$  in the Jacobson radical  $J(Q)$  is a nonunit; hence  $x = rs^{-1}$ , where  $r \in z(R) \subseteq Z(R)$  and  $s \in R^*$ . Then,  $x$  has essential right annihilator  $x^\perp$  in  $Q$ , since  $r^\perp \cap Q$  is essential in  $R$ , a fact that shows that  $J(Q) \subseteq Z(Q)$ . Thus,  $J(Q)$  is nil, so  $Q$  is Artinian by Theorem 1.1 of [F2]. Since  $J(Q)$  is the set of non-units of  $Q$ , it follows that  $Q$  is local.

The converse hinges on the fact that if  $Q$  is Artinian, then  $Q$  is Noetherian and hence Goldie, so  $R$  is Goldie. Furthermore,  $J(Q)$  is nilpotent, and every nilpotent element  $x$  has essential annihilator. (Let  $I$  be any nonzero ideal, and  $x^n = 0$  where  $x^{n-1} \neq 0$ . If  $x^\perp \cap I = 0$ , then  $xI \neq 0$ . Suppose  $i$  is least such that  $x^i I \neq 0$ . Then  $x^\perp \cap I \supseteq x^i I \neq 0$ , a contradiction which shows that  $x^\perp$  is an essential ideal.) Since  $Z(R)$  is nilpotent in an  $acc\perp$  ring (loc. cit.), then  $J(Q) = Z(Q)$ . Since  $Q$  is local and  $J(Q)$  is nilpotent, then every zero divisor  $x$  of  $R$  lies in

$$J(Q) \cap R = Z(Q) \cap R = Z(R),$$

so  $z(R) \subseteq Z(R)$  as needed.  $\square$

The proof of Theorem 1' has the corollary.

2. **Theorem.** *A ring  $R$  has Artinian  $Q$  iff  $R$  is a Goldie ring and the Jacobson radical of  $Q$  coincides with its singular ideal, that is,  $J(Q) = Z(Q)$ .*

*Proof.* If  $J(Q) = Z(Q)$ , then  $R acc\perp \Rightarrow J(Q)$  is nil, so  $R$  Goldie  $\Rightarrow Q$  is Artinian by Theorem 1.1 of [F2]. Conversely,  $Q$  Artinian  $\Rightarrow J(Q)$  is nil, hence  $J(Q) \subseteq Z(Q)$ . But  $Z(Q)$  is nil in an  $acc\perp$  ring, hence  $Z(Q) = J(Q)$ .  $\square$

In a uniform ring every nonzero ideal is essential, so the theorems each imply that any uniform  $acc\perp$  ring  $R$  has Artinian  $Q$ , but  $Q$  is in fact then quasi-Frobenius since  $Q$  has simple socle. With this fact as a motivator, we next derive a more general theorem with the same conclusion (Theorem 2).

A ring  $R$  is  $F$ -injective (=  $\aleph_0$ -injective) if every map  $I \rightarrow R$  of a finitely generated ideal  $I$  is extendable to  $R \rightarrow R$ . (Any  $FP$ -injective ring  $R$  is  $F$ -injective; cf. [F3], p. 189.) Any  $F$ -injective ring  $R$  coincides with its quotient ring  $Q$ . Any valuation ring  $R$  has  $FP$ -injective  $Q$  by a theorem of Facchini ([F-P], p. 96, Corollary 6-10; cf. [F-F]).

3. **Theorem.** *If  $R$  is an  $acc\perp$  ring with  $F$ -injective (e.g.  $FP$  or self-injective) quotient ring  $Q$ , then  $R$  is Kerr, in fact  $Q$  is quasi-Frobenius (=  $QF$ ).*

*Proof.* Every finitely generated ideal  $I$  of an  $F$ -injective ring  $R$  is an annihilator (see, e.g., [F3], p. 189, Prop. 23.21.2). The  $acc\perp$  in  $R$  implies the  $acc\perp$  in  $Q$ , and hence  $Q$  satisfies the  $acc$  on finitely generated ideals, so  $Q$  is Noetherian. But a Noetherian  $F$ -injective ring is self-injective, hence  $QF$ .  $\square$

4. **Corollary.** *Any uniform  $acc\perp$  ring  $R$ , e.g. any  $acc\perp$  valuation ring, is a Kerr ring. Furthermore  $Q$  is Artinian in fact  $QF$ .*

*Proof.*  $Q$  is Artinian by Theorem 1, and has Goldie  $\dim = 1$ , hence has simple socle, which by classical ideal theory (cf. Corollary 2 of [F1]) implies that  $Q$  is  $QF$ .  $\square$

<sup>1</sup>Classically, it is known that a ring  $R$  has local  $Q$  iff the set  $z(R)$  is an ideal  $P$ . In this case,  $P$  is a prime ideal and  $Q = R_P$  is the local ring at  $P$ . Any Artinian ring  $R$  is a finite product of local Artinian rings. See Theorem 2 in this connection.

WHEN IS  $R[X]$  KERR?

We raised the question in [F2]: If  $R$  is Kerr, is  $R[X]$ ?<sup>2</sup>

We cited some obvious examples in [F2], e.g. any subring of a Noetherian ring, and mentioned the Camillo-Guralnick theorem which yields an affirmative answer for an algebra over an uncountable field. We next show that Beck's theorem [B] yields another affirmative answer.

**5. Theorem.** *If  $R$  is an  $\text{acc}\perp$  ring and if  $R_P$  is Noetherian for every associated prime  $P$ , then the same is true of  $R[X]$ . Furthermore, both  $R$  and  $R[X]$  have (flat) embeddings into Noetherian rings, hence each is a Kerr ring.*

*Proof.* In any  $\text{acc}\perp$  ring  $R$ , the set  $\text{Ass}R$  of associated prime ideals is finite (see [F4], Corollary 3.7 and Theorem 3.6), and obviously  $\bigcup_{P \in \text{Ass}R} P$  is the set  $z(R)$  of zero divisors (i.e., every  $x \in z(R)$  is contained in some  $P \in \text{Ass}R$ ). We can now apply Beck's theorem ([B], Theorem 5.1) to conclude that  $R$  has a flat embedding in a Noetherian ring  $T$ , and hence both  $R$  and  $R[X]$  are Kerr, since  $R[X]$  is contained in a Noetherian ring  $T[X]$ .

Next, contraction induces a 1-1 correspondence  $\text{Ass}R[X] \rightarrow \text{Ass}R$  under various conditions including  $\text{acc}\perp$  in  $R$  ([F4], Theorem 3.12 B; any  $\text{acc}\perp$  ring is trivially a zip ring in the terminology employed there).

Thus, if  $P \in \text{Ass}R[X]$ , then

$$P_0 = P \cap R \in \text{Ass}R$$

and

$$R[X]_P = R_{P_0}[X]_{PR_P[X]} = R_M[X]_{PR_M[X]}$$

which holds for any prime ideal  $P$  of  $R[X]$ , and  $M$  any maximal ideal containing  $P_0 = P \cap R$ . (See, e.g., [H], p.73, Lemma 13.1.)

In particular, this shows that

$$R_{P_0} \text{ Noetherian} \Rightarrow R[X]_P \text{ Noetherian,}$$

so  $R[X]$  has the stated property (this also follows from the theorem of Beck since  $R[X] \hookrightarrow T[X]$  is a flat embedding in a Noetherian ring).  $\square$

Since a Kerr ring need not embed in a Noetherian ring, this shows that a Kerr ring  $R$  does not in general localize to Noetherian rings at associated primes.

**6. Corollary.** *If  $R$  satisfies the hypothesis of Theorem 5, then so does the infinite polynomial ring  $S = R[x_1, \dots, x_n, \dots]$ , that is,  $S$  has a flat embedding in a Noetherian ring, hence is Kerr.*

*Proof.* By Theorem 5,  $R$  has a flat embedding in a Noetherian ring  $T$ , and by a remark of D. D. Anderson (cited in [C], p. 75, Remark 2),  $A = T[x_1, \dots, x_n, \dots]$  localized at the ideal  $P$  consisting of all polynomials in  $A$  of content 1 is a Noetherian ring. Thus, since  $A \hookrightarrow A_P$  is a flat embedding in a Noetherian ring,  $S \hookrightarrow A_P$  is, also.  $\square$

<sup>2</sup>In the meanwhile Cedó and Herbera have found a ring  $R$  over which the polynomial ring in  $n$  variables is Kerr but that in  $n + 1$  variables is not (Fax of November 1994). See [C-H].

## NOTE ADDED IN PROOF

The author has discovered an error in the proof of Theorem 2.2 in [F2]. The first sentence should read:

“If  $I$  is an ideal of  $R$ , then  $I$  is an annihilator of  $R$  iff  $IQ$  is an annihilator of  $Q$  and  $IQ \cap R = I$ .”

The third sentence should read:

“This also implies that if  $K$  is an ideal of  $Q$ , then  $K \cap R \in \text{Ass } R$  iff  $K \in \text{Ass } Q$ .”

## REFERENCES

- [B] I. Beck,  $\Sigma$ -injective modules, *J. Algebra* **21** (1972), 232–249. MR **50**:9967
- [C] V. Camillo, *Coherence for polynomial rings*, *J. Algebra* **132** (1990), 72–76. MR **91c**:16018
- [C-H] F. Cedó and D. Herbera, *On polynomial rings over Kerr commutative rings*, preprint, U. Autónoma de Barcelona, 1995.
- [F-F] A. Facchini and C. Faith, *FP-injective quotient rings and elementary divisor rings*, Proceedings of the Fez Conference on Commutative Algebra (1995), Lecture Notes in Pure and Appl. Math., Marcel Dekker, New York and Basel, 1996.
- [F1] C. Faith, *Finitely embedded commutative rings*, *Proc. Amer. Math. Soc.* **112** (1991), 657–659. MR **93f**:13012
- [F2] ———, *Polynomial rings over Goldie-Kerr commutative rings*, *Proc. Amer. Math. Soc.* **120** (1994), 989–993. MR **94k**:13024
- [F3] ———, *Algebra II: Ring theory*, Springer-Verlag, Berlin, Heidelberg, and New York, 1976. MR **55**:383
- [F4] ———, *Annihilators, associated primes and Kasch-McCoy quotient rings of commutative rings*, *Comm. Algebra* **119** (1991), 1867–1892. MR **92g**:16008
- [F-P] C. Faith and P. Pillay, *Classification of commutative FPF rings*, *Notas Mat.*, vol. 4, Univ. Murcia, Murcia.
- [H] J. Huckaba, *Commutative rings with zero divisors*, Monographs Pure Appl. Math., Marcel Dekker, Basel and New York, 1988. MR **89e**:13001
- [K1] J. W. Kerr, *The polynomial ring over a Goldie ring need not be a Goldie ring*, *J. Algebra* **134** (1990), 344–352. MR **91h**:16042
- [K2] ———, *An example of a Goldie ring whose matrix ring is not Goldie*, *J. Algebra* **61** (1979), 590–592. MR **81b**:16016
- [S] L. Small, *Orders in Artinian rings*, *J. Algebra* **4** (1966), 13–41. MR **34**:199

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY 08903

*Permanent address:* 199 LONGVIEW DRIVE, PRINCETON, NEW JERSEY 08540