

## INVARIANT SUBSPACES OF THE HARMONIC DIRICHLET SPACE WITH LARGE CO-DIMENSION

WILLIAM T. ROSS

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. In this paper, we comment on the complexity of the invariant subspaces (under the bilateral Dirichlet shift  $f \rightarrow \zeta f$ ) of the harmonic Dirichlet space  $D$ . Using the sampling theory of Seip and some work on invariant subspaces of Bergman spaces, we will give examples of invariant subspaces  $\mathcal{F} \subset D$  with  $\dim(\mathcal{F}/\zeta\mathcal{F}) = n$ ,  $n \in \mathbb{N} \cup \{\infty\}$ . We will also generalize this to the Dirichlet classes  $D_\alpha$ ,  $0 < \alpha < \infty$ , as well as the Besov classes  $B_p^\alpha$ ,  $1 < p < \infty$ ,  $0 < \alpha < 1$ .

### 1. INTRODUCTION

For a space  $X$  of functions on the unit circle  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$  for which the shift operator

$$T : X \rightarrow X, \quad (Tf)(\zeta) = \zeta f(\zeta)$$

is continuous and invertible (e.g.  $L^p(\mathbb{T}, |d\zeta|)$ ,  $C^n(\mathbb{T})$ ,  $C^\infty(\mathbb{T})$ ,  $W_n^p(\mathbb{T})$  the Sobolev classes,  $B_p^\alpha(\mathbb{T})$  the Besov classes,  $D_\alpha$  the Dirichlet classes,  $BL_{p,\theta}^l(\mathbb{T})$  the Triebel-Lizorkin classes), the problem of characterizing the subspaces (closed linear manifolds)  $\mathcal{F} \subset X$  for which  $T\mathcal{F} \subset \mathcal{F}$ , the so called *invariant subspaces*, is a very difficult and open problem. There are two types of invariant subspaces to consider: *simply invariant* (or 1-invariant)  $T\mathcal{F} \neq \mathcal{F}$ , and *2-invariant*  $T\mathcal{F} = \mathcal{F}$ . The 2-invariant subspaces are often described by their zero sets in  $\mathbb{T}$  [1], [10], [11], [12], while the 1-invariant subspaces are known to be much more complicated and for most of the classes mentioned above, a complete characterization seems a long way off [3], [6], [8], [9].

In this paper, we focus our attention on the *harmonic Dirichlet space*  $D$  of functions  $f \in L^2(\mathbb{T}, |d\zeta|)$  with finite norm

$$\|f\|^2 = \sum_{n=-\infty}^{\infty} (1 + |n|) |\hat{f}(n)|^2,$$

where  $\{\hat{f}(n)\}$  are the Fourier coefficients of  $f$ . The 2-invariant subspaces of  $D$  can be characterized by their zero sets on  $\mathbb{T}$  [12], while the simply invariant subspaces of  $D$  are much more complicated. In this paper, we remark on the complexity of the invariant subspaces  $\mathcal{F}$  with  $D_A \subset \mathcal{F} \subset D$ , where  $D_A$  is the analytic Dirichlet

---

Received by the editors October 31, 1994 and, in revised form, December 9, 1994.

1991 *Mathematics Subject Classification*. Primary 30H05; Secondary 30C15.

*Key words and phrases*. Dirichlet spaces, invariant subspaces, co-dimension, Bergman spaces.

This research was supported in part by a grant from the National Science Foundation.

space  $\{f \in D : \hat{f}(n) = 0 \ \forall n < 0\}$ , by examining their *index*. Here for an invariant subspace  $\mathcal{F} \subset D$ , we define the *index* (or sometimes called the *co-dimension*) to be

$$\text{ind } \mathcal{F} = \dim(\mathcal{F}/\zeta\mathcal{F}).$$

For the analytic Dirichlet space  $D_A$ , every non-zero invariant subspace has index 1 [13]. In stark contrast to this, it was observed in [1] that there are examples of invariant subspaces  $D_A \subset \mathcal{F} \subset D$  with  $\text{ind } \mathcal{F} = n$  for any  $n \in \mathbb{N} \cup \{\infty\}$ . We will give a new and more direct proof of this fact as well as specific examples of these types of subspaces. To do this, we will employ the natural one-to-one correspondence between the invariant subspaces  $D_A \subset \mathcal{F} \subset D$  and the invariant subspaces (under  $f \rightarrow zf$ ) of the Bergman space  $L^2_a(\mathbb{D})$ . We will then use some recent results of Seip, Hedenmalm, and Richter [4], [5], [14] which yield specific examples of invariant subspaces of the Bergman space with large index. We then transfer this information back to the Dirichlet space to obtain the following theorem.

**Theorem 1.1.** *Given  $n \in \mathbb{N} \cup \{\infty\}$ , there are sequences  $A_j$ ,  $0 \leq j < n$ , of points in  $\mathbb{D}$  such that*

$$\mathcal{F} = \bigcap_{0 \leq j < n} \text{span}\{\overline{B_a}H^2 \cap D : a \in A_j\}$$

*is a simply invariant subspace of  $D$  with  $\text{ind } \mathcal{F} = n$ .*

Note that  $H^2 = \{f \in L^2 : \hat{f}(n) = 0 \ \forall n < 0\}$  is the usual Hardy space on the circle and

$$(1.1) \quad B_a(\zeta) = \frac{\bar{a}}{|a|} \frac{a - \zeta}{1 - \bar{a}\zeta}, \quad \zeta \in \mathbb{T}.$$

We also use  $\text{span } X$  to denote the closed linear span of  $X$ , and  $\bar{z}$  to denote the complex conjugate of a complex number  $z$ .

We will remark at the end that a similar result also holds for the Dirichlet spaces  $D_\alpha$  and the Besov classes  $B_p^\alpha(\mathbb{T})$ .

## 2. BERGMAN SPACES

The sequences  $A_j$  in Theorem 1.1 will be zero sequences of the Bergman space, and thus we begin our discussion with some basic Bergman space facts. The *Bergman space*  $L^2_a(\mathbb{D})$  is the space of analytic functions  $f$  on  $\mathbb{D}$  such that

$$\int_{\mathbb{D}} |f(z)|^2 dA(z) = \sum_{n=0}^{\infty} \frac{|a_n|^2}{n+1} < \infty.$$

Here  $dA$  is area measure on  $\mathbb{D}$  normalized so it has mass 1, and  $\{a_n\}$  are the power series coefficients of  $f$ . It is well known that  $L^2_a(\mathbb{D})$  is a closed subspace of  $L^2(\mathbb{D}, dA)$  and the operator  $f \rightarrow zf$  is continuous. The subspaces  $\mathcal{M} \subset L^2_a(\mathbb{D})$  with  $z\mathcal{M} \subset \mathcal{M}$  (also called *invariant subspaces*) are tremendously complicated and poorly understood. As before, we define the *index* (or *co-dimension*) of an invariant subspace  $\mathcal{M} \subset L^2_a(\mathbb{D})$  to be  $\text{ind } \mathcal{M} = \dim(\mathcal{M}/z\mathcal{M})$ . It is known [2] that given  $n \in \mathbb{N} \cup \{\infty\}$  there exists an invariant subspace  $\mathcal{M} \subset L^2_a(\mathbb{D})$  with  $\text{ind } \mathcal{M} = n$ . Moreover, using sampling theory, Hedenmalm, Richter, and Seip [4], [5] have been able to generate specific examples of these types of invariant subspaces by using zero-based invariant subspaces.

Given a sequence  $A \subset \mathbb{D}$ , we let

$$\mathcal{I}(A) = \{f \in L^2_a(\mathbb{D}) : f|_A = 0\}.$$

In the case where  $\mathcal{I}(A) \neq 0$  one can show that  $\mathcal{I}(A)$  is a closed invariant subspace of  $L_a^2(\mathbb{D})$  with  $\text{ind } \mathcal{I}(A) = 1$ . We now state the Seip, Hedenmalm, Richter result [4], [5].

**Theorem 2.1** (Hedenmalm, Richter, Seip). *Given  $n \in \mathbb{N} \cup \{\infty\}$  there are sequences  $A_j, 0 \leq j < n$ , of points in  $\mathbb{D}$  such that*

$$\mathcal{M} = \text{span}\{\mathcal{I}(A_j) : 0 \leq j < n\}$$

*is an invariant subspace of  $L_a^2(\mathbb{D})$  with  $\text{ind } \mathcal{M} = n$ .*

### 3. THE CORRESPONDENCE

As mentioned in the introduction, there will be a natural correspondence between the invariant subspaces  $D_A \subset \mathcal{F} \subset D$  and the invariant subspaces of the Bergman space. This type of correspondence was observed by Makarov [9] in the  $C^\infty(\mathbb{T})$  case where he observed a natural correspondence between the invariant subspaces  $C_A^\infty(\mathbb{T}) \subset \mathcal{F} \subset C^\infty(\mathbb{T})$  and the invariant subspaces (under  $f \rightarrow zf$ ) of  $A^{-\infty}$ , the analytic distributions. Here  $C_A^\infty(\mathbb{T}) = \{f \in C^\infty : \hat{f}(n) = 0 \forall n < 0\}$ , and  $A^{-\infty}$  are the distributions  $g$  with  $\hat{g}(n) = 0$  for all  $n < 0$ . We make a similar observation for the Dirichlet space and also prove an interesting index formula.

The dual space of  $D$ , which we will denote by  $D'$ , is the space of distributions  $g$  with

$$\sum_{n=-\infty}^{\infty} T_\infty \frac{|\hat{g}(n)|^2}{|n|+1} < \infty.$$

The pairing between  $D$  and  $D'$  is given by

$$\langle f, g \rangle = \sum_{n=-\infty}^{\infty} \hat{f}(n)\overline{\hat{g}(n)}, \quad f \in D, g \in D'.$$

Let  $D'_A$  be the analytic distributions of  $D'$

$$D'_A = \{g \in D' : \hat{g}(n) = \langle \zeta^n, g \rangle = 0 \forall n < 0\}.$$

Making  $\{\hat{g}(n)\}$  the power series coefficients of an analytic function  $g(z)$  on the disk, we see that  $D'_A$  can be naturally identified with the Bergman space  $L_a^2(\mathbb{D})$  and moreover

$$(3.1) \quad \langle f, g \rangle = \lim_{r \rightarrow 1^-} \int_{\mathbb{T}} f(r\zeta)\overline{g(r\zeta)} \frac{|d\zeta|}{2\pi}, \quad f \in D, g \in D'_A.$$

**Theorem 3.1.** (1) *If  $D_A \subset \mathcal{F} \subset D$  is invariant, then  $\mathcal{M} = (\overline{\zeta\mathcal{F}})^\perp$  is an invariant subspace of  $L_a^2(\mathbb{D})$ .*

(2) *If  $\mathcal{M} \subset L_a^2(\mathbb{D})$  is invariant, then  $\mathcal{F} = \overline{\zeta(\mathcal{M})}^\perp$  is invariant and  $D_A \subset \mathcal{F} \subset D$ .*

(3)  $\text{ind } \mathcal{M} = \text{ind } \mathcal{F}$ .

*Proof.* (1) Letting  $g \in \mathcal{M}$ , we see that  $\langle \overline{\zeta f}, g \rangle = 0$  for all  $f \in \mathcal{F}$ . But since  $\zeta^n \in \mathcal{F}$  for all  $n \geq 0$ , then  $\langle \overline{\zeta}^{n+1}, g \rangle = \hat{g}(-n-1) = 0$  for all  $n \geq 0$ , i.e.  $g \in D'_A = L_a^2(\mathbb{D})$ . Also, by (3.1),

$$\langle \overline{\zeta f}, zg \rangle = \langle \overline{\zeta}(\overline{\zeta f}), g \rangle = 0 \quad \forall f \in \mathcal{F}$$

since  $\mathcal{F}$  is invariant. Thus  $zg \in \mathcal{M}$ , which makes  $\mathcal{M}$  an invariant subspace of  $L_a^2(\mathbb{D})$ .

(2) If  $f \in \mathcal{F}$ , then  $\zeta f \in (\overline{\mathcal{M}})^\perp$  and so

$$\langle \zeta(\zeta f), \overline{g} \rangle = \langle \zeta f, \overline{zg} \rangle = 0 \quad \forall g \in \mathcal{M}$$

since  $\mathcal{M}$  is invariant. Thus  $\zeta f \in \overline{\zeta(\overline{\mathcal{M}})}^\perp = \mathcal{F}$ . Also,  $\mathcal{M} \subset L_a^2(\mathbb{D}) = D'_A$  and so

$$0 = \langle \zeta^{n+1}, \overline{g} \rangle = \langle \zeta \zeta^n, \overline{g} \rangle \quad \forall n \geq 0, g \in \mathcal{M},$$

which implies  $\zeta^n \in \overline{\zeta(\overline{\mathcal{M}})}^\perp = \mathcal{F}$ . Thus  $D_A \subset \mathcal{F} \subset D$  and  $\mathcal{F}$  is invariant.

(3) To prove the index formula, first we notice from the Hahn-Banach theorem and basic linear algebra that

$$\begin{aligned} \dim(\mathcal{F}/\zeta\mathcal{F}) &= \dim((\zeta\mathcal{F})^\perp/\mathcal{F}^\perp) \\ &= \dim(\overline{\mathcal{M}}/(\overline{\zeta(\overline{\mathcal{M}})}^\perp)^\perp) \\ &= \dim(\mathcal{M}/(\zeta(\mathcal{M}^\perp))^\perp). \end{aligned}$$

The proof will be finished once we show that

$$(3.2) \quad (\zeta(\mathcal{M}^\perp))^\perp = z\mathcal{M}.$$

To this end, we let  $g \in \mathcal{M}$ ; then  $zg \in \mathcal{M}$  and for all  $f \in \mathcal{M}^\perp$ ,

$$\langle \zeta f, zg \rangle = \langle \overline{\zeta} f, g \rangle = \langle f, g \rangle = 0.$$

Thus  $zg \in (\zeta(\mathcal{M}^\perp))^\perp$ .

For the other direction we observe that

$$D_A \subset \mathcal{F} = \overline{\zeta(\overline{\mathcal{M}})}^\perp$$

and so  $\overline{\zeta D_A} \subset \mathcal{M}^\perp$ . Thus if  $g \in (\zeta(\mathcal{M}^\perp))^\perp$ , then

$$0 = \langle g, \zeta \overline{\zeta} \rangle = \langle g, 1 \rangle = \hat{g}(0).$$

Next we observe (using the invariance of  $\mathcal{M}$ ) that if  $h \in \mathcal{M}^\perp$ ,

$$0 = \langle h, k \rangle = \langle h, zk \rangle = \langle \overline{\zeta} h, k \rangle \quad \forall k \in \mathcal{M}.$$

Thus  $\overline{\zeta} h \in \mathcal{M}^\perp$ . Hence  $0 = \langle g, \zeta \overline{\zeta} h \rangle = \langle g, h \rangle$  for all  $h \in \mathcal{M}^\perp$ . Thus  $g \in (\mathcal{M}^\perp)^\perp = \mathcal{M}$ . Now using the fact that  $g \in L_a^2(\mathbb{D})$  and  $g(0) = 0$ , we get that  $\frac{1}{z}g \in L_a^2(\mathbb{D})$  and so, by (3.1),

$$0 = \langle g, \zeta h \rangle = \left\langle z \frac{1}{z} g, \zeta h \right\rangle = \left\langle \frac{1}{z} g, h \right\rangle \quad \forall h \in \mathcal{M}^\perp.$$

Thus  $\frac{1}{z}g \in \mathcal{M}$  and hence  $g \in z\mathcal{M}$ . So (3.2) has been established and the proof is complete.  $\square$

Recall from (1.1) that for  $w \in \mathbb{D}$ ,  $B_w(\zeta)$  is the single Blaschke factor with zero at  $w$ . Using the fact that  $\mathcal{I}(\{w\}) = \text{span}\{z^n B_w(z) : n \geq 0\}$ , and the F. and M. Riesz theorem, one can prove the following.

**Lemma 3.2.** *If  $\mathcal{M} = \mathcal{I}(\{w\})$ , then  $\mathcal{F} = \overline{\zeta(\overline{\mathcal{M}})}^\perp = \overline{B_w} H^2 \cap D$ .*

We are now ready to prove our main theorem.

*Proof of Theorem 1.1.* By Theorem 2.1, the invariant subspace

$$\mathcal{M} = \text{span}\{\mathcal{I}(A_j) : 0 \leq j < n\}$$

has index  $n$ . Thus by Theorem 3.1,  $\mathcal{F} = \overline{\zeta}(\overline{\mathcal{M}})^\perp$  also has index  $n$ . We now identify  $\mathcal{F}$ :

$$\begin{aligned} \mathcal{F} &= \overline{\zeta}(\overline{\mathcal{M}})^\perp \\ &= \overline{\zeta} \left( \text{span}\{\overline{\mathcal{I}(A_j)} : 0 \leq j < n\} \right)^\perp \\ &= \overline{\zeta} \left( \bigcap_{0 \leq j < n} \overline{\mathcal{I}(A_j)}^\perp \right) \\ &= \bigcap_{0 \leq j < n} \overline{\zeta} \left( \bigcap_{a \in A_j} \overline{\mathcal{I}(\{a\})} \right)^\perp \quad \text{since mult. by } \overline{\zeta} \text{ is invertible} \\ &= \bigcap_{0 \leq j < n} \overline{\zeta} \text{ span}\{\overline{\mathcal{I}(\{a\})}^\perp : a \in A_j\} \\ &= \bigcap_{0 \leq j < n} \text{span}\{\overline{\zeta}(\overline{\mathcal{I}(\{a\})}^\perp) : a \in A_j\} \quad \text{since mult. by } \zeta \text{ is cont. and invertible} \\ &= \bigcap_{0 \leq j < n} \text{span}\{\overline{B_a}H^2 \cap D : a \in A_j\} \quad \text{by Lemma 3.2.} \quad \square \end{aligned}$$

#### 4. GENERALIZATIONS

In this last section, we remark that the analog of Theorem 1.1 is also true for the Dirichlet classes  $D_\alpha$ ,  $0 < \alpha < \infty$ , of  $f \in L^2(\mathbb{T}, |d\zeta|)$  with

$$\sum_{n=-\infty}^{\infty} (1 + |n|)^\alpha |\hat{f}(n)|^2 < \infty,$$

and the Besov classes  $B_p^\alpha$ ,  $1 < p < \infty$ ,  $0 < \alpha < 1$ , of  $f$  with

$$\int_0^\pi \int_0^{2\pi} \frac{|f(e^{i(\theta+h)}) - f(e^{i\theta})|^p}{h^{1+\alpha p}} d\theta dh < \infty.$$

Here the appropriate Bergman spaces to consider are the weighted Bergman spaces  $A_p^\alpha$ ,  $0 < \alpha < \infty$ ,  $1 < p < \infty$ , of analytic functions on  $\mathbb{D}$  with

$$\int_{\mathbb{D}} |f(z)|^p (1 - |z|)^{\alpha p - 1} dA(z) < \infty.$$

The analog of Theorem 2.1 is true for  $A_p^\alpha$  where, of course,  $\mathcal{I}(A) = \{f \in A_p^\alpha : f|_A = 0\}$  [4], [5]. For  $D_\alpha$ , the analytic distributions (in the dual) can be identified with the weighted Bergman space  $A_2^{\alpha/2}$  via (3.1) [7]. For  $B_p^\alpha$ , the analytic distributions (in the dual) can be identified with  $A_q^\alpha$ , where  $q = p(p - 1)^{-1}$  via (3.1). For both these spaces, the analog of Theorem 3.1 remains true. Thus we have the following.

**Theorem 4.1.** *Let  $X = D_\alpha$ ,  $0 < \alpha < \infty$ , or  $B_p^\alpha$ ,  $1 < p < \infty$ ,  $0 < \alpha < 1$ , and  $n \in \mathbb{N} \cup \{\infty\}$ . Then there are sequences  $A_j$ ,  $0 \leq j < n$ , of points in  $\mathbb{D}$  so that*

$$\mathcal{F} = \bigcap_{0 \leq j < n} \text{span}_X\{\overline{B_a}H^2 \cap X : a \in A_j\}$$

*is a simply invariant subspace of  $X$  with  $\text{ind } \mathcal{F} = n$ .*

## REFERENCES

- [1] A. Aleman, S. Richter, and W.T. Ross, 'Bergman spaces on disconnected domains, *Canad. J. Math.* (to appear).
- [2] H. Bercovici, C. Foiaş, and C. Pearcy, *Dual algebras with applications to invariant subspaces and dilation theory*, CBMS Regional Conf. Ser. in Math., no. 56, Amer. Math. Soc., Providence, RI., 1985. MR **87g**:47091
- [3] M. Hasumi and T.P. Srinivasan, 'Invariant subspaces of continuous functions', *Canad. J. Math.* **22** (1965), 643 - 651. MR **31**:3871
- [4] H. Hedenmalm, 'An invariant subspace of the Bergman space having the co-dimension two property', *J. Reine Angew. Math.* **443** (1993), 1 - 9. MR **94k**:30092
- [5] H. Hedenmalm, S. Richter, K. Seip, 'Zero sequences and invariant subspaces in the Bergman space', preprint.
- [6] H. Helson, *Lectures on invariant subspaces*, New York and London, 1964. MR **30**:1409
- [7] S. Khrushchev and V. Peller, 'Hankel operators, best approximation, and stationary Gaussian processes', *Russian Math. Surveys*, **37** (1982), 61 - 144. MR **84e**:47036
- [8] N.G. Makarov, 'Sets of 1-invariant and  $\lambda$ -invariant subspaces (smooth functions)', *Soviet Math. Dokl.* **25** (1982), 191 - 194. MR **83h**:46053
- [9] N.G. Makarov, 'Invariant subspaces of the space  $C^\infty(\mathbb{T})$ ', *Math. USSR Sbornik*, **47** (1984), 1 - 26. MR **84d**:30088
- [10] B. Malgrange, *Ideals of differentiable functions*, Tata Inst. Fund. Res., Bombay and Oxford Univ. Press, London, 1967. MR **35**:3446
- [11] Y.U. Netrusov 'Spectral synthesis in spaces of smooth functions', *Russian Acad. Dokl. Math.*, **46** (1993), 135 - 138. MR **94c**:46068
- [12] S. Richter, W. T. Ross, and C. Sundberg, 'Hyperinvariant subspaces of the harmonic Dirichlet space', *J. Reine Angew. Math.* **448** (1994), 1 - 26. MR **95e**:47045
- [13] S. Richter and A. Shields, 'Bounded analytic functions in the Dirichlet space', *Math. Z.* **198** (1988), 151 - 159. MR **89c**:46039
- [14] K. Seip, 'Beurling type density theorems in the unit disk', *Invent. Math.* **113** (1993), 26 - 39. MR **94g**:30033

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF RICHMOND, RICHMOND, VIRGINIA 23173  
E-mail address: rossb@mathcs.urich.edu