

**THE INDICES, THE NULLITIES AND THE STABILITY OF
TOTALLY GEODESIC SUBMANIFOLDS IN THE COMPLEX
QUADRATIC HYPERSURFACES: $Q_m = SO(m+2)/SO(m) \times SO(2)$**

ZHAO QIANG

(Communicated by Roe W. Goodman)

ABSTRACT. In the paper, the stability of totally geodesic submanifolds in the complex quadratic hypersurfaces: $Q_m = SO(m+2)/SO(m) \times SO(2)$ ($m > 1$) is discussed, and the indices, the nullities and the Killing nullities of totally geodesic submanifolds in Q_m are calculated.

1. INTRODUCTION

It is always an interesting and important problem to find all stable minimal submanifolds in each symmetric space. In 1980, B.Y.Chen, P.F.Leung and T.Nagano gave the algorithm of determining the stability of totally geodesic submanifolds in compact symmetric spaces which was reformulated in 1987 by Y.Ohnita (cf. [1] and [2]). In [3], B.Y.Cheng and T.Nagano completely classified complete, connected, totally geodesic submanifolds of the complex quadratic hypersurface: $Q_m = SO(m+2)/SO(2) \times SO(m)$, $m > 1$. Using their results, we determine the indices and nullities of all totally geodesic submanifolds in Q_m , and discuss their stability.

Suppose M is a compact minimal submanifold of a Riemannian manifold N with metric h . The isometric immersion $\phi : M \rightarrow N$ is called **stable** if the second derivative of the volume $Vol(M, \phi_t^* h)$ at $t = 0$ is nonnegative for every smooth variation $\{\phi_t\}$ of ϕ with $\phi_0 = \phi$. Then we say that M is a **stable minimal submanifold of N** . Choose a smooth variation $\{\phi_t\}$ of ϕ with $\phi_0 = \phi$ and $(\partial/\partial t)\phi_t(x)|_{t=0} = V_x(x \in M)$ for any vector field $V \in \Gamma(\phi^{-1}T(N))$. Then the classical second variational formula is given as follows:

$$(d^2/dt^2)Vol(M, \phi_t^* h)|_{t=0} = \int_M \langle \mathcal{J}(V^N), V^N \rangle dv$$

where dv denotes the Riemannian measure of (M, g) and V^N the component of V normal to M . \mathcal{J} is a self-adjoint strongly elliptic linear differential operator of order 2 acting on the space $\Gamma(N(M))$ of smooth sections of the normal bundle $N(M)$, called **the Jacobi operator** of ϕ . \mathcal{J} has discrete eigenvalues $\mu_1 < \mu_2 < \dots \rightarrow \infty$. Set

$$E_\mu = \{V \in \Gamma(N(M)) \mid \mathcal{J}(V) = \mu V\}.$$

Received by the editors August 5, 1994 and, in revised form, January 18, 1995.
1991 *Mathematics Subject Classification*. Primary 53C35; Secondary 22E70.
Key words and phrases. Indices, nullities, stability.

We call the number $\sum_{\mu < 0} \dim E_\mu$ **the index** of ϕ or the index of M in N ; denote it by $i(\phi)$ or $i(M)$. It is easy to see that ϕ is stable if and only if $i(\phi) = 0$. The number $\dim E_0$ is called **the nullity** of ϕ and denoted by $n(\phi)$ or $n(M)$. Define

$$P = \{X^N \mid X \text{ is a Killing vector field on } N\} \subset \Gamma(N(M)).$$

Then $P \subset E_0$. The $\dim P$ is called **the Killing nullity** of ϕ and denoted by $n_k(\phi)$ or $n_k(M)$.

In particular, if M is an m -dimensional compact totally geodesic submanifold immersed in a compact Riemannian symmetric space N with metric g_N , the immersion $\phi: M \rightarrow N$ can be expressed as follows: There are compact symmetric pairs (U, L) and (G, K) with $N = U/L$, $M = G/K$ and

$$\begin{aligned} \phi: M = G/K &\rightarrow N = U/L, \\ gK &\rightarrow \rho(g)L \end{aligned}$$

where $\rho: G \rightarrow U$ is an analytic homomorphism with $\rho(K) \subset L$ and the injective differential $\rho: g \rightarrow u$ satisfying $\rho(m) \subset p$. Here $u = l + p$ and $g = k + m$ are the Cartan decompositions of u and g , respectively. There exists an $\text{ad}U$ -invariant inner product (\cdot, \cdot) on u such that (\cdot, \cdot) induces the metric g_N on N . By (\cdot, \cdot) we also denote the $\text{ad}G$ -invariant inner product on g induced from (\cdot, \cdot) through ρ . Let m^\perp be the smooth orthogonal complement of $\rho(m)$ with p relative to (\cdot, \cdot) , and k^\perp the orthogonal complement of $\rho(k)$ in l . Put $g^\perp = k^\perp + m^\perp$. Then g^\perp is the orthogonal complement of $\rho(g)$ in u relative to (\cdot, \cdot) , and g^\perp is $\text{ad}\rho(G)$ -invariant. Let θ be the involutive automorphism of the symmetric pair (U, L) . Choose an orthogonal decomposition $g^\perp = g_1^\perp \oplus \cdots \oplus g_t^\perp$ such that each g_i^\perp is an irreducible $\text{ad}\rho(G)$ -invariant subspace with $\theta(g_i^\perp) = g_i^\perp$. Then the Casimir operator C of the representation of G on each g_i^\perp is $a_i I$ for $a_i \in \mathbf{C}$. Put $g_i^\perp = k_i^\perp + m_i^\perp$, where $k_i^\perp = k^\perp \cap g_i^\perp$ and $m_i^\perp = m^\perp \cap g_i^\perp$.

Theorem 1.1 ([1]). *The index, nullity and Killing nullity of ϕ are given as follows:*

$$\begin{aligned} (1) \quad i(M) &= \sum_{i=1}^t \sum_{\lambda \in D(G), a_\lambda > a_i} \dim \text{Hom}_K(V(\lambda), (m_i^\perp)^\mathbf{C}) \dim V(\lambda); \\ (2) \quad n(M) &= \sum_{i=1}^t \sum_{\lambda \in D(G), a_\lambda = a_i} \dim \text{Hom}_K(V(\lambda), (m_i^\perp)^\mathbf{C}) \dim V(\lambda); \\ (3) \quad n_K(M) &= \sum_{i=1, m_i^\perp \neq 0}^t \dim g_i^\perp, \end{aligned}$$

where $D(G)$ is the set of irreducible representations of G and a_λ is the eigenvalue for the Casimir operator of the irreducible G -module (λ, V_λ) relative to (\cdot, \cdot) .

2. THE SPACE Q_m AND ITS TOTALLY GEODESIC SUBMANIFOLDS

The main result of [3] is the following theorem. It gave the classification of all complete, connected, totally geodesic submanifolds of the complex quadratic hypersurface: $Q_m = SO(2+m)/SO(2) \times SO(m)$, $m > 1$.

Theorem 2.1 ([3]). *If M is a maximal totally geodesic submanifold of Q_m , M is one of the following three spaces:*

- (1) Q_{m-1} ;

- (2) a local Riemannian product of two spheres S^p and S^q , $p + q = m$;
- (3) the complex projective space $P(\mathbf{C}^{n+1})$ of complex dimension n , $2n = m$.

If M is a nonmaximal, totally geodesic submanifold of Q_m , M is either contained in Q_{m-1} in an appropriate position in Q_m , or the real projective space $P(\mathbf{R}^{n+1})$ of real dimension n , $2n = m$, which is the intersection of $P(\mathbf{C}^{n+1})$ in (3) and the local product space in (2) with $p = q = n$.

Now we have Riemannian symmetric spaces

$$N = Q_m = U/L, \quad U = SO(m + 2), \quad L = SO(2) \times SO(m).$$

Then $u = l + p$ is the corresponding Cartan decomposition, where

$$l = \left\{ \left(\begin{array}{cc} A & 0 \\ 0 & B \end{array} \right) \mid A = \begin{pmatrix} 0 & a \\ -a & 0 \end{pmatrix}, a \in \mathbf{R}, B \in gl(m), B^t = -B \right\},$$

$$p = \left\{ \left(\begin{array}{cc} 0 & A \\ -A^t & 0 \end{array} \right) \mid A \text{ is a matrix of } 2 \text{ rows and } m \text{ columns over } \mathbf{R} \right\}.$$

From now on, we denote the set of matrices of m rows and n columns with elements in \mathbf{R} by $\mathbf{R}(m, n)$.

If M is a compact totally geodesic submanifold immersed in N , the immersion $\phi : M \rightarrow N$ can be expressed as follows: There exist a compact symmetric pair (G, K) with $M = G/K$ and a homomorphism $\rho : G \rightarrow U$ such that ϕ has form as $gK \rightarrow \rho(g)L$. Then $\rho(K) \subset L$ and the injective differential $\rho : g \rightarrow u$ satisfies $\rho(m) \subset p$. We choose an adU -invariant inner product on u as $(X, Y) = TrXY$ ($X, Y \in u$); it induces the metric on Q_m . Let $g = k + m$ be the corresponding Cartan decomposition of g . Below we list the corresponding ρ , k and m in every case.

1. $M = Q_n$ ($n < m$), $\rho : g \rightarrow u$ is

$$A \longrightarrow \begin{pmatrix} A & \\ & 0 \end{pmatrix} \in u, \quad A \in g,$$

$$k = \left\{ \begin{pmatrix} A & & \\ & B & \\ & & 0 \end{pmatrix} \mid A \in gl(2), B \in gl(n), A^t = -A, B^t = -B \right\},$$

$$m = \left\{ \begin{pmatrix} 0 & A & 0 \\ -A^t & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \mid A \in \mathbf{R}(2, n) \right\}.$$

2. $M = S^p \times S^q$ ($p + q = m$). $\rho : g \rightarrow u$ is

$$(A, B) \longrightarrow \begin{pmatrix} b_{11} & 0 & b_{12} & b_{13} & \cdots & b_{1,q+1} \\ 0 & A & 0 & 0 & \cdots & o \\ b_{21} & 0 & b_{22} & b_{23} & \cdots & b_{2,q+1} \\ b_{31} & 0 & b_{32} & b_{33} & \cdots & b_{3,q+1} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ b_{q+1,1} & 0 & b_{q+1,2} & b_{q+1,3} & \cdots & b_{q+1,q+1} \end{pmatrix}.$$

Here $A = (a_{ij}) \in gl(p+1)$, $B = (b_{ij}) \in gl(q+1)$, $A^t = -A$, $B^t = -B$.

$$k = \left\{ \left(\begin{array}{cc} 0 & \\ & A & \\ & & B \end{array} \right) \mid A \in gl(p), B \in gl(q), A^t = -A, B^t = -B \right\},$$

$$m = \left\{ \left(\begin{array}{cccc} 0 & 0 & 0 & X \\ 0 & 0 & Y & 0 \\ 0 & -Y^t & 0 & 0 \\ -X^t & 0 & 0 & 0 \end{array} \right) \mid X \in \mathbf{R}(1, q), Y \in \mathbf{R}(1, p) \right\}.$$

3. $M = \mathbf{C}P^n = SU(n+1)/SU(n)$ ($m = 2n$). $\rho : su(n+1) \rightarrow so(2n+2)$ is

$$A + iB \longrightarrow P_{n+1}(A, B).$$

Here $P_{n+1}(A, B)$ is

$$\left(\begin{array}{cccccc} a_{11} & b_{11} & a_{12} & b_{12} & \cdots & a_{1,n+1} & b_{1,n+1} \\ -b_{11} & a_{11} & -b_{12} & a_{12} & \cdots & -b_{1,n+1} & a_{1,n+1} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{n+1,1} & b_{n+1,1} & a_{n+1,2} & b_{n+1,2} & \cdots & a_{n+1,n+1} & b_{n+1,n+1} \\ -b_{n+1,1} & a_{n+1,1} & -b_{n+1,2} & a_{n+1,2} & \cdots & -b_{n+1,n+1} & a_{n+1,n+1} \end{array} \right)$$

where $A = (a_{ij})$, $B = (b_{ij}) \in gl(n+1)$, $A^t = -A$, $B^t = -B$, $TrB = 0$.

$$k = \left\{ \left(\begin{array}{cc} 0 & 0 \\ 0 & P_n(B, C) \end{array} \right) \mid B^t = -B, C^t = C, TrC = 0 \right\},$$

$$m = \left\{ \left(\begin{array}{cc} 0 & O(A, B) \\ -O(A, B)^t & 0 \end{array} \right) \mid O(A, B) \in \mathbf{R}(2, 2n) \right\}$$

where

$$O(A, B) = \left(\begin{array}{cccc} a_1 & b_1 & \cdots & a_n & b_n \\ -b_1 & a_1 & \cdots & -b_n & a_n \end{array} \right).$$

4. $M = \mathbf{R}P^n = SO(n+1)/SO(n)$ ($m = 2n$). $\rho : so(n+1) \rightarrow so(2n+2)$ is $A \rightarrow P_{n+1}(A)$, and $P_{n+1}(A)$ equals

$$\left(\begin{array}{cccccc} a_{11} & 0 & a_{12} & 0 & \cdots & a_{1,n+1} & 0 \\ 0 & a_{11} & 0 & a_{12} & \cdots & 0 & a_{1,n+1} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{n+1,1} & 0 & a_{n+1,2} & 0 & \cdots & a_{n+1,n+1} & 0 \\ 0 & a_{n+1,1} & 0 & a_{n+1,2} & \cdots & 0 & a_{n+1,n+1} \end{array} \right)$$

where $A = (a_{ij}) \in so(n+1)$.

$$k = \left\{ \left(\begin{array}{cc} 0 & 0 \\ 0 & P_n(A) \end{array} \right) \mid A \in so(n) \right\},$$

$$m = \left\{ \left(\begin{array}{cc} 0 & T(A) \\ -T(A)^t & 0 \end{array} \right) \mid T(A) \in \mathbf{R}(2, 2n) \right\},$$

$$T(A) = \left(\begin{array}{cccc} a_1 & 0 & a_2 & 0 & \cdots & a_n & 0 \\ 0 & a_1 & 0 & a_2 & \cdots & 0 & a_n \end{array} \right).$$

3. THE INDICES, NULLITIES AND STABILITY OF THE GEODESIC SUBMANIFOLDS IN Q_m

We choose the inner product $(X, Y) = TrXY$ on $u = so(m + 2)$ which induces the metric on Q_m . In every case, we first calculate g^\perp , m^\perp and the orthogonal decomposition of the G -module $(g^\perp)^\mathbb{C}$.

1. $M = Q_n$ ($n < m$),

$$g^\perp = \left\{ \begin{pmatrix} 0 & B \\ -B^t & C \end{pmatrix} \mid B \in \mathbf{R}(n + 2, m - n), C \in gl(m - n), C^t = -C \right\}$$

$$= \left\{ \begin{pmatrix} 0 & 0 \\ 0 & C \end{pmatrix} \right\} \oplus \left\{ \begin{pmatrix} 0 & B \\ -B^t & 0 \end{pmatrix} \right\} = g_1^\perp \oplus g_2^\perp,$$

$$m_1^\perp = 0,$$

$$m_2^\perp = \left\{ \begin{pmatrix} 0 & 0 & D \\ 0 & 0 & 0 \\ -D^t & 0 & 0 \end{pmatrix} \mid D \in \mathbf{R}(2, m - n) \right\}.$$

Obviously the representation of G on g_1^\perp is trivial. Below we investigate the G -module $(g_2^\perp)^\mathbb{C}$. Let E_{ij} be the square matrix with entry 1 where the i -th row and j -th column meet, all other entries being 0.

1) If $n + 2 = 2s, m + 2 = 2(s + t)$, set $H_i = E_{2i-1,2i} - E_{2i,2i-1}$ ($1 \leq i \leq s$); then $h = span_{\mathbb{C}}\{H_i\}_1^s$ is a Cartan subalgebra of $g^\mathbb{C}$. Let $F_{ij} = E_{ij} - E_{ji}$ and

$$G_{jk}^\pm = F_{2j-1,2k-1} \pm F_{2j,2k} + i(F_{2j-1,2k} \mp F_{2j,2k-1}) \quad (j \neq k).$$

It is obvious that

$$(g_2^\perp)^\mathbb{C} = span_{\mathbb{C}}\{G_{jk}^\pm \mid 1 \leq j \leq s < k \leq s + t \text{ or } 1 \leq k \leq s < j \leq s + t\},$$

$$(adH)G_{jk}^+ = (e_j(H) - e_k(H))G_{jk}^+,$$

$$(adH)G_{jk}^- = sgn(j - k)(e_j(H) + e_k(H))G_{jk}^-$$

for $H \in h$, where $e_j(H_k) = -i\delta_{jk}$. So the set of weights of the G -module $(g_2^\perp)^\mathbb{C}$ is

$$\Phi = \underbrace{\{\pm e_j, \dots, \pm e_j\}}_{2t} \mid 1 \leq j \leq s.$$

We can choose a set of simple roots of $(g)^\mathbb{C}$ as

$$R = \{\alpha_j = e_j - e_{j+1}, \alpha_s = e_{s-1} + e_s \mid 1 \leq j \leq s\}.$$

In this orientation, all the dominant weights which are also highest weights are $\underbrace{\{e_1, \dots, e_1\}}_{2t}$. But the representation having the highest weight e_1 is exactly the

first basic representation ω_1 of $so(n + 2)$, so

$$(g_2^\perp)^\mathbb{C} = \underbrace{\omega_1 \oplus \dots \oplus \omega_1}_{2t}.$$

Generally, for the representation having the highest weight $\lambda = \sum_i a_i \omega_i$ of g , the Casimir operator C has the form $C = a_\lambda I$, where

$$a_\lambda = -(a_i \omega_i + 2\rho, a_j \omega_j) = -(a_i a_j g_{ij} + 2 \sum_{j,k} a_j g_{jk}).$$

Here ρ is half the sum of positive roots, and the information about g_{ij} may be found in [4].

Therefore, the Casimir operator C of g on $(g_i^\perp)^\mathbf{C}$ has the action $a_i I$ ($i = 1, 2$); here $a_1 = 0$ and $a_2 = -(2s - 1)$. So

$$\begin{aligned} \{\lambda \in D(G) \mid a_\lambda > a_2\} &= \{0\}, \\ \{\lambda \in D(G) \mid a_\lambda = a_2\} &= \{\omega_1\}. \end{aligned}$$

It is easy to see that $k^\mathbf{C} = \mathbf{C}H_1 \oplus so(n, \mathbf{C})$. Now $m_1^\perp = 0$, and the set of weights of the $\mathbf{C}H_1$ -module $(m_2^\perp)^\mathbf{C}$ is

$$\underbrace{\{1, \dots, 1\}}_{2t}, \underbrace{\{-1, \dots, -1\}}_{2t}.$$

Since the action of $so(n, \mathbf{C})$ on $(m_2^\perp)^\mathbf{C}$ is trivial, as a K -module,

$$(m_2^\perp)^\mathbf{C} = \bigoplus_{2t} (\lambda(1, 0, \dots, 0) \oplus \lambda(-1, 0, \dots, 0)).$$

By the branching rule of representations [5], $\dim \omega_1 = 2s = n + 2$,

$$\omega_1 = \begin{cases} \lambda(1, 1, 0, \dots, 0) \oplus \lambda(1, 0, \dots, 0) \oplus \lambda(-1, 0, \dots, 0), & s > 3 \\ \lambda(1, 0, 1, 0, \dots, 0) \oplus \lambda(1, 0, \dots, 0) \oplus \lambda(-1, 0, \dots, 0), & s = 3. \end{cases}$$

So we have

- (1) $i(M) = 0,$
- (2) $n(M) = (n + 2)(m - n),$
- (3) $n_K(M) = (n + 2)(m - n).$

2) If $n + 2 = 2s, m + 2 = 2(s + t) + 1$, let

$$D_j^\pm = F_{2j-1, m+2} \pm iF_{2j, m+2}, \quad 1 \leq j \leq s.$$

Then

$$(adH)D_j^\pm = \pm e_j(H)D_j^\pm, \quad 1 \leq j \leq s, \quad H \in h.$$

The set of dominant weights which are also the highest weights of the g -module $(g_2^\perp)^\mathbf{C}$ is

$$\underbrace{\{\omega_1, \dots, \omega_1\}}_{2t+1}.$$

Using methods similar to 1) we get

- (4) $i(M) = 0,$
- (5) $n(M) = 2s(2t + 1) = (n + 2)(m - n),$
- (6) $n_K(M) = (n + 2)(m - n).$

For other cases of $m + 2$ and $n + 2$, we can get the same results by similar discussion.

2. $\phi : M = S^p \times S^q \longrightarrow Q_m (p+q = m)$. By a series of similarity transformations, we have

$$\begin{aligned}
 g &= \left\{ \begin{pmatrix} A & & & \\ & B & & \\ & & & \\ & & & \end{pmatrix} \mid A \in so(p+1), B \in so(q+1) \right\}, \\
 k &= \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & A & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & B \end{pmatrix} \mid A \in so(p), B \in so(q) \right\}, \\
 m &= \left\{ \begin{pmatrix} 0 & X & 0 & 0 \\ -X^t & 0 & 0 & 0 \\ 0 & 0 & 0 & Y \\ 0 & 0 & -Y^t & 0 \end{pmatrix} \mid X \in \mathbf{R}(1,p), Y \in \mathbf{R}(1,q) \right\}, \\
 g^\perp &= \left\{ \begin{pmatrix} 0 & B \\ -B^t & 0 \end{pmatrix} \mid B \in \mathbf{R}(p+1, q+1) \right\}, \\
 m^\perp &= \left\{ \begin{pmatrix} 0 & 0 & 0 & Y \\ 0 & 0 & X & 0 \\ 0 & -X^t & 0 & 0 \\ -Y^t & 0 & 0 & 0 \end{pmatrix} \mid X \in \mathbf{R}(1,p), Y \in \mathbf{R}(1,q) \right\}.
 \end{aligned}$$

If $p + 1 = 2s, q + 1 = 2t$, choose $H_i = E_{2i-1,2i} - E_{2i,2i-1} (1 \leq i \leq s + t)$; then $h = span\{H_i\}_1^{s+t}$ is a Cartan subalgebra of $g^{\mathbf{C}} = (so(2s) \oplus so(2t))^{\mathbf{C}}$. By a similar discussion as before, we know $(g^\perp)^{\mathbf{C}}$ is an irreducible g -module. It has the highest weight $\lambda(1, 0, \dots, 0 - 1, 0, \dots, 0)$, so it is the tensor product of the first basic representations of $so(2s)$ and $so(2t)$. And

$$\begin{aligned}
 a &= -(\lambda(1, 0, \dots, 0 - 1, 0, \dots, 0) + 2\rho, \lambda(1, 0, \dots, 0 - 1, 0, \dots, 0)) \\
 &= -(\lambda_1(\underbrace{1, 0, \dots, 0}_{s-1}) + 2\rho_1, \lambda_1(1, 0, \dots, 0)) \\
 &\quad -(\lambda_2(\underbrace{1, 0, \dots, 0}_{t-1}) + 2\rho_2, \lambda_2(1, 0, \dots, 0)) \\
 &= -2(s + t - 1), \\
 \{\lambda \in D(G) \mid a_\lambda > a\} &= \{0, \lambda(1, 0, \dots, 0 - 0, \dots, 0), \lambda(0, \dots, 0 - 1, 0, \dots, 0)\}, \\
 \{\lambda \in D(G) \mid a_\lambda = a\} &= \{\lambda(1, 0, \dots, 0 - 1, 0, \dots, 0)\}.
 \end{aligned}$$

Considering $k = so(p) \oplus so(q)$, we have

$$(m^\perp)^{\mathbf{C}} = (m_1^\perp)^{\mathbf{C}} \oplus (m_2^\perp)^{\mathbf{C}}$$

as a K -module, where

$$\begin{aligned}
 m_1^\perp &= \left\{ \begin{pmatrix} & & & 0 \\ & & X & \\ & -X^t & & \\ 0 & & & \end{pmatrix} \mid X \in \mathbf{R}(1,p) \right\}, \\
 m_2^\perp &= \left\{ \begin{pmatrix} & & & Y \\ & & 0 & \\ & & & \\ -Y^t & 0 & & \end{pmatrix} \mid Y \in \mathbf{R}(1,q) \right\}.
 \end{aligned}$$

It is obvious that the actions of $so(p)$ on m_1^\perp and $so(q)$ on m_2^\perp are their standard representations. Thus, as a K -module,

$$(m^\perp)^\mathbb{C} = \lambda'(1, 0, \dots, 0 - 0, \dots, 0) \oplus \lambda'(0, \dots, 0 - 1, 0, \dots, 0).$$

But

$$\begin{aligned} \lambda(1, 0, \dots, 0 - 0, \dots, 0) &= \lambda'(0, \dots, 0 - 0, \dots, 0) \oplus \lambda'(1, 0, \dots, 0 - 0, \dots, 0), \\ \dim \lambda(1, 0, \dots, 0 - 0, \dots, 0) &= p + 1, \dim \lambda(0, \dots, 0 - 1, 0, \dots, 0) = q + 1, \\ \lambda(1, 0, \dots, 0 - 1, 0, \dots, 0) &= \lambda'(1, \dots, 0 - 1, 0, \dots, 0) \\ &\oplus \lambda'(1, 0, \dots, 0 - 0, \dots, 0) \oplus \lambda'(0, \dots, 0 - 1, 0, \dots, 0) \oplus \lambda'(0, \dots, 0 - 0, \dots, 0), \\ \dim \lambda(1, 0, \dots, 0 - 1, 0, \dots, 0) &= (p + 1)(q + 1). \end{aligned}$$

So we have

- (7) $i(M) = m + 2,$
- (8) $n(M) = 2(p + 1)(q + 1),$
- (9) $n_K(M) = (p + 1)(q + 1).$

We have the same results for the other cases of p and q by similar discussion.

3. $\phi: \mathbf{C}P^n = SU(n + 1)/SU(n) \rightarrow Q_{2n} (m = 2n),$

$$\begin{aligned} g^\perp &= \{Q(A, B) | A, B \in so(n + 1)\}, \\ m^\perp &= \left\{ \begin{pmatrix} 0 & s(a, b) \\ -s(a, b)^t & 0 \end{pmatrix} \mid s(a, b) \in \mathbf{R}(2, 2n) \right\}. \end{aligned}$$

Here, for $A = (a_{ij}), B = (b_{ij}), Q(A, B)$ is

$$\begin{pmatrix} a_{11} & b_{11} & a_{12} & b_{12} & \cdots & a_{1,n+1} & b_{1,n+1} \\ b_{11} & -a_{11} & b_{12} & -a_{12} & \cdots & b_{1,n+1} & -a_{1,n+1} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{n+1,1} & b_{n+1,1} & a_{n+1,2} & b_{n+1,2} & \cdots & a_{n+1,n+1} & b_{n+1,n+1} \\ b_{n+1,1} & -a_{n+1,1} & b_{n+1,2} & -a_{n+1,2} & \cdots & b_{n+1,n+1} & -a_{n+1,n+1} \end{pmatrix},$$

$$s(a, b) = \begin{pmatrix} a_1 & b_1 & \cdots & a_n & b_n \\ b_1 & -a_1 & \cdots & b_n & -a_n \end{pmatrix}.$$

A Cartan subalgebra of g can be imbedded in u as

$$h = \left\{ \sum_{i=1}^{n+1} c_i H_i \mid \sum_i c_i = 0 \right\}.$$

The set of the weights of the g -module $(g^\perp)^\mathbb{C} = \sum_{j \neq k} \mathbf{C}G_{jk}^-$ is

$$\{\pm(e_j + e_k) \mid j \neq k\}.$$

Choose $\{e_1 - e_2, e_2 - e_3, \dots, e_n - e_{n+1}\}$ as the set of simple roots of $g^\mathbb{C}$. The dominant weights of $(g^\perp)^\mathbb{C}$ are $e_1 + e_2$ and $-e_n - e_{n+1}$. By comparing the dimensions among $(g^\perp)^\mathbb{C}$ and the spaces of the representations having the highest weights $\{e_1 + e_2, -e_n - e_{n+1}\}$, we know

$$(g^\perp)^\mathbb{C} = \lambda(0, 1, 0, \dots, 0) \oplus \lambda(0, \dots, 0, 1, 0).$$

Since $(m^\perp)^\mathbb{C} = \sum_{k>1} (\mathbf{C}G_{1k}^- \oplus \mathbf{C}G_{k1}^-)$ and $\{\sum_{i=2}^{n+1} c_i H_i \mid \sum_i c_i = 0\}$ is a Cartan subalgebra of $k = su(n)$, the set of the weights of $(m^\perp)^\mathbb{C}$ is $\{\pm e_k\}_{k=2}^{n+1}$. If we choose

$\{e_i - e_{i-1} | 2 \leq i \leq n\}$ as the set of simple roots of k , the dominant weights of the k -module $(m^\perp)^{\mathbf{C}}$ are e_2 and $-e_{n+1}$. So

$$\begin{aligned} (m^\perp)^{\mathbf{C}} &= \lambda'(1, 0, \dots, 0) \oplus \lambda'(0, \dots, 0, 1), \\ a &= -(\lambda(0, 1, 0, \dots, 0) + 2\rho, \lambda(0, 1, 0, \dots, 0)) \\ &= -(\lambda(0, \dots, 0, 1, 0) + 2\rho, \lambda(0, \dots, 0, 1, 0)) \\ &= -2(n-1)(n+2)/(n+1), \\ a_{\lambda_k} &= -k(n-k+1)(n+2)/(n+1) \text{ for } \lambda_k = \underbrace{\lambda(0, \dots, 0, 1, 0, \dots, 0)}_k; \\ \{\lambda \in D(G) | a_\lambda > a\} &= \{0, \lambda_1, \lambda_n\}, \\ \{\lambda \in D(G) | a_\lambda = a\} &= \{\lambda_2, \lambda_{n-1}\}. \end{aligned}$$

By the branching rule of representations, we know

$$\begin{aligned} \lambda_1 &= \lambda'(1, 0, \dots, 0) \oplus \lambda'(0, \dots, 0), \quad \lambda_n = \lambda'(0, \dots, 0, 1) \oplus \lambda'(0, \dots, 0), \\ \lambda_2 &= \lambda'(0, 1, 0, \dots, 0) \oplus \lambda'(1, 0, \dots, 0), \\ \lambda_{n-1} &= \lambda'(0, \dots, 0, 1) \oplus \lambda'(0, \dots, 0, 1, 0), \\ \dim \lambda_1 &= \dim \lambda_n = n + 1, \quad \dim \lambda_2 = \dim \lambda_{n-1} = n(n + 1)/2. \end{aligned}$$

Therefore we get

$$\begin{aligned} (10) \quad & i(M) = 2(n + 1), \\ (11) \quad & n(M) = n(n + 1), \\ (12) \quad & n_K(M) = n(n + 1). \end{aligned}$$

4. $\mathbf{R}P^n = SO(n + 1)/SO(n) \rightarrow Q_{2n} (m = 2n)$. Through a series of similarity transformations, we have

$$\begin{aligned} g &= \left\{ \begin{pmatrix} A & \\ & A \end{pmatrix} \mid A \in so(n + 1) \right\}, \\ k &= \left\{ \begin{pmatrix} 0 & & & \\ & A & & \\ & & 0 & \\ & & & A \end{pmatrix} \mid A \in so(n) \right\}, \\ m &= \left\{ \begin{pmatrix} 0 & X & 0 & 0 \\ -X^t & 0 & 0 & 0 \\ 0 & 0 & 0 & X \\ 0 & 0 & -X^t & 0 \end{pmatrix} \mid X \in \mathbf{R}(1, n) \right\}, \\ g^\perp &= g_1^\perp \oplus g_2^\perp \oplus g_3^\perp, \\ m^\perp &= m_1^\perp \oplus m_2^\perp \oplus m_3^\perp, \end{aligned}$$

where

$$\begin{aligned}
 g_1^\perp &= \left\{ \begin{pmatrix} C & \\ & -C \end{pmatrix} \mid C \in so(n+1) \right\}, \\
 g_2^\perp &= \left\{ \begin{pmatrix} 0 & B \\ -B^t & 0 \end{pmatrix} \mid B = B^t \in gl(n+1) \right\}, \\
 g_3^\perp &= \left\{ \begin{pmatrix} 0 & B \\ -B^t & 0 \end{pmatrix} \mid B \in so(n+1) \right\}, \\
 m_1^\perp = 0, \quad m_2^\perp &= \left\{ \begin{pmatrix} & & & X \\ & & -X^t & \\ & X & & \\ -X^t & & & \end{pmatrix} \mid X \in \mathbf{R}(1, n) \right\}, \\
 m_3^\perp &= \left\{ \begin{pmatrix} & & & X \\ & & X^t & \\ & -X & & \\ -X^t & & & \end{pmatrix} \mid X \in \mathbf{R}(1, n) \right\}.
 \end{aligned}$$

It is easy to see

$$(g_2^\perp)^\mathbf{C} = (g_{21}^\perp)^\mathbf{C} \oplus (g_{22}^\perp)^\mathbf{C},$$

where

$$\begin{aligned}
 (g_{21}^\perp)^\mathbf{C} &= \lambda(0, \dots, 0), \quad (g_{22}^\perp)^\mathbf{C} = \lambda(2, 0, \dots, 0), \\
 (m_{21}^\perp)^\mathbf{C} &= 0, \quad (m_{22}^\perp)^\mathbf{C} = (m_2^\perp)^\mathbf{C}.
 \end{aligned}$$

Without loss of generality, suppose $n = 2s - 1$; then

$$(g_1^\perp)^\mathbf{C} = \lambda(0, 1, 0, \dots, 0), \quad (g_3^\perp)^\mathbf{C} = \lambda(0, 1, 0, \dots, 0).$$

As a K -module, it is easy to see

$$(m_2^\perp)^\mathbf{C} = \lambda'(1, 0, \dots, 0), \quad (m_3^\perp)^\mathbf{C} = \lambda'(1, 0, \dots, 0).$$

So $a_{21} = -(4s - 2)$, $a_3 = -4(s - 1)$. For $\lambda_k = \lambda(\underbrace{0, \dots, 0}_k, 1, 0, \dots, 0)$, we have

$$a_{\lambda_k} = -k(2s - k) \quad (1 \leq k \leq s - 2), \quad a_{\lambda_s} = a_{\lambda_{s-1}} = -s^2/2.$$

$$\begin{aligned}
 \{\lambda \in D(G) \mid a_\lambda > a_{21}\} &= \begin{cases} \{\lambda_1, \lambda_2\}, & s > 8, \\ \{\lambda_1, \lambda_2, \lambda_{s-1}, \lambda_s\}, & 3 \leq s \leq 7, \\ \{\lambda_1, \lambda_2, \lambda_3\}, & s = 3, \end{cases} \\
 \{\lambda \in D(G) \mid a_\lambda = a_{21}\} &= \lambda(2, 0, \dots, 0), \\
 \{\lambda \in D(G) \mid a_\lambda > a_3\} &= \begin{cases} \lambda_1, & s > 6, \\ \{\lambda_1, \lambda_{s-1}, \lambda_s\}, & s \leq 6, \end{cases} \\
 \{\lambda \in D(G) \mid a_\lambda = a_3\} &= \lambda_2.
 \end{aligned}$$

By the branching rule of representations, if $s > 3$ we get

$$\begin{aligned} \lambda_1 &= \lambda'(1, 0, \dots, 0) \oplus \lambda'(0, \dots, 0), \dim \lambda_1 = n + 1, \\ \lambda(2, 0, \dots, 0) &= \lambda'(2, 0, \dots, 0) \oplus \lambda'(1, 0, \dots, 0) \oplus \lambda'(0, \dots, 0), \\ \dim \lambda(2, 0, \dots, 0) &= \frac{(n + 1)(n + 2)}{2} - 1, \\ \lambda_2 &= \lambda'(0, 1, 0, \dots, 0) \oplus \lambda'(1, 0, \dots, 0), \dim \lambda_2 = n(n + 1)/2, \\ \lambda_s &= \lambda'(0, \dots, 0, 1) = \lambda_{s-1}, \\ \dim \lambda_{s-1} &= \dim \lambda_s = 2^{s-1}. \end{aligned}$$

If $s = 3$

$$\begin{aligned} \lambda(2, 0, 0) &= \lambda'(2, 0), \quad \lambda_1 = \lambda'(1, 0) = \lambda_3, \\ \lambda_2 &= \lambda'(0, 0) \oplus \lambda'(0, 1), \\ \dim \lambda(2, 0, 0) &= 10, \quad \dim \lambda_1 = \dim \lambda_3 = 4, \quad \dim \lambda_2 = 6. \end{aligned}$$

Thus we have

$$(13) \quad i(M) = \begin{cases} (n + 1)(n + 4)/2, & n > 5, \\ 16, & n = 5, \end{cases}$$

$$(14) \quad n(M) = \begin{cases} (n + 1)^2 - 1, & n > 5, \\ 0, & n = 5, \end{cases}$$

$$(15) \quad n_K(M) = (n + 1)^2 - 1.$$

Finally, we get the following theorem:

Theorem 3.1. *The indices, the nullities and the Killing nullities of the totally geodesic submanifolds in Q_m are listed in the following table. Among all the totally geodesic submanifolds in Q_m , only $Q_n (n < m)$ are stable; the nullity and the Killing nullity are equivalent except for $S^p \times S^q (p + q = m)$.*

M	$i(M)$	$n(M)$	$n_K(M)$
Q_n	0	$(n + 2)(m - n)$	$(n + 2)(m - n)$
$S^p \times S^q (p + q = m)$	$m + 2$	$2(p + 1)(q + 1)$	$(p + 1)(q + 1)$
$CP^n (m = 2n)$	$2(n + 1)$	$n(n + 1)$	$n(n + 1)$
$RP^n (m = 2n > 10)$	$(n + 1)(n + 4)/2$	$(n + 1)^2 - 1$	$(n + 1)^2 - 1$

REFERENCES

1. Y.Ohmita, On stability of minimal submanifolds in compact symmetric spaces, *Compositio Math.*, 64(1987), 157-189. MR **88k**:53082
2. B.Y.Chen, *Geometry of slant submanifolds*, Katholieke Universiteit Leuven, 1990. MR **92d**:53047
3. B.Y.Chen and T.Nagano, Totally geodesic submanifolds of symmetric spaces I, *Duke Math. J.*, 44(1977), 745-755. MR **56**:16543
4. E.B.Dynkin, Semisimple subalgebras of semisimple Lie algebras, *Amer. Math. Soc. Transl. Ser. 2*, vol. 6, Amer. Math. Soc., Providence, RI, 1957, pp. 111-244. (Russian original) MR **13**:904c

5. W.G.Mckay and J.Patera, Tables of dimensions, indices, and branching rules for representations of simple Lie algebras, Lecture Notes in Pure and Applied Mathematics, vol. 69, Marcel Dekker, New York and Basel, 1981. MR **82i**:17008

DEPARTMENT OF MATHEMATICS, BEIJING UNIVERSITY, BEIJING, 100871, PEOPLE'S REPUBLIC OF CHINA

DEPARTMENT OF MATHEMATICS, NORTHWEST NORMAL UNIVERSITY, LANZHOU, 730070, PEOPLE'S REPUBLIC OF CHINA