

RINGS OF WEAK DIMENSION ONE AND SYZYGETIC IDEALS

FRANCESC PLANAS-VILANOVA

(Communicated by Wolmer V. Vasconcelos)

ABSTRACT. We prove that rings of weak dimension one are the rings with all (three-generated) ideals syzygetic. This leads to a characterization of these rings in terms of the André-Quillen homology.

Let I be an ideal of a commutative ring A . There is a canonical morphism of graded A -algebras $\alpha : \mathbf{S}(I) \rightarrow \mathbf{R}(I)$ from the symmetric algebra of I onto its Rees algebra. The ideal I is said to be of linear type if α is an isomorphism. If $\alpha_2 : \mathbf{S}_2(I) \rightarrow I^2$ is an isomorphism, I is said to be syzygetic.

In [C] (Theorem 4), Costa showed that a domain A is Prüfer if and only if I is of linear type for every two-generated ideal I of A and I is syzygetic for every three-generated ideal I of A . In this note we show that the preliminary hypothesis that A is a domain can be removed by changing the Prüfer condition to the condition $wd(A) \leq 1$, weak dimension of A one or less. Moreover, the condition that every two-generated ideal of A be of linear type is not necessary. Concretely,

Theorem 1. *Let A be a commutative ring. The following conditions are equivalent:*

- i) $wd(A) \leq 1$.*
- ii) Every ideal of A is of linear type.*
- iii) Every ideal of A is syzygetic.*
- iv) Every three-generated ideal of A is syzygetic.*

Recall that $wd(A)$ is the supremum of the flat dimensions of all A -modules. The von Neumann regular rings are those of weak dimension zero. Semihereditary rings (i.e. rings with all their finitely generated ideals projective) have weak dimension one or less. In fact, A is a semihereditary ring if and only if $wd(A) \leq 1$ and A is coherent. A semihereditary domain is called a Prüfer ring. For a domain A , to be Prüfer is equivalent to $wd(A) \leq 1$ (see [B] or [R]). In particular, if A is a domain, Theorem 1 characterizes Prüfer rings as domains with every three-generated ideal being syzygetic.

Before proving Theorem 1 we need the following two lemmas.

Lemma 2. *Let (A, \mathfrak{m}) be a local ring. Suppose that there exist two nonzero elements $a, b \in A$ with $ab = 0$. If (a) is syzygetic, then (a, b) is not.*

Received by the editors November 4, 1994 and, in revised form, April 18, 1995.

1991 *Mathematics Subject Classification.* Primary 13F05, 13A30; Secondary 13D03.

Key words and phrases. Ideal of linear type, syzygetic ideal, weak dimension, homology of commutative rings.

Proof. Let $I = (a, b) \subset \mathfrak{m}$ be the ideal of A generated by the zero divisors a, b . If $x \in (a) \cap (b)$, $x = ac = bd$, and multiplying by $a, c \in (0 : a^2)$. Since (a) is syzygetic, $c \in (0 : a)$ (see [V], page 31) and $x = ca = 0$. Therefore, $(a) \cap (b) = 0$.

Let $0 \rightarrow Z_1 \rightarrow A^2 \xrightarrow{f} I \rightarrow 0$ be the free presentation of I defined by $f((1, 0)) = a, f((0, 1)) = b$. Consider $0 \rightarrow N \rightarrow A[X, Y] \xrightarrow{s} \mathbf{R}(I) \rightarrow 0$, the induced free presentation of $\mathbf{R}(I) = \bigoplus_{q \geq 0} I^q t^q$, defined by $s(X) = at, s(Y) = bt$.

If $I = (a, b)$ were syzygetic, the quadratic relation XY on a, b could be written in terms of linear relations on a, b ([V], page 29), i.e.

$$(1) \quad XY = (a_1X + b_1Y)(c_1X + d_1Y) + \dots + (a_rX + b_rY)(c_rX + d_rY)$$

with $c_iX + d_iY \in N_1 = Z_1$. In particular, $c_i a = -d_i b \in (a) \cap (b) = 0$. Therefore, $c_i a = d_i b = 0$ and if $a, b \neq 0$, then c_i, d_i would be zero divisors, in particular, elements of \mathfrak{m} . Comparing the coefficients of XY on both sides of (1) we would get the contradiction $1 = \sum_{i=1}^r a_i d_i + b_i c_i \in \mathfrak{m}$. \square

Lemma 3. *Let (A, \mathfrak{m}, k) be a local ring. Let I be a nonprincipal finitely generated ideal of A . If I is syzygetic, then I^2 is not.*

Proof. As I is not principal, $\dim_k(I/\mathfrak{m}I) = n > 1$. By hypothesis, α_2 is an isomorphism and hence, $\alpha_2 \otimes 1_k$ is also an isomorphism. Therefore,

$$\dim_k(I^2/\mathfrak{m}I^2) = \dim_k(\mathbf{S}_2^k(I/\mathfrak{m}I)) = \frac{n(n+1)}{2} = p.$$

Thus, $\dim_k(\mathbf{S}_2^k(I^2/\mathfrak{m}I^2)) = \frac{p(p+1)}{2} = \frac{n(n+1)(n^2+n+2)}{8}$. Since α_4 is an epimorphism, $\alpha_4 \otimes 1_k$ is also an epimorphism. So, one has

$$\dim_k(I^4/\mathfrak{m}I^4) \leq \dim_k(\mathbf{S}_4^k(I/\mathfrak{m}I)) = \frac{n(n+1)(n^2+5n+6)}{24}.$$

Finally, one observes that if $n \neq 0, 1$, then $3(n^2 + n + 2) > (n^2 + 5n + 6)$. In particular, $\mathbf{S}_2^k(I^2/\mathfrak{m}I^2) \not\subseteq I^4/\mathfrak{m}I^4$ and $\mathbf{S}_2^A(I^2) \not\subseteq I^4$. \square

Proof of the theorem. Recall $wd(A) \leq 1$ is equivalent to every ideal of A being flat ([R]). If I is a flat ideal of A , then I is an ideal of linear type (see Proposition 3 [MR] or [P]). This proves $i) \Rightarrow ii)$. Let us show that $iv) \Rightarrow i)$.

Let I be an ideal of A . To show I is flat, one can suppose I is finitely generated, since any ideal is the direct limit of finitely generated ideals, and the direct limit of flat modules is again a flat module. Write $I = (x_1, \dots, x_n)$ and let us see that $I_{\mathfrak{m}}$ is a free $A_{\mathfrak{m}}$ -module for every maximal ideal \mathfrak{m} of A . If $I \not\subseteq \mathfrak{m}$, $I_{\mathfrak{m}} = A_{\mathfrak{m}}$. If $I \subseteq \mathfrak{m}$, consider $J = (x_1, x_2) \subseteq I$. By hypothesis $iv)$, J and J^2 are syzygetic ideals of A . Localizing at \mathfrak{m} , Lemma 3 provides an element $z \in J$ with $J_{\mathfrak{m}} = (\frac{z}{1})$. Therefore $I_{\mathfrak{m}} = (\frac{z}{1}, \frac{x_3}{1}, \dots, \frac{x_n}{1})$. Now, take $J' = (z, x_3) \subseteq I$. Repeating the process we deduce that $I_{\mathfrak{m}}$ is a principal ideal of $A_{\mathfrak{m}}$. By hypothesis $iv)$ and Lemma 2, $A_{\mathfrak{m}}$ is a domain, in particular, $I_{\mathfrak{m}}$ is an ideal generated by a nonzero divisor, i.e. a free $A_{\mathfrak{m}}$ -module. \square

Remark 4. From Lemma 2 and Costa's Theorem 3 of [C] one deduces that for a commutative ring A to be a locally an integrally closed domain is equivalent to being of linear type for every two-generated ideal of A . Moreover, the same example given by Costa in his paper shows that this last condition is strictly stronger than every two-generated ideal of A being syzygetic.

In terms of the André-Quillen homology (see [A]) and as a corollary of Theorem 1, rings of weak dimension one are characterized as follows:

Corollary 5. *Let A be a commutative ring. The following conditions are equivalent:*

- i) $wd(A) \leq 1$.*
- ii) $H_2(A, B, \cdot) = 0$ for every quotient ring $B = A/I$ of A .*
- iii) $H_2(A, B, B) = 0$ for every quotient ring $B = A/I$ of A by a three-generated ideal I of A .*

Proof. It follows from the fact that if I is an ideal of A , $B = A/I$ and $\alpha_2 : \mathbf{S}_2(I) \rightarrow I^2$ is the canonical morphism, then $H_2(A, B, B) = \text{Ker } \alpha_2$. Moreover, if I is syzygetic, $H_2(A, B, W) = \text{Tor}_1^B(I/I^2, W)$ for any B -module W (see, for instance, [BR]). \square

ACKNOWLEDGEMENTS

I am very grateful to José M. Giral for the very helpful discussions concerning this paper.

REFERENCES

- [A] M. André: Homologie des algèbres commutatives. Grundlehren **206**. Heidelberg: Springer 1974. MR **50**:4707
- [BR] J. Barja, A. G. Rodicio: *Syzygetic Ideals, Regular sequences and a Question of Simis*. J. Algebra **121** (1989), 310-314. MR **90b**:13017
- [B] N. Bourbaki: Algèbre. Chapitre 10. Algèbre homologique. Masson. Paris, 1980. MR **82j**:18022
- [C] D. Costa: *On the Torsion-Freeness of the Symmetric Powers of an Ideal*. J. Algebra **80** (1983), 152-158. MR **85e**:13019
- [MR] A. Micali, N. Roby: *Algèbres symétriques et syzygies*. J. Algebra **17** (1971), 460-469. MR **44**:198
- [P] F. Planas Vilanova: Ideals de tipus lineal i homologia d'André-Quillen. Ph.D. Thesis. Universitat de Barcelona, 1994.
- [R] J. Rotman: An Introduction to Homological Algebra. Academic Press, 1979. MR **80k**:18011
- [V] W. V. Vasconcelos: Arithmetic of Blowup Algebras. LMS Lecture Note Series **195**. Cambridge University Press, 1994. MR **95g**:13005

DEPARTAMENT DE MATEMÀTICA APLICADA I, ETSEIB, UNIVERSITAT POLITÈCNICA DE CATALUNYA, DIAGONAL 647, E-08028 BARCELONA, SPAIN

E-mail address: `planas@ma1.upc.es`