

A SIMPLE PROOF OF SINGER'S REPRESENTATION THEOREM

WOLFGANG HENSGEN

(Communicated by Dale E. Alspach)

ABSTRACT. Let Ω be a compact Hausdorff space and X a Banach space. Singer's theorem states that under the dual pairing $(f, m) \mapsto \int \langle f, dm \rangle$, the dual space of $C(\Omega; X)$ is isometric to $rcabv(\Omega; X')$. Using the Hahn-Banach theorem and the (scalar) Riesz representation theorem, a proof of Singer's theorem is given which appears to be simpler than the proofs supplied earlier by Singer (1957, 1959) and Dinculeanu (1959, 1967).

The notation of the abstract is supposed to be self-explanatory. A Borel measure $m \in cabv(\Omega; X)$ is called regular ($m \in rcabv$) iff its variation $|m|$ is regular. (As a corollary of our approach it is shown below that this is equivalent to the weakest possible notion of regularity.)

Proof of Singer's representation theorem. a) Let $m \in cabv(\Omega; X')$; then routine verifications show that $\varphi_m(f) := \int \langle f, dm \rangle$ defines a functional $\varphi_m \in C(\Omega; X)'$ of norm $\|\varphi_m\| \leq \|m\|$ ($= |m|(\Omega)$). Under the additional requirement that $\langle x, m \rangle$ be regular $\forall x \in X$, m is uniquely determined by φ_m , by the uniqueness part of the (scalar) Riesz theorem.

b) Conversely, let $\varphi \in C(\Omega; X)'$ be given. Equip the unit ball $B_{X'}$ with the weak* topology and consider the isometric embedding $C(\Omega; X) \hookrightarrow C(\Omega \times B_{X'})$ sending f into the scalar function $(\omega, x') \mapsto \langle f(\omega), x' \rangle$. The Hahn-Banach theorem combined with the Riesz representation theorem [R, 6.19] produces a complex regular Borel measure ν on $\Omega \times B_{X'}$ satisfying $\|\nu\| = \|\varphi\|$ and $\forall f \in C(\Omega; X) : \varphi(f) = \int_{\Omega \times B_{X'}} \langle f(\omega), x' \rangle \nu(d\omega, dx')$. In particular, $\forall u \in C(\Omega) \forall x \in X$:

$$\varphi(u \otimes x) = \int_{\Omega \times B_{X'}} u(\omega) \langle x, x' \rangle \nu(d\omega, dx') = \int_{\Omega} u d\mu_x$$

for this Borel measure μ_x on Ω :

$$\mu_x(A) := \int_{\Omega \times B_{X'}} \mathbf{1}_A(\omega) \langle x, x' \rangle \nu(d\omega, dx').$$

Received by the editors April 21, 1995.

1991 *Mathematics Subject Classification.* Primary 46E15, 46E40.

Key words and phrases. Vector-valued continuous functions, regular vector measures.

©1996 American Mathematical Society

Obviously

$$\begin{aligned} |\mu_x(A)| &\leq \|x\| \int_{\Omega \times B_{X'}} \mathbf{1}_A(\omega) |\nu|(d\omega, dx') \\ &= \|x\| \int_{\Omega} \mathbf{1}_A(\omega) \nu_1(d\omega) = \|x\| \nu_1(A), \end{aligned}$$

where $\nu_1 := |\nu| \circ \text{pr}_1^{-1}$ is a positive *regular* Borel measure on Ω , as is easily verified. Defining, for Borel sets $A \subset \Omega$ and $x \in X$, $m(A)(x) := \mu_x(A)$, we have immediately $m(A) \in X'$, $\|m(A)\| \leq \nu_1(A)$, hence $|m| \leq \nu_1$ so that $m \in \text{rcabv}(\Omega; X')$ and $\|m\| \leq \nu_1(\Omega) = \|\nu\| = \|\varphi\|$. By construction, m represents φ on $C(\Omega) \otimes X$ which is dense in $C(\Omega; X)$ [DU, p.225]. Invoke a) to conclude that m represents φ on $C(\Omega; X)$. \square

The following corollary is proved directly in [S1, Lemma 3].

Corollary. *For $m \in \text{cabv}(\Omega; Y)$, the following are equivalent:*

- 1° $|m|$ is regular.
- 2° $\langle m, y' \rangle$ is regular $\forall y' \in X$, a norming subspace of Y' .

Proof. To prove 2° \Rightarrow 1°, we can assume $Y = X'$. Suppose $\langle x, m \rangle$ is regular $\forall x \in X$. The functional $\varphi_m \in C(\Omega; X)'$ is represented by an $m' \in \text{rcabv}(\Omega; X')$ which implies $m = m'$ as noted in a) above. \square

REFERENCES

- [DU] J. Diestel, J.J. Uhl, Jr., *Vector Measures*, AMS, Providence, 1977 (Math. Surveys **15**). MR **56**:11216
- [D1] N. Dinculeanu, *Sur la représentation intégrale des certaines opérations linéaires. III*, Proc. AMS **10** (1959), 59–68. MR **21**:2909
- [D2] N. Dinculeanu, *Vector Measures*, Pergamon Press, Oxford etc., 1967. MR **34**:6011b
- [R] W. Rudin, *Real and Complex Analysis*, 3rd ed., McGraw Hill, New York etc., 1987. MR **88k**:00002
- [S1] I. Singer, *Linear functionals on the space of continuous mappings of a compact Hausdorff space into a Banach space* (in Russian), Rev. Roum. Math. Pures Appl. **2** (1957), 301–315. MR **20**:3445
- [S2] I. Singer, *Sur les applications linéaires intégrales des espaces de fonctions continues. I*, Rev. Roum. Math. Pures Appl. **4** (1959), 391–401. MR **22**:5883

NWF I – MATHEMATIK, UNIVERSITÄT REGENSBURG, D– 93040 REGENSBURG, GERMANY
E-mail address: wolfgang.hensgen@mathematik.uni-regensburg.de