

CHARACTERIZATION OF CLASSICAL GROUPS BY ORBIT SIZES ON THE NATURAL MODULE

MARTIN W. LIEBECK

(Communicated by Ronald M. Solomon)

ABSTRACT. We show that if V is a finite vector space, and G is a subgroup of $P\Gamma L(V)$ having the same orbit sizes on 1-spaces as an orthogonal or unitary group on V , then, with a few exceptions, G is itself an orthogonal or unitary group on V .

Let C be a finite orthogonal or unitary group, with associated vector space V , and let G be a subgroup of $P\Gamma L(V)$ having the same orbit sizes as C on the set of 1-dimensional subspaces of V . We shall show that, with a few exceptions, under these hypotheses G must itself be an orthogonal or unitary group on V . The precise result is stated in the Theorem below.

This paper was written in response to a question of Prof. S. Abhyankar, who makes use of the result in [Ab].

Theorem. *Let q be a prime power, $d \geq 3$ an integer, and $V = V_d(q)$ a vector space of dimension d over \mathbb{F}_q . Suppose that G is a subgroup of $P\Gamma L_d(q)$ such that the sizes of the orbits of G on the 1-spaces of V are as in one of cases (1)–(5) in Table 1 below. If $d \leq 7$, assume that $q > 2$; and if $d \leq 4$, assume that $q > 3$.*

(a) *If the orbit sizes are as in (4) or (5) of Table 1, then either $G \triangleright PSU_d(q^{1/2})$, or $d = 3, q = 4, G \triangleright 3^2 = O_3(PSU_3(2))$ (and if also $G \leq PGL_3(4)$, then $G \triangleright PSU_3(2)$).*

(b) *If the sizes are as in (2), then either $G \triangleright P\Omega_{2m}^+(q)$, or $G \triangleright PSU_m(q)$ (m even), or $d = 8, G \triangleright \Omega_7(q)$ (embedded irreducibly in $PSL_8(q)$), or $d = 4, q = 5, G \triangleright A_6$.*

(c) *If the sizes are as in (3), then either $G \triangleright P\Omega_{2m}^-(q)$, or $G \triangleright PSU_m(q)$ (m odd), or $d = 4, G \triangleright {}^2B_2(q)$ (q an odd power of 2), or $d = 6, q = 3, G \triangleright L_3(4)$, or $d = 4, q = 4, G \leq \Gamma L_1(2^8)$.*

(d) *If the orbit sizes are as in (1) with q odd, then either $G \triangleright \Omega_{2m+1}(q)$, or $d = 7, G \triangleright G_2(q)$.*

(e) *If the orbit sizes are as in (1) with q even, then $G \triangleright Sp_{2a}(q^b)$ ($2ab = 2m$) or $G \triangleright G_2(q^b)$ ($6b = 2m$), embedded naturally in $Sp_{2m}(q)$ acting indecomposably on V (fixing a $2m$ -dimensional subspace W , with V/W trivial).*

Conversely, all the groups arising in (a)-(e) do have orbit sizes as in Table 1.

Remarks. 1. The classical groups arising in conclusions (a)-(d) all act in the natural way on V .

Received by the editors March 20, 1995.

1991 *Mathematics Subject Classification.* Primary 20G40.

2. If we relax the assumptions in the Theorem made on q for small d , more examples occur, but it would not be hard to list these.

TABLE 1

| Case | d | orbit sizes |
|------|----------|--|
| (1) | $2m + 1$ | $\frac{q^{2m}-1}{q-1}, \frac{1}{2}q^m(q^m + 1), \frac{1}{2}q^m(q^m - 1)$ |
| (2) | $2m$ | $\frac{(q^m-1)(q^{m-1}+1)}{q-1}, q^{m-1}(q^m - 1)$ or $\frac{(q^m-1)(q^{m-1}+1)}{q-1}, \frac{1}{2}q^{m-1}(q^m - 1), \frac{1}{2}q^{m-1}(q^m - 1)$ (q odd) |
| (3) | $2m$ | $\frac{(q^m+1)(q^{m-1}-1)}{q-1}, q^{m-1}(q^m + 1)$ or $\frac{(q^m+1)(q^{m-1}-1)}{q-1}, \frac{1}{2}q^{m-1}(q^m + 1), \frac{1}{2}q^{m-1}(q^m + 1)$ (q odd) |
| (4) | $2m$ | $\frac{(q^m-1)(q^{m-\frac{1}{2}}+1)}{q-1}, \frac{q^{m-\frac{1}{2}}(q^m-1)}{q^{\frac{1}{2}}+1}$ (q square) |
| (5) | $2m + 1$ | $\frac{(q^{m+\frac{1}{2}}+1)(q^m-1)}{q-1}, \frac{q^m(q^{m+\frac{1}{2}}+1)}{q^{\frac{1}{2}}+1}$ (q square) |

In the proof we shall use *primitive prime divisors*: if $n \geq 3$ and $(q, n) \neq (2, 6)$, then by [Zs] there is a prime which divides $q^n - 1$ but does not divide $q^i - 1$ for $1 \leq i \leq n - 1$. Such a prime is called a primitive prime divisor of $q^n - 1$ and is denoted by q_n ; note that $q_n \equiv 1 \pmod n$. Write q_n^* for the product of all primitive prime divisors of $q^n - 1$, counting multiplicities.

The proof of the Theorem is a fairly routine application of the results in [Li] and [GPPS]. The paper [Li] determines the irreducible subgroups of $P\Gamma L_d(q)$ having exactly two orbits on 1-spaces, and can be used to handle such cases in the Theorem. And [GPPS] lists subgroups of $P\Gamma L_d(q)$ which have order divisible by q_e for some $e > \frac{1}{2}d$; since our group G is in general divisible by such a prime, this applies to our problem. Despite the routine nature of the proof, we feel that the result may be of some interest, especially in view of the application [Ab].

Proof of the Theorem. Let $G \leq P\Gamma L_d(q)$ be as in the statement of the Theorem. Write $q = p^f$ with p prime. Assume first that $d \geq 5$, and also that $q > 2$ if $d \leq 8$; we shall handle the excluded cases later. Referring to Table 1, we see that $|G|$ is divisible by q_d^*, q_{d-1}^* or q_{d-2}^* in case (1), (2) or (3), and by $(q^{1/2})_{2d}^*$ or $(q^{1/2})_{2d-2}^*$ in case (4) or (5).

Suppose first that G is reducible on V . Then $G \leq P_i$, the stabilizer in $P\Gamma L(V)$ of an i -space, for some i . The orbit sizes of P_i on 1-spaces are $\frac{q^i-1}{q-1}$ and $\frac{q^i(q^{d-i}-1)}{q-1}$, so one of these is an orbit size of G . The only possibility is that $d = 2m + 1, i = 2m$ and the orbit sizes of G are as in case (1) of Table 1. Thus $G \leq P_{2m} = QL$, where $Q \cong (\mathbf{F}_q)^{2m}$ is the unipotent radical and $L \triangleright SL_{2m}(q)$ is a Levi subgroup. The orbit sizes of P_{2m} are $\frac{q^{2m}-1}{q-1}, q^{2m}$; G is transitive on the first orbit, and Q is transitive on the second. Therefore $Q \not\leq G$. As G acts irreducibly on Q , it follows that $G \cap Q = 1$, whence G is isomorphic to a subgroup of L which is transitive on the 1-spaces of $V_{2m}(q)$. The list of all transitive linear groups is given in [Li, Appendix 1], and the only possibilities which are divisible by the orbit sizes in (1), hence by $\frac{1}{2}q^m(q^{2m} - 1)/(q - 1)$, are as follows:

- (i) $G \leq \Gamma L_1(p^{2fm})$ (where $q = p^f$);

- (ii) $G \triangleright S = Sp_{2a}(q^b)$ (where $2ab = 2m, a \geq 1$);
- (iii) $G \triangleright S = SL_a(q^b)$ (where $ab = 2m, a \geq 3$);
- (iv) $G \triangleright S = G_2(q^b)$ (where $6b = 2m, q$ even).

In case (i) the divisibility condition forces p^{fm} to divide $4fm$, whence $p = 2, fm = 4$; but the subgroup $G \cap GL_1(2^8)$, being of odd order, fixes a 1-space of V , so G has an orbit size dividing $|G : G \cap GL_1(2^8)|$, hence dividing 8, which is not the case.

In case (iii), or in case (ii) with q odd, we have $H^1(S, Q) = 0$ by [JP]. It follows that the subgroup S of G is conjugate to a subgroup of L , hence fixes a 1-space of V . But then G has an orbit of size dividing $|G : S|$, which is not so.

In the remaining cases ((ii) with q even, and (iv)), $H^1(S, Q)$ has dimension 1 by [JP]; by the argument of the previous paragraph, S does not fix a 1-space, so S is not conjugate to a subgroup of L . It follows that conclusion (e) of the Theorem holds. Here S lies in a subgroup $Sp_{2m}(q)$ acting indecomposably on V , with orbit sizes as in (1) of Table 1 and point stabilizers P_1 (a parabolic), $O_{2m}^+(q)$ and $O_{2m}^-(q)$. The group S is transitive on each of the orbits, since $Sp_{2m}(q)$ factorizes as $S.P_1 = S.O_{2m}^+(q) = S.O_{2m}^-(q)$ (see [LPS, Tables 1 and 2]).

Now assume that G is irreducible on V . Write $Z = Z(GL_d(q))$. Choose an integer b , maximal such that $G \leq \Gamma L_a(q^b)/Z$ ($ab = d$) in its usual embedding in $P\Gamma L_d(q)$. If $a = 1$ then $|G|$ divides $(q^d - 1)df$ (where $q = p^f$), which is impossible since orbit sizes in Table 1 divide $|G|$. Hence $a \geq 2$.

The subgroups of $\Gamma L(V)$ having two orbits on 1-spaces, and their orbit sizes, are listed in [Li, Appendix 2]. A glance at this list shows that the only such groups having orbit sizes as in Table 1 satisfy conclusion (a), (b) or (c) of the Theorem. Thus we may assume that G has three orbits on 1-spaces; the orbit sizes are then as in (1), (2) or (3) of Table 1, with q odd in cases (2), (3). In particular, $|G|$ is divisible by q_e^* , where $e = d, d - 1$ or $d - 2$.

Let X be one of the classical groups $SL_a(q^b), Sp_a(q^b), \Omega_a(q^b), U_a(q^{b/2})$, chosen to be minimal such that $G \leq N_{\Gamma L(V)}(X)/Z$. Write $\bar{X} = X/X \cap Z$. If G contains \bar{X} then X must be orthogonal or unitary, and from the orbit sizes of X we see that $b = 1$ in the orthogonal case, $b = 1$ or 2 in the unitary case; hence G is as in (a)-(d) of the Theorem. Consequently we may assume that $\bar{X} \not\leq G$.

At this point we apply the main result of [As] on the subgroups of the classical group $N_{P\Gamma L(V)}(\bar{X})$. According to this result, either G lies in a member of one of the families $\mathcal{C}_1, \dots, \mathcal{C}_8$ of subgroups of this group, or $G \in \mathcal{S}$, a certain collection of almost simple subgroups. A discussion of this result can be found in [KL, Chapter 1], and detailed descriptions of the members of the families \mathcal{C}_i in [KL, Chapter 4].

Suppose first that $G \in \mathcal{C}_i$ for some i . As G is irreducible, and by choice of b and X , i is not 1, 3 or 8; also subgroups in \mathcal{C}_i for $i = 4, 5, 7$ do not have order divisible by q_e^* . If $G \leq M \in \mathcal{C}_2$, then G stabilizes a decomposition $V = V_1 \oplus \dots \oplus V_k$, where each V_i has \mathbb{F}_q -dimension $r, rk = d$ and $G \cap PGL(V) \leq (GL_r(q) \text{ wr } S_k)/Z$. As q_e^* divides $|G|$ and $e \geq d - 2 > d/2$, we must have $r = 1, k = d$ and $q_e^* = d - 1$ or d . But G has at least k orbits on 1-spaces, so this is impossible when $d \geq 5$. Finally, suppose that $G \leq M \in \mathcal{C}_6$. Then $|M \cap PGL(V)|$ divides $r^{2k}|Sp_{2k}(r)|$, where r is prime, $a = r^k$ and $r|q^b - 1$. Since q_e^* divides $|G|$, this means that $r = 2, a = d = 2^k$ and $q_e^* = 2^k \pm 1$ with $e = d$ or $d - 2$. A result of Hering [He, 3.9] determines all (q, e) such that $q_e^* = e + 1$, and this implies that $(q, e) = (3, 4), (3, 6)$ or $(5, 6)$. In the first case $d = 4$, contrary to assumption; in the second case $G \leq 2^6.Sp_6(2) < L_8(3)$, and

the orbit sizes of $2^6 \cdot Sp_6(2)$ on 1-spaces are 720, 2560 by [Li, Appendix 2], neither of which is an orbit size of G ; and in the last case the orbit sizes of G do not divide $2^6 |Sp_6(2)|$.

It remains to deal with the case where $G \in \mathcal{S}$. Here G is almost simple; write $S = F^*(G)$. In [GPPS, Examples 2.6 - 2.9], all possibilities for subgroups in \mathcal{S} which are divisible by a primitive prime divisor q_{ib} , $i > a/2$, are listed. Clearly $e = ib$ with $i > a/2$, so our group S is in this list.

Suppose first that S is of Lie type in characteristic p . Then S is given by [GPPS, Example 2.8]. The only possibilities with $|S|$ divisible by q_e^* ($e \geq d-2$) are $(S, d, e) = (L_2(q^3), 8, 6)$, $(\Omega_7(q), 8, 6)$, $(G_2(q)$ or ${}^2G_2(q), 7, 6)$ (q odd), $(G_2(q^2), 14, 12)$ (q odd), $(U_3(q), 8 - \delta_{3,p}, 6)$ or $(U_3(q^2), 14, 12)$ ($p = 3$). Of these, the only cases where $|\text{Aut } S|$ is divisible by the orbit sizes in Table 1 are $S = \Omega_7(q)$, $G_2(q)$, as in conclusions (b), (d) of the Theorem. In these cases S lies in $\Omega_8^+(q)$, $\Omega_7(q)$ respectively, and is transitive on each orbit of these groups on 1-spaces (see [LPS] again).

Now assume that S is alternating, sporadic, or of Lie type in p' -characteristic. From the lists in [GPPS, Examples 2.6, 2.7 and 2.9], we see that one of the following holds:

(i) there is only *one* primitive prime divisor of $q^e - 1$ dividing $|G|$, and this is equal to $e + 1$ or $2e + 1$; moreover, this prime divides $|G|$ to the first power only;

(ii) $S = L_2(s)$ with s prime, $d = (s \pm 1)/2$, $e = (s - 1)/2$ and $q_e^* = (e + 1)(2e + 1)$.

Consequently q_e^* must be $e + 1$, $2e + 1$ or $(e + 1)(2e + 1)$. Hence the possibilities for (q, e) are given by [He, 3.9]. In case (ii), (q, e) is $(3, 18)$ or $(17, 6)$. But then either $S = L_2(37) < L_d(3)$ ($d = 18$ or 19), or $S = L_2(13) < L_d(17)$ ($d = 6$ or 7), and $|\text{Aut } S|$ is not divisible by the orbit sizes. Therefore (i) holds, and the possibilities for q, e are as follows:

$$\begin{aligned} q = 2 : & \quad e = 3, 4, 8, 10, 12, 18 \text{ or } 20 \\ q = 3 : & \quad e = 4 \text{ or } 6 \\ q = 4 : & \quad e = 3 \text{ or } 6 \\ q = 5 : & \quad e = 6. \end{aligned}$$

Suppose that $q = 2$ or 4 . Since we are assuming G to have three orbits on 1-spaces, we must have $d = 2m + 1$ and $|G|$ divisible by the orbit sizes $\frac{q^{2m}-1}{q-1}$, $\frac{1}{2}q^m(q^m + 1)$, $\frac{1}{2}q^m(q^m - 1)$. (In particular, $e = 2m = d - 1$.) From [GPPS], we see that the only possibilities for $S = F^*(G)$ satisfying these conditions are $S = PSp_4(5)$ or $PSp_6(3)$, with $q = 2, d = 13$. However, $PSp_4(5), PSp_6(3)$ are not subgroups of $L_{13}(2)$ by Lagrange's theorem.

Now let $q = 3$. Here $e = 4$ or 6 , so $5 \leq d \leq 8$. If $d = 5$, the orbit sizes 40, 45, 36 divide $|G|$, and [GPPS] shows that $S = M_{11}$; however, $M_{11} < L_5(3)$ has only two orbits on 1-spaces, by [Li]. If $d = 6$ then the orbit sizes of G are 130, 117, 117 or 112, 126, 126, whence $|G|$ is divisible by either $3^2 \cdot 5 \cdot 13$ or $2^4 \cdot 3^2 \cdot 7$; now [GPPS] implies that S is $L_3(4), A_7$ or J_2 . The first case is in conclusion (c) of the Theorem, and $L_3(4) < \Omega_6^-(3)$ is transitive on each $\Omega_6^-(3)$ -orbit on 1-spaces (see [At, p. 52]). In the second case, $G = A_7$ or S_7 has orbit sizes 112, 126, 126; but A_7 and S_7 have no transitive actions of degree 112 (see [At, p.10]). Finally, $J_2 \not\leq L_6(3)$ by Lagrange's theorem. When $d = 7$, [GPPS] gives no possibilities for G of order divisible by the orbit sizes $2^2 \cdot 7 \cdot 13$, $2 \cdot 3^3 \cdot 7$, $3^3 \cdot 13$. Now suppose $d = 8$. The orbit sizes imply that $|G|$ is divisible by either $2^5 \cdot 3^3 \cdot 5 \cdot 7$ (case (2) of Table 1) or by $13 \cdot 41$ (case (3)). By [GPPS], the former holds, and $S = Sp_6(2), \Omega_8^+(2), A_9$ or $L_3(4)$. In the first three cases the 8-dimensional representation of S is uniquely determined (see the 3-modular character

tables of these groups in [At2]), and embeds $S \leq \Omega_8^+(2) < \Omega_8^+(3) < L_8(3)$. But in this representation, $\Omega_8^+(2)$ has an orbit of size 120 on 1-spaces (see [Li, p.505, case 3(e,f)]). Finally, if $S = L_3(4)$, one checks using [At, p.23] that the group G with $F^*(G) = S$ has no transitive action of degree $2^3 \cdot 3^3 \cdot 5$, which is one of the required orbit sizes.

Now suppose that $q = 5$. Then $e = 6$ and $6 \leq d \leq 8$. When $d = 6$, $|G|$ is divisible by either 13.31 (case (2) of Table 1) or $2^2 \cdot 3^3 \cdot 5^2 \cdot 7$ (case (3)). By [GPPS], the latter holds, and $S = J_2$. But $J_2 < L_6(5)$ has only two orbits on 1-spaces, by [Li]. Finally, when $d = 7$ or 8, [GPPS] shows that there are no possibilities for S with $|G|$ divisible by the required orbit sizes.

This completes the proof of the Theorem under our initial assumption that $d \geq 5$ and that $q > 2$ if $d \leq 8$. By the hypotheses of the Theorem, it remains to handle the cases $d = 3, 4$ with $q \geq 4$, and $d = 8, q = 2$. The argument given at the beginning of the proof of the Theorem (second paragraph) shows that if G is reducible then $d = 3, q$ is even and $G \triangleright Sp_2(q)$ as in conclusion (e). Thus we suppose that G is irreducible. In the case $d = 8, q = 2$, G has two orbits on 1-spaces, and we check that conclusion (b) or (c) holds using [Li]. Thus we suppose from now on that $d = 3$ or 4 and $q \geq 4$.

As in the proof above, we choose b maximal such that $G \leq \Gamma L_a(q^b)/Z$, where $ab = d$. If $a = 1$ then the orbit sizes must divide $(q^d - 1) \cdot d \log_p q$, which implies that $d = 4$ and $q = 4, 8$ or 16 (and the orbit sizes are as in (3) of Table 1). The subgroups of $\Gamma L_1(q^d)$ having two orbits on nonzero vectors are given by [FK, §3], from which we see that an example arises if and only if $q = 4$, as in conclusion (c).

Hence we now assume that $a \geq 2$; and $(a, b) = (d, 1)$ or $(2, 2)$. Again choose a classical group X of dimension a over \mathbf{F}_{q^b} , minimal such that $G \leq N_{\Gamma L(V)}(X)/Z$. If G contains $\bar{X} = X/X \cap Z$, then one of (a)-(d) of the Theorem holds, so assume $\bar{X} \not\leq G$. Suppose that G is contained in a member M of one of the families \mathcal{C}_i of subgroups of $N(\bar{X})$. Then $i \neq 1, 3, 4, 7, 8$ by choice of b and X . If $i = 2$ or 5 then the orbit sizes of M are not compatible with those of G . And if $M \in \mathcal{C}_6$, then $|G \cap L_d(q)|$ divides $2^4 \cdot 3^3$ (if $d = 3$), or $2^8 \cdot 3^2 \cdot 5$ (if $d = 4$). The fact that $|G|$ is divisible by orbit sizes in Table 1 forces either $d = 3, q = 4$ or $d = 4, q = 5$. In the first case, $G \leq 3^2 \cdot 2S_4$ (see [At, p.23]) and G has orbit sizes 9, 12 as in (5) of Table 1. Then clearly $G \triangleright 3^2$, as in (a) of the Theorem; moreover, if also $G \leq PGL_3(4)$, then $G \leq 3^2 \cdot 2A_4$, whence from the action on the orbits, G contains $3^2 \cdot Q_8 = PSU_3(2)$. Now consider $d = 4, q = 5$. Here $G \leq 2^4 \cdot Sp_4(2)$ and the orbit sizes of G are 36, 60, 60 or 36, 120. By [Li], the group $2^4 \cdot Sp_4(2)$ has orbit sizes 60, 96. The normal subgroup 2^4 has 15 orbits of size 4 and 16 of size 6, both sets permuted transitively by the factor $Sp_4(2)$ (see [Li, 1.2]). Hence G cannot contain this 2^4 , and so $G \cap 2^4 = 1$. Thus $G = A_6$ or S_6 , as in conclusion (b) of the Theorem.

Thus G lies in the collection \mathcal{S} of almost simple subgroups of $N_{P\Gamma L(V)}(\bar{X})$. Let $S = F^*(G)$. For $d \leq 4$, the members of this collection are well known, and are among the following (see [Kl, Chapter 5]):

$$\begin{aligned} d = 3: & \quad S = A_5, A_6 \text{ or } L_2(7) \\ d = 4: & \quad S = A_5, A_6, A_7, L_2(7), U_4(2), L_3(4) \text{ or } L_2(q). \end{aligned}$$

By [Li], the only possibility for G having two orbits is $d = 4, q = 5, S = A_6$, as in (b) of the Theorem. So assume that G has three orbits; these have sizes $q + 1, \frac{1}{2}q(q + 1), \frac{1}{2}q(q - 1)$, or $(q + 1)^2, \frac{1}{2}q(q^2 - 1), \frac{1}{2}q(q^2 - 1)$, or $q^2 + 1, \frac{1}{2}q(q^2 + 1), \frac{1}{2}q(q^2 + 1)$. The only possibilities with $|\text{Aut } S|$ divisible by orbit sizes are as

follows:

$$d = 3, q = 4, S = A_5 \text{ or } A_6, \text{ and}$$

$$d = 4, q = 5, S = A_6 \text{ or } A_7.$$

In the case $d = 3$, A_6 has orbit sizes 6,15 (see [Li]), and there is no irreducible A_5 in $L_3(4)$. And in the case $d = 4$, A_6 has two orbits ([Li]) and there is no A_7 in $L_4(5)$.

This completes the proof of the Theorem.

REFERENCES

- [Ab] S. Abhyankar, "Again nice equations for nice groups", *Proc. Amer. Math. Soc.* **124** (1996), 3567–3576.
- [As] M. Aschbacher, "On the maximal subgroups of the finite classical groups", *Invent. Math.* **76** (1984), 469–514. MR **86a**:20054
- [At] J.H. Conway, R.T. Curtis, S.P. Norton, R.A. Parker and R.A. Wilson, *Atlas of finite groups*, Oxford University Press, 1985. MR **88g**:20025
- [FK] D.A. Foulser and M.J. Kallaher, "Solvable, flag-transitive, rank 3 collineation groups", *Geom. Ded.* **7** (1978), 111–130. MR **57**:12263
- [GPPS] R. Guralnick, T. Penttila, C.E. Praeger and J. Saxl, "Linear groups with orders having certain primitive prime divisors", preprint, University of Western Australia, 1994.
- [He] C. Hering, "Transitive linear groups and linear groups which contain irreducible subgroups of prime order", *Geom. Ded.* **2** (1974), 425–460. MR **49**:439
- [At2] C. Jansen, K. Lux, R.A. Parker and R.A. Wilson, *Atlas of modular characters*, Oxford University Press, 1995.
- [JP] W. Jones and B. Parshall, "On the 1-cohomology of finite groups of Lie type", *Proc. Conf. on Finite Groups*, eds. W. Scott and F. Gross, Academic Press, (1976). MR **53**:8272
- [Kl] P.B. Kleidman, *The subgroup structure of some finite simple groups*, Ph.D. Thesis, University of Cambridge, 1987.
- [KL] P.B. Kleidman and M.W. Liebeck, *The Subgroup Structure of the Finite Classical Groups*, London Math. Soc. Lecture Note Series **129**, Cambridge University Press, Cambridge, 1990. MR **91g**:20001
- [Li] M.W. Liebeck, "The affine permutation groups of rank three", *Proc. London Math. Soc.* **54** (1987), 477–516. MR **88m**:20004
- [LPS] M.W. Liebeck, C.E. Praeger and J. Saxl, "The maximal factorizations of the finite simple groups and their automorphism groups", *Mem. Amer. Math. Soc.* **86**, No. 432 (1990), 1–151. MR **90k**:20048
- [Zs] K. Zsigmondy, "Zur Theorie der Potenzreste", *Monatsh. fur Math. und Phys.* **3** (1892), 265–284.

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE, LONDON SW7 2BZ, UNITED KINGDOM
E-mail address: m.liebeck@ic.ac.uk