

THE STANDARD CANTOR FUNCTION IS SUBADDITIVE

JOZEF DOBOŠ

(Communicated by Andreas R. Blass)

ABSTRACT. In this paper the subadditivity of the Cantor function $\phi: [0, 1] \rightarrow [0, 1]$ is proved.

Let us begin by recalling that a function $f: \mathbb{R} \rightarrow \mathbb{R}$ is said to be subadditive, if it satisfies the inequality $f(x + y) \leq f(x) + f(y)$ whenever $x, y \in \mathbb{R}$. (See [2] and [4].)

The usual definition of the standard Cantor function involves the classic middle-thirds description of the standard Cantor set. (See [1] and [3].) We offer an alternate definition of this function.

Define a sequence of functions $\phi_n: \mathbb{R} \rightarrow [0, 1]$ by

$$\phi_0(x) = \begin{cases} 0 & \text{if } x \leq 0, \\ x & \text{if } 0 \leq x \leq 1, \\ 1 & \text{if } x \geq 1, \end{cases} \quad \phi_{n+1}(x) = \begin{cases} \frac{1}{2} \cdot \phi_n(3x) & \text{if } x \leq \frac{2}{3}, \\ \frac{1}{2} + \frac{1}{2} \cdot \phi_n(3x - 2) & \text{if } x \geq \frac{1}{3}. \end{cases}$$

It is easy to check that each ϕ_n is non-decreasing, that $\phi_n(x) = 0$ for all $x \leq 0$, that $\phi_n(x) = 1$ for all $x \geq 1$, and that the two lines in the definition of ϕ_{n+1} agree in the overlap of their domains, both giving $\phi_{n+1}(x) = \frac{1}{2}$ when $\frac{1}{3} \leq x \leq \frac{2}{3}$.

Put $\phi = \lim_{n \rightarrow +\infty} \phi_n$. It is not difficult to verify that the restriction of ϕ to $[0, 1]$ is the standard Cantor function. The functions ϕ_n are polygonal approximations of ϕ .

Theorem. *The standard Cantor function is subadditive.*

Proof. The function ϕ is the pointwise limit of the functions ϕ_n as $n \rightarrow +\infty$. So to prove the subadditivity of ϕ , it suffices to prove the subadditivity of all ϕ_n , which we do by induction on n . The case $n = 0$ is trivial, so we proceed to the induction step from n to $n + 1$. Let $x, y \in \mathbb{R}$, $x \geq y$. Here we consider several cases.

Case 1: $y \leq 0$. This case is trivial as ϕ_{n+1} is monotone.

Case 2: $y \geq \frac{1}{3}$. In this case,

$$\phi_{n+1}(x + y) \leq 1 = \frac{1}{2} + \frac{1}{2} \leq \phi_{n+1}(x) + \phi_{n+1}(y).$$

Case 3: $x \leq \frac{1}{3}$. As x, y and $x + y$ are all $\leq \frac{2}{3}$, we have

$$\begin{aligned} \phi_{n+1}(x + y) &= \frac{1}{2} \cdot \phi_n(3x + 3y) \\ &\leq \frac{1}{2} \cdot \phi_n(3x) + \frac{1}{2} \cdot \phi_n(3y) = \phi_{n+1}(x) + \phi_{n+1}(y). \end{aligned}$$

Received by the editors September 2, 1994 and, in revised form, May 11, 1995.

1991 *Mathematics Subject Classification.* Primary 26D15.

Key words and phrases. Subadditive functions.

Case 4: $0 \leq y \leq \frac{1}{3} \leq x$. As $x + y \geq \frac{1}{3}$, we have

$$\begin{aligned}\phi_{n+1}(x+y) &= \frac{1}{2} + \frac{1}{2} \cdot \phi_n(3x+3y-2) \\ &\leq \frac{1}{2} + \frac{1}{2} \cdot \phi_n(3x-2) + \frac{1}{2} \cdot \phi_n(3y) = \phi_{n+1}(x) + \phi_{n+1}(y).\end{aligned}$$

These four cases exhaust all the possibilities, so the proof is complete. \square

Let us recall that a modulus of continuity is a function f defined, continuous, nondecreasing and subadditive on $[0, 1]$ with $f(0) = 0$.

Corollary. *The standard Cantor function is a modulus of continuity.*

The author wishes to thank the referee for his useful suggestions.

REFERENCES

1. Darst, R.B., *Some Cantor sets and Cantor functions*, Math. Mag. **45** (1972), 2–7. MR **46**:311
2. Matkowski, J. and Świątkowski, T., *On subadditive functions*, Proc. Amer. Math. Soc. **119** (1993), 187–197. MR **93k**:26002
3. Richey, M., *Mapping the Cantor set onto $[0, 1]$: a new construction*, preprint.
4. Rosenbaum, R.A., *Subadditive functions*, Duke Math. J. **17** (1950), 227–247. MR **12**:164a

DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY, 042 00 KOŠICE, SLOVAKIA
E-mail address: `dobos@ccsun.tuke.sk`