

COLORING FINITE SUBSETS OF UNCOUNTABLE SETS

PÉTER KOMJÁTH AND SAHARON SHELAH

(Communicated by Andreas R. Blass)

ABSTRACT. It is consistent for every $1 \leq n < \omega$ that $2^\omega = \omega_n$ and there is a function $F : [\omega_n]^{<\omega} \rightarrow \omega$ such that every finite set can be written in at most $2^n - 1$ ways as the union of two distinct monocolored sets. If GCH holds, for every such coloring there is a finite set that can be written at least $\frac{1}{2} \sum_{i=1}^n \binom{n+i}{n} \binom{n}{i}$ ways as the union of two sets with the same color.

1. INTRODUCTION

In [6] we proved that for every coloring $F : [\omega_n]^{<\omega} \rightarrow \omega$ there exists a set $A \in [\omega_n]^{<\omega}$ which can be written at least $2^n - 1$ ways as $A = H_0 \cup H_1$ for some $H_0 \neq H_1$, $F(H_0) = F(H_1)$, and that for $n = 1$ there is in fact a function F for which this is sharp. Here we show that for every $n < \omega$ it is consistent that $2^\omega = \omega_n$ and for some function F as above for every finite set A there are at most $2^n - 1$ solutions of the above equation. We use historic forcing, which was first used in [1] and [7], then in [5] and [4]. Under GCH, we improve the positive result of [6] by showing that for every F as above some finite set can be written at least $T_n = \frac{1}{2} \sum_{i=1}^n \binom{n+i}{n} \binom{n}{i}$ ways as the union of two sets with the same F value.

With the methods of [6] it is easy to show the following corollary of our independence result. It is consistent that $2^\omega = \omega_n$ and there is a function $f : \mathbb{R} \rightarrow \omega$ such that if x is a real number then x cannot be written more than $2^n - 1$ ways as the arithmetic mean of some $y \neq z$ with $f(y) = f(z)$. ((y, z) and (z, y) are not regarded as distinct.)

Notation. We use the standard set theory notation. If S is a set, κ a cardinal, then $[S]^\kappa = \{A \subseteq S : |A| = \kappa\}$, $[S]^{<\kappa} = \{A \subseteq S : |A| < \kappa\}$, $[S]^{\leq \kappa} = \{A \subseteq S : |A| \leq \kappa\}$. $P(S)$ is the power set of S . If f is a function, A a set, then $f[A] = \{f(x) : x \in A\}$.

2. THE INDEPENDENCE RESULT

Theorem 1. *For $1 \leq n < \omega$ it is consistent that $2^\omega = \omega_n$ and there is a function $F : [\omega_n]^{<\omega} \rightarrow \omega$ such that for every $A \in [\omega_n]^{<\omega}$ there are at most $2^n - 1$ solutions of $A = H_0 \cup H_1$ with $H_0 \neq H_1$, $F(H_0) = F(H_1)$.*

Received by the editors March 31, 1995 and, in revised form, May 9, 1995.

1991 *Mathematics Subject Classification.* Primary 03E35.

Key words and phrases. Axiomatic set theory, independence proofs, combinatorial set theory.

The first author was supported by the Hungarian OTKA, Grant No. T014105.

This paper is number 516 in the cumulative list of Shelah's papers. His research was supported by the Basic Research Foundation of the Israel Academy of Sciences and Humanities.

For $\alpha < \omega_n$ fix a bijection $\varphi_\alpha : \alpha \rightarrow |\alpha|$. For $x \in [\omega_n]^{<\omega}$ define $\gamma_i(x)$ for $i < k = \min(n, |x|)$ as follows: $\gamma_0(x) = \max(x)$,

$$\gamma_{i+1}(x) = \varphi_{\gamma_0(x)}^{-1} \left(\gamma_i(\varphi_{\gamma_0(x)}[x \cap \gamma_0(x)]) \right),$$

and $\gamma(x) = \{\gamma_0(x), \dots, \gamma_{k-1}(x)\}$.

So, for example, if $n = 0$ then $\gamma(x) = \emptyset$, if $n = 1$, $x \neq \emptyset$, then $\gamma(x) = \{\gamma_0(x)\} = \{\max(x)\}$.

Lemma 1. *Given $s \in [\omega_n]^{\leq n}$, there are at most countably many $x \in [\omega_n]^{<\omega}$ such that $\gamma(x) = s$.*

Proof. By induction on n . □

Let $\Phi(s) = \bigcup \{x : \gamma(x) \subseteq s\}$, a countable set for $s \in [\omega_n]^{<\omega}$.

Definition. The two sets $x, y \in [\omega_n]^{<\omega}$ are *isomorphic* if the structures $(x; <, \gamma_0(x), \dots, \gamma_{k-1}(x))$, $(y; <, \gamma_0(y), \dots, \gamma_{k-1}(y))$ are isomorphic, i.e., $|x| = |y|$ and the positions of the elements $\gamma_i(x), \gamma_i(y)$ are the same.

Notice that for every finite j there are just finitely many isomorphism types of j -element sets.

The elements of P , the applied notion of forcing, will be *some* structures of the form $p = (s, f)$ where $s \in [\omega_n]^{<\omega}$ and $f : P(s) \rightarrow \omega$.

The only element of P_0 is $\mathbf{1}_P = (\emptyset, \langle \emptyset, 0 \rangle)$; it will be the largest element of P . The elements of P_1 are of the form $p = (\{\xi\}, f)$ where $f(\emptyset) = 0 \neq f(\{\xi\})$ for $\xi < \omega_n$.

Given P_t , $p = (s, f)$ is in P_{t+1} if the following is true. $s = \Delta \cup a \cup b$ is a disjoint decomposition. $p' = (\Delta \cup a, f')$ and $p'' = (\Delta \cup b, f'')$ are in P_t , where $f' = f|P(\Delta \cup a)$, $f'' = f|P(\Delta \cup b)$. There is $\pi : \Delta \cup a \rightarrow \Delta \cup b$, an isomorphism between $(\Delta \cup a, <, P(\Delta \cup a), f')$ and $(\Delta \cup b, <, P(\Delta \cup b), f'')$. $\pi|_\Delta$ is the identity. For $H \subseteq \Delta \cup a$ the sets H and $\pi[H]$ are isomorphic. $a \cap \Phi(\Delta) = b \cap \Phi(\Delta) = \emptyset$. $f - f' - f''$ is one-to-one and takes only values outside $\text{Ran}(f')$ (which is the same as $\text{Ran}(f'')$). $P = \bigcup \{P_t : t < \omega\}$. We make $p \leq p', p''$ and the ordering on P is the one generated by this.

Lemma 2. (P, \leq) is ccc.

Proof. Assume that $p_\alpha \in P$ ($\alpha < \omega_1$). We can assume by thinning and using the Δ -system lemma and the pigeonhole principle that the following hold. $p_\alpha \in P_t$ for the same $t < \omega$. $p_\alpha = (\Delta \cup a_\alpha, f_\alpha)$ where the structures $(\Delta \cup a_\alpha, <, P(\Delta \cup a_\alpha), f_\alpha)$ and $(\Delta \cup a_\beta, <, P(\Delta \cup a_\beta), f_\beta)$ are isomorphic for $\alpha, \beta < \omega_1$, $\{\Delta, a_\alpha : \alpha < \omega_1\}$ pairwise disjoint. We can also assume that if π is the isomorphism between $(\Delta \cup a_\alpha, <, f_\alpha)$ and $(\Delta \cup a_\beta, <, f_\beta)$ then H and $\pi[H]$ are isomorphic for $H \subseteq \Delta \cup a_\alpha$. Moreover, if we assume that Δ occupies the same positions in the ordered sets $\Delta \cup a_\alpha$ ($\alpha < \omega_1$) then π will be the identity on Δ . As $\Phi(\Delta)$ is countable, by removing countably many indices we can also assume that $\Phi(\Delta) \cap a_\alpha = \emptyset$ for $\alpha < \omega_1$. Now any p_α and p_β are compatible, as we can take $p = (\Delta \cup a_\alpha \cup a_\beta, f) \leq p_\alpha, p_\beta$ where $f \supseteq f_\alpha, f_\beta$ is an appropriate extension, i.e., $f - f_\alpha - f_\beta$ is one-to-one and takes values outside $\text{Ran}(f_\alpha)$. □

Lemma 3. *If $(s, f) \in P$ and $H_0, H_1 \subseteq s$ have $f(H_0) = f(H_1)$, then H_0, H_1 are isomorphic.*

Proof. Set $(s, f) \in P_t$. We prove the statement by induction on t . There is nothing to prove for $t < 2$. Assume now that $(s, f) \in P_{t+1}$, $s = \Delta \cup a \cup b$, $\pi : \Delta \cup a \rightarrow \Delta \cup b$ as in the definition of (P, \leq) . As $f(H_0)$ is a value taken twice by f , both H_0 and H_1 must be subsets of either $\Delta \cup a$ or $\Delta \cup b$. We are done by induction unless $H_0 \subseteq \Delta \cup a$ and $H_1 \subseteq \Delta \cup b$ (or vice versa). Now H_0 and $\pi[H_0]$ are isomorphic and $f(H_0) = f(\pi[H_0]) = f(H_1)$, so by the inductive hypothesis $\pi[H_0]$ and H_1 are isomorphic, and then so are H_0, H_1 . \square

Lemma 4. *If $(s, f) \in P$, $H_0, H_1 \subseteq s$, $f(H_0) = f(H_1)$, $x \in H_0 \cap H_1$, then x occupies the same position in the ordered sets H_0, H_1 .*

Proof. Similarly to the proof of the previous lemma, by induction on t , for $(s, f) \in P_t$. With similar steps, we can assume that $(s, f) = (\Delta \cup a \cup b, f) \leq (\Delta \cup a, f'), (\Delta \cup b, f'')$, $H_0 \subseteq \Delta \cup a$, $H_1 \subseteq \Delta \cup b$. Notice that $x \in \Delta$. Now, as $\pi(x) = x$, x is a common element of $\pi[H_0]$ and H_1 , and also $f''(\pi[H_0]) = f''(H_1)$. By induction we get that x occupies the same position in $\pi[H_0]$ and H_1 , so by pulling back we get that this is true for H_0 and H_1 . \square

Lemma 5. *If $(s, f) \in P$, $A \subseteq s$, $0 \leq j \leq n$, then A can be written at most $2^j - 1$ ways as $A = H_0 \cup H_1$ with H_0, H_1 distinct, $f(H_0) = f(H_1)$, and $|\gamma(H_0) \cap \gamma(H_1)| \geq n - j$.*

Proof. By induction on j and, inside that induction, by induction on t , for $(s, f) \in P_t$. The case $t < 2$ will always be trivial.

Assume first that $j = 0$. In this case our lemma reduces to the following statement. There are no $H_0 \neq H_1$ such that $f(H_0) = f(H_1)$ and $\gamma(H_0) = \gamma(H_1)$. In the inductive argument we assume as usual that $s = \Delta \cup a \cup b$ and so $(s, f) \in P_{t+1}$ was created from $(\Delta \cup a, f')$ and $(\Delta \cup b, f'')$, $H_0 \subseteq \Delta \cup a$, $H_1 \subseteq \Delta \cup b$. As $\gamma(H_0) = \gamma(H_1)$, $\gamma(H_0) \subseteq \Delta$, but then, as $\Phi(\Delta) \cap a = \emptyset$, H_0 can have no points outside Δ and similarly for H_1 , so we can go back, say to $(\Delta \cup a, f') \in P_t$, which concludes the argument.

Assume now that the statement is proved for j and we have $p = (s, f) \in P_{t+1}$, $s = \Delta \cup a \cup b$ and p was created from $p' = (\Delta \cup a, f')$ and $p'' = (\Delta \cup b, f'')$. In $A \subseteq \Delta \cup a \cup b$ we can assume that $y = A \cap a \neq \emptyset$, $z = A \cap b \neq \emptyset$, as otherwise we can pull back to p' or p'' . But then, if $A = H_0 \cup H_1$, then, if, say, $H_0 \subseteq \Delta \cup a$, $H_1 \subseteq \Delta \cup b$ hold, then necessarily $H_0 \cap a = y$, $H_1 \cap b = z$, so $H_0 = x_0 \cup y$, $H_1 = x_1 \cup z$ where $x_0 \cup x_1 = x = A \cap \Delta$. We can create decompositions of $B = x \cup \pi[y] \cup z$ by taking $B = \pi[H_0] \cup H_1$. But some of these decompositions will not be different and it may happen that we get a non-proper (i.e., one-piece) decomposition. This can only happen if $\pi[y] = z$, and then the two decompositions $A = (x_0 \cup y) \cup (x_1 \cup z)$ and $A = (x_1 \cup y) \cup (x_0 \cup y)$ produce the same decomposition of B , namely, $B = (x_0 \cup z) \cup (x_1 \cup z)$, and there is but one decomposition, $A = (x \cup y) \cup (x \cup z)$, which cannot be mapped to a decomposition of B . If this (i.e., $\pi[y] = z$) does not happen, we are done by induction. If this does happen, we know that $\gamma(H_0) = \gamma(x_0 \cup y)$ has an element in y (by the argument at the beginning of the proof). As $f(x_0 \cup y) = f(x_1 \cup z)$, by Lemmas 3 and 4, both $H_0 = x_0 \cup y$ and $H_1 = x_1 \cup z$ have an element in the γ -subset, at the same positions which are mapped onto each other by π . We get that $\gamma(x_0 \cup z) \cap \gamma(x_1 \cup z)$ has at least $n - j$ elements, so by our inductive assumption we have at most $2^j - 1$ decompositions, which gives at most $2 \cdot (2^j - 1) + 1 = 2^{j+1} - 1$ decompositions of A . \square

Let $G \subseteq P$ be a generic subset. Set

$$S = \bigcup \{s : (s, f) \in G\}, \quad F = \bigcup \{f : (s, f) \in G\}.$$

Lemma 6. *There is a $p \in P$ such that $p \Vdash |S| = \aleph_n$.*

Proof. Otherwise $\mathbf{1} \Vdash \sup(S) < \omega_n$. By ccc, there is an ordinal $\xi < \omega_n$ for which $\mathbf{1} \Vdash \sup(S) < \xi$, but this is impossible as there are conditions in P_1 forcing that $\xi \in S$. \square

Now we can conclude the proof of the theorem. If G is generic, and $p \in G$ with the condition p of Lemma 6, then in $V[G]$ F witnesses the theorem by Lemma 5 (for $j = n$) on the ground set S . As $|S| = \omega_n$ we can replace it by ω_n . \square

3. THE GCH RESULT

Set

$$T_n = \frac{1}{2} \sum_{i=1}^n \binom{n+i}{n} \binom{n}{i}.$$

So $T_1 = 1$, $T_2 = 6$, $T_3 = 31$. In general, T_n is asymptotically $c(3 + 2\sqrt{2})^n / \sqrt{n}$ for some c .

Theorem 2 (GCH). *If $F : [\omega_n]^{<\omega} \rightarrow \omega$, then some $A \in [\omega_n]^{<\omega}$ has at least T_n decompositions as $A = H_0 \cup H_1$, $H_0 \neq H_1$, $F(H_0) = F(H_1)$.*

Proof. By the Erdős-Rado theorem (see [2], [3]) there is a set $\{x_\alpha : \alpha < \omega_1\}$ which is $(n-1)$ -end-homogeneous, i.e., for some $g : [\omega_1]^{<\omega} \rightarrow \omega$, if $\alpha_1 < \dots < \alpha_k < \beta_1 < \dots < \beta_{n-1} < \omega_1$ then

$$f(\{x_{\alpha_1}, \dots, x_{\alpha_k}, x_{\beta_1}, \dots, x_{\beta_{n-1}}\}) = g(\alpha_1, \dots, \alpha_k).$$

Select $S_1 \in [\omega_1]^{\omega_1}$ in such a way that $g(\alpha) = c_0$ for $\alpha \in S_1$. Set $\gamma_1 = \min(S_1)$. In general, if γ_i, S_i are given ($1 \leq i < n$), pick $S_{i+1} \in [S_i - (\gamma_i + 1)]^{\omega_1}$ so that $g(\gamma_1, \dots, \gamma_i, \alpha) = c_i$ for $\alpha \in S_{i+1}$, and set $\gamma_{i+1} = \min(S_{i+1})$. Given $\gamma_1, \dots, \gamma_n$ and S_n , let $\gamma_{n+1}, \dots, \gamma_{2n}$ be the n least elements of $S_n - (\gamma_n + 1)$.

Our set will be $A = \{x_{\gamma_1}, \dots, x_{\gamma_{2n}}\}$. For $0 \leq i < n$ the color of any $(n+i)$ -element subset of A containing $x_{\gamma_1}, \dots, x_{\gamma_i}$ will be c_i . We can select $\frac{1}{2} \binom{2n-i}{n} \binom{n}{i}$ different pairs of those sets which cover A . In toto, we get T_n decompositions of A . \square

REFERENCES

1. J. E. Baumgartner and S. Shelah, *Remarks on superatomic Boolean algebras*, *Annals of Pure and Applied Logic* **33** (1987), 109–129. MR **88d**:03100
2. P. Erdős, A. Hajnal, A. Máté, and R. Rado, *Combinatorial Set Theory: Partition Relations for Cardinals*, *Studies in Logic*, **106**, North-Holland, 1984. MR **87j**:04002
3. P. Erdős and R. Rado, *A partition calculus in set theory*, *Bull. Amer. Math. Soc.* **62** (1956), 427–489. MR **18**:458a
4. M. Gilchrist and S. Shelah, *On identities of colorings of pairs for \aleph_n* (to appear).
5. P. Komjáth, *A set mapping with no infinite free subsets*, *Journal of Symbolic Logic* **56** (1991), 1400–1402. MR **92i**:03054

6. P. Komjáth and S. Shelah, *On uniformly antisymmetric functions*, Real Analysis Exchange **19** (1993–1994), 218–225. MR **95b**:26001
7. S. Shelah and L. Stanley, *A theorem and some consistency results in partition calculus*, Annals of Pure and Applied Logic **36** (1987), 119–152. MR **89d**:03045

DEPARTMENT OF COMPUTER SCIENCE, EÖTVÖS UNIVERSITY, BUDAPEST, MÚZEUM KRT. 6–8,
1088, HUNGARY

E-mail address: `kope@cs.elte.hu`

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, GIVAT RAM, 91904, JERUSALEM, ISRAEL

E-mail address: `shelah@math.huji.ac.il`