

## STRONGLY $\pi$ -REGULAR RINGS HAVE STABLE RANGE ONE

PERE ARA

(Communicated by Ken Goodearl)

ABSTRACT. A ring  $R$  is said to be *strongly  $\pi$ -regular* if for every  $a \in R$  there exist a positive integer  $n$  and  $b \in R$  such that  $a^n = a^{n+1}b$ . For example, all algebraic algebras over a field are strongly  $\pi$ -regular. We prove that every strongly  $\pi$ -regular ring has stable range one. The stable range one condition is especially interesting because of Evans' Theorem, which states that a module  $M$  cancels from direct sums whenever  $\text{End}_R(M)$  has stable range one. As a consequence of our main result and Evans' Theorem, modules satisfying Fitting's Lemma cancel from direct sums.

### INTRODUCTION

Let  $R$  be a ring, associative with unity. Recall that  $R$  has *stable range one* provided that, for any  $a, b \in R$  with  $aR + bR = R$ , there exists  $y \in R$  such that  $a + by$  is invertible in  $R$ . See [17] and [18]. In this note we will prove that strongly  $\pi$ -regular rings have stable range one. As a consequence we shall obtain that modules satisfying Fitting's Lemma (over any ring) cancel from direct sums.

A ring  $R$  is said to be *strongly  $\pi$ -regular* if for each  $a \in R$  there exist a positive integer  $n$  and  $x \in R$  such that  $a^n = a^{n+1}x$ . By results of Azumaya [3] and Dischinger [8], the element  $x$  can be chosen to commute with  $a$ . In particular, this definition is left-right symmetric. Strongly  $\pi$ -regular rings were introduced by Kaplansky [12] as a common generalization of algebraic algebras and artinian rings.

In [13], Menal proved that a strongly  $\pi$ -regular ring whose primitive factor rings are artinian has stable range one. In [11], various results concerning algebraic algebras and strongly  $\pi$ -regular rings were obtained. In particular, Goodearl and Menal showed that algebraic algebras over an infinite field have stable range one [11, Theorem 3.1] (in fact they showed the somewhat stronger condition called *unit 1-stable range*), and, in [11, p.271], they conjectured that any algebraic algebra has stable range one. Our Corollary 5 proves this conjecture. Further, they ask whether all strongly  $\pi$ -regular rings have stable range one [11, p.279], proving that the answer is affirmative in several cases. For instance, the strongly  $\pi$ -regular ring

---

Received by the editors April 28, 1995.

1991 *Mathematics Subject Classification*. Primary 16E50, 16U50, 16E20.

*Key words and phrases*. Strongly  $\pi$ -regular ring, stable range one, exchange ring, Fitting's Lemma.

The author was partially supported by DGYCIT grant PB92-0586 and the Comissionat per Universitats i Recerca de la Generalitat de Catalunya.

$R$  has stable range one when either  $R$  is a von Neumann regular ring [11, Theorem 5.8] or every element of  $R$  is a sum of a unit plus a central unit [11, Corollary 6.2].

More recently, Yu [20],[21] and Camillo and Yu [5] proved that strongly  $\pi$ -regular rings have stable range one under some additional hypothesis. For example they show that a strongly  $\pi$ -regular ring such that every power of a regular element is regular, has stable range one [5, Theorem 5], generalizing [11, Theorem 5.8].

Goodearl and Menal also proved that a strongly  $\pi$ -regular ring has stable range one if and only if every nilpotent regular element of any corner of the ring is unit-regular in that corner [11, Theorem 6.1]. We will prove that in fact every nilpotent regular element of every *exchange ring* is unit-regular.

Let  $R$  be any ring. An element  $a \in R$  is said to be *regular* if there exists  $b \in R$  such that  $a = aba$ . It is easy to see that  $a$  is regular if and only if the right annihilator of  $a$  (“the kernel of  $a$ ”) and the right ideal generated by  $a$  (“the image of  $a$ ”) are both direct summands of  $R_R$ . The element  $a \in R$  is said to be *unit-regular* if there exists an invertible element  $u \in R$  such that  $a = aua$ . It is easy to see that  $a$  is unit-regular if and only if  $a$  is regular and  $\text{rann}(a) \cong E$  as right  $R$ -modules, where  $E$  denotes a complement of  $aR$  in  $R_R$  (“the kernel of  $a$ ” is isomorphic to “the cokernel of  $a$ ”).

An element  $a \in R$  is said to be *strongly  $\pi$ -regular* if there exists a positive integer  $n$  and  $b \in R$  such that  $a^n = a^{n+1}b$  and  $ab = ba$ . The ring  $R$  is said to be *strongly  $\pi$ -regular* if every element of  $R$  is strongly  $\pi$ -regular. By combining results of Dischinger [8] and Azumaya [3], one obtains the characterization of strongly  $\pi$ -regular rings as either the left  $\pi$ -regular rings or the right  $\pi$ -regular rings; see [11, p. 300] or [5, Lemma 6].

A right  $R$ -module  $M$  has the *exchange property* (see [7]) if for every module  $A_R$  and any decompositions

$$A = M' \oplus N = \bigoplus_{i \in I} A_i$$

with  $M' \cong M$ , there exist submodules  $A'_i \subseteq A_i$  such that

$$A = M' \oplus \left( \bigoplus_{i \in I} A'_i \right).$$

$M$  has the *finite exchange property* if the above condition is satisfied whenever the index set  $I$  is finite. Clearly a finitely generated module satisfies the exchange property if and only if it satisfies the finite exchange property.

Following [19], we say that a ring  $R$  is an *exchange ring* if  $R_R$  satisfies the (finite) exchange property. By [19, Corollary 2], this definition is left-right symmetric.

Every strongly  $\pi$ -regular ring is an exchange ring [16, Example 2.3]. A great deal is known about strongly  $\pi$ -regular rings and exchange rings; see for example [4], [15], [16], [20] and [1].

## THE RESULTS

The following technical lemma is the key to obtain our main results.

**Lemma 1.** *Let  $R$  be an exchange ring and let  $a$  be a regular element of  $R$ . Let  $K$  denote the right annihilator of  $a$ , and  $E$  be a complement of  $aR$  in  $R_R$ . Then there exist right ideals  $A_i, A'_i, B_i, B'_i, C_i, C'_i$ , for  $i \geq 1$ , such that the following conditions are satisfied:*

- (1)  $R = K \oplus \left( \bigoplus_{j=1}^i (A_j \oplus B_j) \right) \oplus C_i$  for all  $i \geq 1$ . Hence,  $C_i \cong A_{i+1} \oplus B_{i+1} \oplus C_{i+1}$ .

- (2)  $E \cong (A_i \oplus B_i) \oplus (A'_i \oplus B'_i)$  for all  $i \geq 1$ .
- (3)  $K \cong A'_i \oplus B'_i \oplus C'_i$  for all  $i \geq 1$ .
- (4)  $A'_i \oplus B'_i = A_{i+1} \oplus A'_{i+1}$  for all  $i \geq 1$ .
- (5)  $aR = C_1 \oplus C'_1$  and, for  $i \geq 1$ ,  $aA_i \oplus aB_i = B_{i+1} \oplus B'_{i+1}$  and  $aC_i = C_{i+1} \oplus C'_{i+1}$ .

Hence,  $C_{i+1} \subseteq a^{i+1}R$ .

*Proof.* Write  $R = K \oplus L = E \oplus aR$ . By using the exchange property we obtain  $A_1, C_1, A'_1, C'_1$  such that  $R = K \oplus A_1 \oplus C_1$ , and  $E = A_1 \oplus A'_1$ , and  $aR = C_1 \oplus C'_1$ . Note that  $K \oplus A_1 \oplus C_1 = A'_1 \oplus C'_1 \oplus A_1 \oplus C_1$ , and so  $K \cong A'_1 \oplus C'_1$ . Set  $B_1 = B'_1 = 0$ .

Now assume that, for some  $n \geq 1$ , we have constructed right ideals  $A_i, A'_i, B_i, B'_i, C_i, C'_i$ , with  $i \leq n$ , satisfying the desired conditions. We will construct  $A_{n+1}, A'_{n+1}, B_{n+1}, B'_{n+1}, C_{n+1}, C'_{n+1}$ . Using (4) repeatedly and the fact that  $B'_1 = 0$ , we obtain

$$(6) \quad A'_1 \oplus \left(\bigoplus_{i=1}^{n-1} B'_{i+1}\right) = \left(\bigoplus_{i=2}^n A_i\right) \oplus A'_n \oplus B'_n.$$

From (1) we have  $aR = \left(\bigoplus_{i=1}^n (aA_i \oplus aB_i)\right) \oplus aC_n$ . By using this and relations (5) and (6) we obtain

$$\begin{aligned} R &= E \oplus aR = A_1 \oplus A'_1 \oplus \left(\bigoplus_{i=1}^n (aA_i \oplus aB_i)\right) \oplus aC_n \\ &= (A_1 \oplus B_1) \oplus A'_1 \oplus \left(\bigoplus_{i=1}^{n-1} (B_{i+1} \oplus B'_{i+1})\right) \oplus aA_n \oplus aB_n \oplus aC_n \\ &= A_1 \oplus \left(\bigoplus_{i=1}^n B_i\right) \oplus \left(A'_1 \oplus \left(\bigoplus_{i=1}^{n-1} B'_{i+1}\right)\right) \oplus aA_n \oplus aB_n \oplus aC_n \\ &= A_1 \oplus \left(\bigoplus_{i=1}^n B_i\right) \oplus \left(\bigoplus_{i=2}^n A_i\right) \oplus (A'_n \oplus B'_n) \oplus aA_n \oplus aB_n \oplus aC_n \\ &= \left(\bigoplus_{i=1}^n (A_i \oplus B_i)\right) \oplus (A'_n \oplus B'_n) \oplus aA_n \oplus aB_n \oplus aC_n. \end{aligned}$$

Now applying the exchange property to the decompositions

$$\begin{aligned} R &= K \oplus \left(\bigoplus_{i=1}^n (A_i \oplus B_i)\right) \oplus C_n \\ &= \left(\bigoplus_{i=1}^n (A_i \oplus B_i)\right) \oplus (A'_n \oplus B'_n) \oplus (aA_n \oplus aB_n) \oplus aC_n, \end{aligned}$$

we obtain a decomposition

$$R = K \oplus \left(\bigoplus_{i=1}^n (A_i \oplus B_i)\right) \oplus A_{n+1} \oplus B_{n+1} \oplus C_{n+1}$$

such that

$$A_{n+1} \oplus A'_{n+1} = A'_n \oplus B'_n$$

for some right ideal  $A'_{n+1}$ , while

$$B_{n+1} \oplus B'_{n+1} = aA_n \oplus aB_n$$

for some right ideal  $B'_{n+1}$ , and

$$C_{n+1} \oplus C'_{n+1} = aC_n$$

for some right ideal  $C'_{n+1}$ . So we obtain (1), (4) and (5).

Since  $(\bigoplus_{i=1}^{n+1} (A_i \oplus B_i)) \oplus C_{n+1}$  is a common complement of both  $K$  and  $A'_{n+1} \oplus B'_{n+1} \oplus C'_{n+1}$ , we obtain (3).

Now we will prove (2). We have

$$\begin{aligned} E &\cong (A_n \oplus B_n) \oplus (A'_n \oplus B'_n) \\ &\cong aA_n \oplus aB_n \oplus A_{n+1} \oplus A'_{n+1} \\ &= B_{n+1} \oplus B'_{n+1} \oplus A_{n+1} \oplus A'_{n+1} \\ &= (A_{n+1} \oplus B_{n+1}) \oplus (A'_{n+1} \oplus B'_{n+1}). \end{aligned}$$

This completes the inductive step.  $\square$

**Theorem 2.** *Let  $R$  be an exchange ring and let  $a$  be a nilpotent regular element of  $R$ . Then  $a$  is unit-regular.*

*Proof.* Assume that  $a^{n+2} = 0$  for some  $n \geq 0$ . Let  $A_i, A'_i, B_i, B'_i, C_i, C'_i$  be right ideals as in Lemma 1. Then  $C_{n+1} \subseteq K$  by (5) of Lemma 1, and so  $C_{n+1} = 0$  by (1). By (5), we have  $C'_{n+1} = aC_n \cong C_n$  so that, using (1), we obtain  $C'_{n+1} \cong C_n \cong A_{n+1} \oplus B_{n+1} \oplus C_{n+1} = A_{n+1} \oplus B_{n+1}$ . Now using this fact and (2), (3), we have

$$\begin{aligned} K &\cong A'_{n+1} \oplus B'_{n+1} \oplus C'_{n+1} \\ &\cong (A'_{n+1} \oplus B'_{n+1}) \oplus (A_{n+1} \oplus B_{n+1}) \cong E. \end{aligned}$$

We conclude that  $a$  is unit-regular.  $\square$

**Theorem 3.** *Let  $R$  be an exchange ring and let  $a$  be a regular element of  $R$ . If  $a$  is strongly  $\pi$ -regular, then  $a$  is unit-regular.*

*Proof.* Let  $a$  be a regular, strongly  $\pi$ -regular element of  $R$ . Let  $b \in R$  be such that  $a^n = a^{n+1}b$  for some  $n \geq 1$ , and  $ab = ba$ . Set  $e = a^n b^n$  and note that  $e$  is idempotent. Moreover,  $ea = ae$  is invertible in  $eRe$ , with inverse  $a^n b^{n+1}$ , and  $a(1-e) = (1-e)a \in (1-e)R(1-e)$  is a regular nilpotent element with  $(a(1-e))^n = 0$ . Since  $(1-e)R(1-e)$  is an exchange ring [19, Theorem 2], it follows from Theorem 2 that  $a(1-e)$  is unit-regular in  $(1-e)R(1-e)$ . Consequently,  $a$  is unit-regular in  $R$ .  $\square$

**Theorem 4.** *Strongly  $\pi$ -regular rings have stable range one.*

*Proof.* By [16, Example 2.3], any strongly  $\pi$ -regular ring is an exchange ring. So the result follows from Theorem 3 and [5, Theorem 3]. Alternatively, one can use Theorem 2 and [11, Theorem 6.1].  $\square$

Our next result proves the conjecture made by Goodearl and Menal in [11, p.271].

**Corollary 5.** *Any algebraic algebra over a field has stable range one.*

*Proof.* Clearly, an algebraic algebra over a field is strongly  $\pi$ -regular. So, the result follows from Theorem 4.  $\square$

A module  $M$  is said to *satisfy Fitting's Lemma* if for each  $f \in \text{End}_R(M)$  there exists an integer  $n \geq 1$  such that  $M = \text{Ker}(f^n) \oplus f^n(M)$ . By [2, Proposition 2.3],  $M$  satisfies Fitting's Lemma if and only if  $\text{End}_R(M)$  is strongly  $\pi$ -regular. It was proved in [6] that modules satisfying Fitting's Lemma have power cancellation. Theorem 4 enables us to improve this result, as follows.

**Corollary 6.** *Let  $M$  be a module satisfying Fitting's Lemma. Then  $M$  cancels from direct sums.*

*Proof.* By [2, Proposition 2.3],  $E := \text{End}_R(M)$  is a strongly  $\pi$ -regular ring. Now, Theorem 4 gives that the stable range of  $E$  is one and, by Evans' Theorem [9, Theorem 2],  $M$  cancels from direct sums.  $\square$

Now we will obtain some cancellation results for finitely generated modules over certain strongly  $\pi$ -regular rings. We need the following concept, introduced by Goodearl [10].

**Definition** ([10]). An element  $u$  of a ring  $R$  is said to be *right repetitive* provided that for each finitely generated right ideal  $I$  of  $R$ , the right ideal  $\sum_{n=0}^{\infty} u^n I$  is finitely generated. The ring  $R$  is *right repetitive* if each element of  $R$  is right repetitive.

Note that every algebraic algebra is (right and left) repetitive.

**Corollary 7.** *Let  $R$  be a strongly  $\pi$ -regular, right repetitive ring. Then any cyclic right  $R$ -module cancels from direct sums.*

*Proof.* Apply [14, Theorem 19], Theorem 4 and Evans' Theorem [9, Theorem 2].  $\square$

**Corollary 8.** *If  $R$  is a ring such that all matrix rings  $M_n(R)$  are strongly  $\pi$ -regular and right repetitive, then any finitely generated right  $R$ -module cancels from direct sums.*

*Proof.* Apply [10, Theorem 8], Theorem 4 and Evans' Theorem [9, Theorem 2].  $\square$

#### ACKNOWLEDGEMENTS

It is a pleasure to thank Ferran Cedó and Ken Goodearl for their useful suggestions.

#### REFERENCES

1. P. Ara, K.R. Goodearl, K. O'Meara and E. Pardo, *Separative cancellation for projective modules over exchange rings*, Preprint.
2. E.P. Armendariz, J.W. Fisher and R.L. Snider, *On injective and surjective endomorphisms of finitely generated modules*, Communications in Algebra **6**(7) (1978), 659-672. MR **57**:9754
3. G. Azumaya, *Strongly  $\pi$ -regular rings*, J. Fac. Sci. Hokkaido Univ. **13** (1954), 34-39. MR **16**:788
4. W.D. Burgess and P. Menal, *On strongly  $\pi$ -regular rings and homomorphisms into them*, Communications in Algebra **16**(8) (1988), 1701-1725. MR **89f**:16015
5. V.P. Camillo and Hua-Ping Yu, *Stable range one for rings with many idempotents*, Trans. Amer. Math. Soc. **347** (1995), 3141-3147. CMP 95:12
6. R. Camps and P. Menal, *Power cancellation for artinian modules*, Communications in Algebra **19**(7) (1991), 2081-2095. MR **92m**:16006
7. P. Crawley and B. Jónsson, *Refinements for infinite direct decompositions of algebraic systems*, Pacific J. Math. **14** (1964), 797-855. MR **30**:49
8. M.F. Dischinger, *Sur les anneaux fortement  $\pi$ -réguliers*, C.R. Acad. Sci. Paris, Ser. A **283** (1976), 571-573. MR **54**:10321

9. E.G. Evans, *Krull-Schmidt and cancellation over local rings*, Pacific J. Math **46** (1973), 115-121. MR **48**:2170
10. K.R. Goodearl, *Surjective endomorphisms of finitely generated modules*, Communications in Algebra **15**(3) (1987), 589-609. MR **88d**:16010
11. K.R. Goodearl and P. Menal, *Stable range one for rings with many units*, J. Pure Applied Algebra **54** (1988), 261-287. MR **89h**:16011
12. I. Kaplansky, *Topological representations of algebras II*, Trans. Amer. Math. Soc. **68** (1950), 62-75. MR **11**:317
13. P. Menal, *On  $\pi$ -regular rings whose primitive factor rings are artinian*, J. Pure Applied Algebra **20** (1981), 71-78. MR **81k**:16015
14. P. Menal, *Cancellation modules over regular rings*, in "Ring Theory, Proceedings, Granada 1986", J.L. Bueso, P. Jara and B. Torrecillas (eds.), LNM **1328**, Springer-Verlag, 1988, pp. 187-209. MR **89i**:16010
15. W.K. Nicholson, *Lifting idempotents and exchange rings*, Trans. Amer. Math. Soc. **229** (1977), 269-278. MR **55**:12757
16. J. Stock, *On rings whose projective modules have the exchange property*, J. Algebra **103** (1986), 437-453. MR **88e**:16038
17. L.N. Vaserstein, *Stable rank of rings and dimensionality of topological spaces*, Funct. Ana. Appl. **5** (1971), 102-110. MR **44**:1701
18. L.N. Vaserstein, *Bass's first stable range condition*, J. Pure Applied Algebra **34** (1984), 319-330. MR **86c**:18009
19. R.B. Warfield, Jr., *Exchange rings and decompositions of modules*, Math. Ann. **199** (1972), 31-36. MR **48**:11218
20. Hua-Ping Yu, *Stable range one for exchange rings*, J. Pure Applied Algebra **98** (1995), 105-109.
21. Hua-Ping Yu, *On strongly  $\pi$ -regular rings of stable range one*, Bull. Austral. Math. Soc. **51** (1995), 433-437. CMP 95:12

DEPARTAMENT DE MATEMÀTIQUES, EDIFICI CC, UNIVERSITAT AUTÒNOMA DE BARCELONA,  
08193 BELLATERRA, BARCELONA, SPAIN  
E-mail address: para@mat.uab.es