

## A NOTE ON FISCHER-MARSDEN'S CONJECTURE

YING SHEN

(Communicated by Peter Li)

ABSTRACT. In this paper, we borrowed some ideas from general relativity and find a Robinson-type identity for the overdetermined system of partial differential equations in the Fischer-Marsden conjecture. We proved that if there is a nontrivial solution for such an overdetermined system on a 3-dimensional, closed manifold with positive scalar curvature, then the manifold contains a totally geodesic 2-sphere.

Let  $\mathcal{M}$  denote the set of smooth Riemannian metrics on an  $n$ -dimensional closed manifold  $M$  whose derivatives are  $L_2$ -integrable. Then for any  $g \in \mathcal{M}$ , its scalar curvature  $R_g$  is an element in the space  $\mathcal{W}$  of  $C^\infty$  functions. From the formula for  $R_g$  in local coordinates, we see that the scalar curvature map from  $\mathcal{M}$  to  $\mathcal{W}$  defines a quasi-linear differential operator of second order. The derivative  $R'_g$  at  $g \in \mathcal{M}$  is given by

$$(1) \quad R'_g(h) = -\Delta_g(\operatorname{tr}_g h) + \delta_g^* \delta_g(h) - g(\operatorname{Ric}_g, h)$$

where  $\delta$  is the divergence operator on the symmetric  $p$ -tensor on  $M$ ,  $\operatorname{Ric}_g$  is the Ricci curvature tensor of  $g$ ,  $\Delta$  is the Laplacian, and  $\delta^*$  is the formal adjoint of  $\delta$ . In the Riemannian case, if  $\{e_i\}_{i=1}^n$  is a local orthonormal basis of vector fields, then

$$(\delta\alpha)(e_1, \dots, e_p) = -\sum_{i=1}^n (D_{e_i}\alpha)(e_i, e_1, \dots, e_p).$$

In particular, for any one-form  $\alpha$ , we have

$$(2) \quad \delta^*\alpha = \frac{1}{2}L_{\hat{\alpha}}g$$

where  $L_{\hat{\alpha}}$  is the Lie derivative of the vector field  $\hat{\alpha}$  and  $\hat{\alpha}$  is the dual of  $\alpha$ .

It is easy to compute that the  $L_2$ -adjoint operator of  $R'_g$  is

$$(3) \quad R_g'^*(f) = -(\Delta f)g + D_g d(f) - f\operatorname{Ric}_g.$$

Since we can regard  $R'_g$  as a linear map from the space of symmetric tensors to the space of functions, so it is known that the following decompositions are true [Be-Eb]:

$$(\operatorname{Im}(R'_g))^\perp = \ker(R_g'^*)$$

or

$$(4) \quad \mathcal{W} = \operatorname{Im}(R'_g) \oplus \ker(R_g'^*).$$

---

Received by the editors June 12, 1995.

1991 *Mathematics Subject Classification*. Primary 53C21, 53C42.

**Definition 1.** We say that the metric  $g$  is singular if  $R'_g$  is not a surjective map.

A natural question is: *What can we say about the singular metric  $g$  on a closed manifold  $M$ ?*

From the decomposition (4), we know that  $g$  being singular is the same as the existence of a nontrivial solution to the following equation:

$$(5) \quad D_g d(v) = v Ric_g + (\Delta v)g.$$

Now we see that equation (5) is an overdetermined system, so the existence of a nontrivial solution should provide us with some information on the geometry of the underlying manifold  $M$ . Actually, Bourguignon [Bo] and Fischer-Marsden [F-M] proved the following:

**Lemma 1.** *Let  $(M, g)$  be a closed Riemannian manifold admitting a nontrivial solution to equation (5). Then either  $(M, g)$  is Ricci-flat and  $Ker(R'_g)^* = \mathcal{W}$ , or  $R_g$  is a positive constant.*

Based on this lemma, Fischer-Marsden made the following conjecture:

**Conjecture 1.**  $(M, g)$  must be an Einstein manifold if  $g$  is a singular metric.

We can easily rewrite (5) as:

$$(6) \quad D_g dv + \frac{R_g}{n(n-1)}vg = vT_g$$

or

$$(7) \quad v_{ij} + v \frac{R_g}{n(n-1)}g_{ij} = vT_{ij}$$

where  $T_g$  and  $T_{ij}$  are the traceless-Ricci tensors.

In order to prove the Conjecture, one only needs to show that  $(M, g)$  is Einstein if  $R_g$  is a positive constant. Now if  $(M, g)$  is an Einstein manifold with positive scalar curvature, then a theorem of Obata [Ob] tells us that  $(M, g)$  must be a round sphere. So the Fischer-Marsden Conjecture can be rephrased as the following:

**Conjecture 2.** If  $g$  is a singular metric on a Riemannian manifold  $M$  such that its scalar curvature is positive, then the existence of a nontrivial solution of the overdetermined system (5) implies that  $(M, g)$  is a round sphere.

The counterexample of the Fischer-Marsden Conjecture was provided by O. Kobayashi [Ko] and J. Lafontaine [Laf] independently. They proved that if we assume further that  $(M, g)$  is conformally flat, then  $(M, g)$  must be isometric to one of the following:

- (a) Euclidean sphere  $S^n$ .
- (b) Finite quotient of  $(S^1, dt^2) \times (S^{n-1}, g_0)$ ,  $g_0$  is the canonical metric.
- (c) Finite quotient of a product torus  $(S^1 \times S^{n-1}, dt^2 + h^2(t)g_0)$ .

By looking at the examples listed above, one observes that  $(M, g)$  always contains a totally geodesic  $(n-1)$ -sphere if  $(M, g)$  is conformally flat. So it is interesting to know whether this is a general phenomenon without assuming that  $(M, g)$  is conformally flat. The present note gives a positive answer to the question if  $\dim M = 3$ .

**Theorem 1.** *If  $g$  is a singular metric on a 3-dimensional closed Riemannian manifold  $M$  such that its scalar curvature  $R_g$  is positive, then  $(M, g)$  contains a totally geodesic 2-sphere.*

The idea of the proof is borrowed from general relativity.

Recall that a geodesically complete spacetime  $(N, \overset{4}{g})$  is called static if and only if there is an orientable, spacelike, 3-manifold  $(\Sigma, g)$  such that  $N$  is diffeomorphic to  $\Sigma \times \mathbb{R}$ , and

$$\overset{4}{g} = -U^2 dt^2 + g$$

where  $U$  is called the gravitational scalar potential of the static field.

The Einstein equations for a static perfect fluid with mass-energy density  $\rho$  and pressure  $p$  can be decomposed into:

$$(8) \quad Ric(g)_{ij} = U^{-1}U_{;ij} + 4\pi(\rho - p)g_{ij},$$

$$(9) \quad \Delta U = 4\pi U(\rho + 3p).$$

Therefore, we know that our equation (5) is the equation for static perfect fluid. There has been active research among relativists for the proof of the spherical symmetries for stellar models. We only provide the interested readers with a short list of papers: [Rob], [B-Mas], [I], [K], [Mas], [Lind]. The essential step in proving the spherical symmetry is to find some nice Robinson type identity that generally has the form that “divergence equals positive quantity”.

*Proof of the Theorem.* We are going to construct a Robinson type identity first and then prove the theorem.

We may assume, without loss of generality, that  $R_g = n(n - 1)$ . Hence equation (6) becomes

$$(10) \quad D_g dv + vg = vT_g$$

and

$$(11) \quad \Delta v + nv = 0.$$

It is easy to see that  $\{x \in M | v(x) \neq 0\}$  is dense in  $M$  (see [Bo] or [F-M]). Let  $M_0$  be a connected component of  $\{x \in M | v(x) > 0\}$  such that  $dv \neq 0$  on  $\Sigma = \partial M_0$ .

Define  $W = |\nabla v|^2$  and  $W_0 = 1 - v^2$ . Since the scalar curvature is constant on  $\partial M_0$ , we can construct the Robinson type identity for the system (10):

$$(12) \quad \nabla^a [v^{-1} \nabla_a (W - W_0)] = 2v|T|^2.$$

Equation (12) can be easily verified by straightforward computation.

From equation (10), it is easy to check that

$$|W| = constant$$

on  $\Sigma$ . When restricted to  $\Sigma$ , we know that  $v_{;j} = 0$ . It is straightforward to check that  $\Sigma$  is totally geodesic and  $-\frac{\nabla v}{W^{\frac{1}{2}}}$  is a unit outward normal vector field to  $\Sigma$ . We let  $e_1, e_2, e_3$  be an orthonormal basis such that along  $\Sigma$ ,

$$e_3 = \frac{\nabla v}{W^{\frac{1}{2}}}.$$

We remark that although  $\{e_1, e_2, e_3\}$  is a local frame,  $e_3$  is globally defined on  $\Sigma$  due to the embeddedness of  $\Sigma$ . The Gauss equation tells us that

$$(13) \quad R_{1212} = K$$

where  $R_{ijkl}$  is the full curvature tensor of  $(M, g)$  and  $K$  is the Gaussian curvature of  $\Sigma$  with the induced metric.

Integrating equation (12) and using Stoke's formula, we obtain

$$(14) \quad \int_{\Sigma} W^{-\frac{1}{2}} R^{ab} v_{;a} v_{;b} - 2W^{\frac{1}{2}} S_0 = - \int_{M_0} v |T|^2$$

where  $S_0$  is the area of  $\Sigma$ .

By using the facts

$$(15) \quad R_{11} = R_{1313} + K,$$

$$(16) \quad R_{22} = R_{2323} + K,$$

$$(17) \quad R_{33} = R_{1313} + R_{2323},$$

$$(18) \quad R_{11} + R_{22} + R_{33} = 6,$$

we know that

$$(19) \quad \int_{\Sigma} W^{-\frac{1}{2}} R^{ab} v_{;a} v_{;b} = 3W^{\frac{1}{2}} S_0 - W^{\frac{1}{2}} \int_{\Sigma} K.$$

Therefore we get

$$(20) \quad \int_{\Sigma} K \geq S_0 > 0,$$

and the Gauss-Bonnet theorem immediately implies that  $\Sigma$  is a 2-sphere.

*Remark.* The area of the sphere  $\Sigma$ ,  $S_0$ , is in general less than or equal to  $4\pi$ . In case  $S_0 = 4\pi$ , then it is easy to see that  $(M_0, g)$  is isometric to the canonical 3-sphere since the manifold is Einstein.

#### ACKNOWLEDGEMENTS

We would like to thank Professor Richard Schoen for bringing the problem to our attention and for his stimulating discussion and constant encouragement. We would also like to thank H.-D. Cao, M. Gursky, D. Pollack, and P. Yang for their interest in the problem and their encouragement of writing this up. Finally, we wish to thank the referee for many helpful comments and suggestions.

#### REFERENCES

- [Be-Eb] M. Berger and D. Ebin, *Some decompositions of the space of symmetric tensors on a Riemannian manifold*, J. Diff. Geom. **3** (1969), 379–392. MR **42**:993
- [Bes] A. Besse, *Einstein Manifolds*, Springer-Verlag, Berlin, 1987. MR **88f**:53087
- [Bo] J.P. Bourguignon, *Une stratification de l'espace des structures riemanniennes*, Compositio Math. **30** (1975), 1–41. MR **54**:6189
- [B-Mas] G. L. Bunting and A.K.M. Masood-ul-Alam, *Nonexistence of Multiple Black Holes in Asymptotically Euclidean Static Vacuum Space-Time*, General Relativity and Gravitation **19** (1987), 147–154. MR **88e**:83031
- [F-M] A.E. Fischer and J.E. Marsden, *Linearization stability of nonlinear partial differential equations*, Proc. Symp. Pure Math. **27** (1975), 219–262. MR **53**:4337
- [I] W. Israel, *Event horizons in static vacuum space-times*, Phys. Rev. **164** (1967), 1776–1779.
- [Ko] O. Kobayashi, *A differential equation arising from scalar curvature function*, J. Math. Soc. Japan **34**, No. **4** (1982), 665–675. MR **84a**:53046
- [K] J.P. Kunzle, *On the Spherical Symmetry of a Static Perfect Fluid*, Comm. Math. Phys. **20** (1971), 85–100. MR **43**:1586
- [Laf] J. Lafontaine, *Sur la geometrie d'une generalisation de l'equation d'Obata*, J. Math. Pures Appliquees **62** (1983), 63–72. MR **84i**:53047

- [Lind] L. Lindblom, *Some properties of static general relativistic stellar models*, J. Math. Phys. **21**, No. **6** (1980), 1455–1459.
- [Mas] A.K. M. Masood-ul-Alam, *On spherical symmetry of static perfect fluid spacetimes and positive-mass theorem*, Class. Quantum Grav. **4** (1987), 625–633. MR **88f**:83054
- [Ob] M. Obata, *Certain conditions for a riemannian manifold to be isometric with a sphere*, J. Math. Soc. Japan **14** (1962), 333–340. MR **25**:5479
- [Rob] D. C. Robinson, *A simple proof of the generalization of Israel's Theorem*, General Relativity and Gravitation **8**, No. **8** (1977), 695–698.

DEPARTMENT OF MATHEMATICS, TEXAS A & M UNIVERSITY, COLLEGE STATION, TEXAS 77843

*E-mail address*: `yshen@math.tamu.edu`

*Current address*: Department of Mathematics, Dartmouth College, Hanover, New Hampshire 03755

*E-mail address*: `ying.shen@dartmouth.edu`