

CONSTRUCTION OF ANR TOPOLOGIES ON CERTAIN GROUPS

BERND GÜNTHER

(Communicated by Thomas Goodwillie)

ABSTRACT. A method is shown to construct ANR-topologies on topological groups of suitable homotopy type.

It is well known that every simple countable CW-complex with vanishing Postnikov invariants has the homotopy type of a topological Abelian group G , that can be taken as realization of a countable simplicial Abelian group and hence is a countable CW-complex [5, Ch.V]. In particular this means that the homotopy groups $\pi_n(G)$ can be chosen as an arbitrarily prescribed sequence of countable Abelian groups. However, there are situations where it would be preferable to have G as an ANR-space instead of a CW-complex. One would guess that a suitable topology exists on G , because by subdivision it can be turned into a polyhedron and then can get the metric topology, but the subdivision process necessarily leaves the range of simplicial Abelian groups and need not produce a group topology. Instead, we follow a method developed by Cauty in [1].

Theorem. *Let G be a locally contractible, not necessarily Abelian group, every open subset of which is σ -compact, such that G is the union of a sequence of finite dimensional compact metric spaces and has the homotopy type of a CW-complex. Then G carries a metrizable group topology, coarser than the original one but of the same homotopy type, which turns G into an ANR-space.*

All these requirements are satisfied, for instance, if G itself is a countable CW-complex.

Proof. We construct a decreasing sequence of open neighborhoods V_n of the identity $1 \in G$ such that

1. $\bigcap_{n=1}^{\infty} V_n = \{1\}$,
2. the inclusion map $V_{n+1} \hookrightarrow V_n$ is null homotopic,
3. if $H_m : V_{m+1} \times I \rightarrow V_m$ is the null homotopy from 2, then for any $g \in V_{m+1}$ and each $n \geq m$ there exist $N \geq n$ and $\varepsilon > 0$ such that $g' \in gV_N$, $|t' - t| < \varepsilon \Rightarrow H_m(g', t') \in H_m(g, t)V_n$,
4. $V_n^{-1} = V_n$ and $V_{n+1}V_{n+1} \subseteq V_n$,
5. for each n and every $g \in G$ there exists $N \geq n$ with $g^{-1}V_Ng \subseteq V_n$.

Received by the editors April 1, 1996.

1991 *Mathematics Subject Classification.* Primary 54H11, 54C55, 22A05.

Key words and phrases. Homotopy type of topological groups, ANR-spaces, CW-complexes.

All of these properties are immediately clear with the exception of 3 and 5. We construct V_n inductively. Let U_n be a sequence of neighborhoods of the identity with $\bigcap_{n=1}^{\infty} U_n = \{1\}$; it exists because the open set $G \setminus \{1\}$ is σ -compact and therefore $\{1\}$ is a G_δ . Choose V_1 with $V_1 = V_1^{-1} \subseteq U_1$, otherwise arbitrarily. Now suppose that V_1, \dots, V_k are already constructed such that 2, 4, $V_k \subseteq U_k$ and the following properties hold:

- 3a. For $k < m \leq n$ there exist a sequence of neighborhoods of the identity $\{W_i^{km} \mid i \in \mathbb{N}\}$ and a sequence of numbers $\varepsilon_i^{km} > 0$ such that for each $g \in V_{k+1}$ there exists i with $g' \in gW_i^{km}$, $|t' - t| < \varepsilon_i^{km} \Rightarrow H_k(g', t') \in H_k(g, t)V_m$.
- 3b. $V_n \subseteq \bigcap_{k < m < n, i \leq n} W_i^{km}$.
- 5a. For every $g \in K_n$ we have $g^{-1}V_{n+1}g \subseteq V_n$, where K_m is an increasing sequence of compacta covering G .

To pass from n to $n + 1$ choose V_{n+1} subject to conditions 2, 3b, 4, 5a and $V_{n+1} \subseteq V_n \cap U_{n+1}$. It now remains to construct the neighborhoods $W_i^{k,n+1}$ for $k \leq n$. By assumption we can find a sequence of compact sets $C_i \subseteq V_{k+1}$ with $\bigcup_{i=1}^{\infty} C_i = V_{k+1}$. For each of them we choose a neighborhood of the identity $W_i^{k,n+1}$ with $C_i W_i^{k,n+1} \subseteq V_{k+1}$ and a number $\varepsilon_i^{k,n+1} > 0$ such that $g \in C_i$, $g' \in gW_i^{k,n+1}$ and $|t' - t| < \varepsilon_i^{k,n+1}$ implies $H_k(g', t') \in H_k(g, t)V_m$. It is clear that the whole sequence of neighborhoods V_n satisfies 1–5 and hence defines a metrizable group topology on G , which is coarser than the original topology. Denote by \tilde{G} the group G topologized by our neighborhood system V_n . 3 ensures that the homotopies H_n are continuous with respect to our metrizable topology, and hence that \tilde{G} is locally contractible. It now follows from [3] that \tilde{G} is an ANR-space and from [4, Theorem 1] that the identity map $G \rightarrow \tilde{G}$ is a homotopy equivalence. \square

In a similar way it can be proved that the construction is “almost” functorial in the sense that for any family of continuous homomorphisms $f_\lambda : G_\lambda \rightarrow H_\lambda$ between groups satisfying the assumptions of our theorem, such that each group appears at most countably often among the source groups G_λ the ANR group topology can be chosen in such a way that all homomorphisms $f_\lambda : \tilde{G}_\lambda \rightarrow \tilde{H}_\lambda$ are continuous. The countability assumption seems to be essential.

REFERENCES

1. R. Cauty: *Un espace métrique linéaire qui n'est pas un rétracte absolu*; Fund. Math. 146 (1994) 85–99. MR **95j**:54022
2. A. Dold, R. Thom: *Quasifaserungen und unendliche symmetrische Produkte*; Ann. of Math. 67 (1958) 239–281. MR **20**:3542
3. W.E. Haver: *Locally contractible spaces that are absolute neighborhood retracts*; Proc. Amer. Math. Soc. 40 (1973) 280–284. MR **48**:9645
4. G. Kozłowski: *Factorization of Certain Maps up to Homotopy*; Proc. Amer. Math. Soc. 21 (1969) 88–92. MR **38**:6588
5. J.P. May: *Simplicial Objects in Algebraic Topology*; University of Chicago Press, Midway Reprint (1992). MR **93m**:55025

FACHBEREICH MATHEMATIK, JOHANN WOLFGANG GOETHE-UNIVERSITÄT, ROBERT-MAYER-STRASSE 6–10, 60054 FRANKFURT, GERMANY

E-mail address: guenther@mathematik.uni-frankfurt.d400.de