

NOTES ON HOMOLOGY COBORDISMS OF PLUMBED HOMOLOGY 3-SPHERES

NIKOLAI SAVELIEV

(Communicated by Ronald A. Fintushel)

ABSTRACT. The gauge-theoretical invariants of Donaldson and Seiberg-Witten are used to detect some infinite order elements in the homology cobordism group of integral homology 3-spheres.

This paper is concerned with the homology cobordism group $\Theta_{\mathbb{Z}}^3$ of oriented integral homology 3-spheres. We use S. Donaldson's [D] and M. Furuta's [F] (see also [A]) theorems and the $\bar{\mu}$ -invariant introduced by W. Neumann [N] and L. Siebenmann [S] to prove the following two theorems.

Theorem 1. *Let a homology sphere Σ be the link of an algebraic singularity. If Σ is homology cobordant to zero, then $\bar{\mu}(\Sigma) \geq 0$.*

All Seifert fibered homology spheres are the links of algebraic singularities. Therefore, one can apply this theorem to recover some of the results of R. Fintushel and R. Stern [FS] on homology cobordisms of Seifert fibered homology spheres. For example, it easily follows that for any relatively prime positive integers p and q , Seifert fibered homology spheres $\Sigma(p, q, pqk - 1)$, $k \geq 1$, have infinite order in the group $\Theta_{\mathbb{Z}}^3$. Theorem 1 implies some new results as well; see Section 2.

Theorem 2. *Let Σ be a plumbed homology sphere and assume that $\bar{\mu}(\Sigma) \neq 0$. If Σ bounds a plumbed 4-manifold with even intersection form of rank not exceeding $10|\bar{\mu}(\Sigma)|$, then Σ has infinite order in the group $\Theta_{\mathbb{Z}}^3$.*

One can easily see [NR] that any Seifert sphere $\Sigma(a_1, \dots, a_n)$ with one of the a_i 's even bounds an (essentially unique) plumbed manifold with even intersection form. This intersection form is very often, though not always, of the desired rank not exceeding $10|\bar{\mu}|$. Moreover, in many cases the rank can be reduced with the help of Kirby calculus without changing $\bar{\mu}$ so that the conclusion of Theorem 2 still holds. Such an approach leads in particular to the following result [S1].

Corollary 1. *For any relatively prime integers $p, q \geq 2$ and any odd integer $k \geq 1$ the Seifert fibered homology sphere $\Sigma(p, q, pqk + 1)$ has infinite order in the group $\Theta_{\mathbb{Z}}^3$.*

This result cannot be extended to $\Sigma(p, q, pqk + 1)$ with even k , because, for instance, $\Sigma(2, 3, 13)$ is known to bound a contractible manifold.

Received by the editors November 19, 1996 and, in revised form, February 11, 1997.
1991 *Mathematics Subject Classification.* Primary 57M25, 57R80.

Key words and phrases. Homology 3-sphere, homology cobordism, gauge theory.

It is worth mentioning that in fact $\bar{\mu}(\Sigma)$ vanishes for all known examples of plumbed homology spheres Σ homology cobordant to zero. In [N], W. Neumann conjectured that $\bar{\mu}$ is a homology cobordism invariant. The results of this paper give further evidence which weighs for a positive solution to this conjecture.

1. DEFINITION AND BASIC PROPERTIES OF THE $\bar{\mu}$ -INVARIANT

We recall the definition of the invariant $\bar{\mu}$ by W. Neumann [N]. Note that our definition differs from the original one by a factor of 1/8.

Let Γ be a connected plumbing graph, that is, a connected graph with no cycles, each of whose vertices carries an integer weight e_i , $i = 1, \dots, s$. The matrix $A(\Gamma) = (a_{ij})_{i,j=1,\dots,s}$ with the entries

$$a_{ij} = \begin{cases} e_i, & \text{if } i = j, \\ 1, & \text{if } i \text{ is connected to } j \text{ by an edge,} \\ 0, & \text{otherwise,} \end{cases}$$

is the intersection matrix of the 4-dimensional manifold $P(\Gamma)$ obtained by plumbing D^2 -bundles over 2-spheres according to Γ . This manifold is simply connected.

Disconnected graphs are also allowed. Namely, if $\Gamma = \Gamma_0 + \Gamma_1$ is a disjoint union of Γ_0 and Γ_1 , then $P(\Gamma)$ is the boundary connected sum $P(\Gamma_0) \natural P(\Gamma_1)$.

If Γ is a plumbing graph as above, then $M(\Gamma) = \partial P(\Gamma)$ is an integral homology sphere if and only if $\det A(\Gamma) = \pm 1$. For example, all Seifert fibered homology spheres $\Sigma(a_1, \dots, a_n)$ are of the form $\partial P(\Gamma)$ where Γ is a star-shaped graph; see [NR].

If $M(\Gamma)$ is a homology sphere, there is a unique homology class $w \in H_2(P(\Gamma); \mathbb{Z})$ satisfying the following two conditions. First, w is *characteristic*, that is (dot represents intersection number)

$$(1) \quad w \cdot x \equiv x \cdot x \pmod{2} \quad \text{for all } x \in H_2(P(\Gamma); \mathbb{Z}),$$

and second, all coordinates of w are either 0 or 1 in the natural basis of $H_2(P(\Gamma); \mathbb{Z})$. We call w *the integral Wu class* for $P(\Gamma)$. It follows from [N] that the integer $\text{sign } P(\Gamma) - w \cdot w$ depends only on $M(\Gamma)$ and not on Γ . This integer is divisible by 8, so one can define the Neumann-Siebenmann invariant by the formula

$$(2) \quad \bar{\mu}(M(\Gamma)) = \frac{1}{8}(\text{sign } P(\Gamma) - w \cdot w).$$

It is also easily seen that (1) implies that no two coordinates of w corresponding to adjacent vertices in Γ can both be 1, so it follows that w is spherical, and the modulo 2 reduction of the $\bar{\mu}$ -invariant is the usual Rohlin invariant μ ; see [NR].

2. THE $\bar{\mu}$ -INVARIANT OF ALGEBRAIC LINKS

The plumbed homology spheres have been classified in [EN]. In particular, it has been shown that a plumbed homology sphere Σ is an algebraic link if and only if there exists a plumbing graph Γ such that the intersection form of the manifold $P(\Gamma)$ with $\Sigma = \partial P(\Gamma)$ is negative definite. The simplest case is the Seifert fibered case: any Seifert fibered homology sphere $\Sigma(a_1, \dots, a_n)$ is the link of the singularity of $f^{-1}(0)$, where $f : \mathbb{C}^n \rightarrow \mathbb{C}^{n-2}$ is a map of the form

$$f(z_1, \dots, z_n) = \left(\sum_{k=1}^n b_{1,k} z_k^{a_k}, \dots, \sum_{k=1}^n b_{n-2,k} z_k^{a_k} \right)$$

with sufficiently general coefficient matrix $(b_{i,j})$; see [NR]. For instance,

$$\Sigma(p, q, r) = \{z \in \mathbb{C}^3 \mid \|z\| = \varepsilon \text{ and } z_1^p + z_2^q + z_3^r = 0\}$$

for $\varepsilon > 0$ small enough.

The next simplest case is the following: if p, q, r are pairwise relatively prime integers, as are p', q', r' , then the homology sphere Σ obtained by splicing $\Sigma(p, q, r)$ and $\Sigma(p', q', r')$ along the singular fibers of degrees r and r' is the link of singularity if and only if $rr' > pp'qq'$; see [NW], § 4.

Theorem 1. *Let a homology sphere Σ be the link of an algebraic singularity. If Σ is homology cobordant to zero, then $\bar{\mu}(\Sigma) \geq 0$.*

Proof. Since Σ is an algebraic link, one may assume that Σ is the boundary of a plumbed 4-manifold $P(\Gamma)$ whose intersection form is negative definite. Suppose that Σ bounds a smooth homology ball M . Let us consider the manifold $W = P(\Gamma) \cup_{\Sigma} (-M)$. This is a smooth closed oriented manifold whose intersection form is naturally isomorphic to the intersection form of $P(\Gamma)$, in particular, is negative definite. By S. Donaldson's Theorem 1 from [D], this form is diagonalizable over the integers.

We use this fact to evaluate $\bar{\mu}(\Sigma)$. Obviously, $\text{sign } P(\Gamma) = -s$, where s is the number of vertices in the graph Γ , so we only need to find the Wu class w . In the standard basis associated with the plumbing, the matrix A of the intersection form of $P(\Gamma)$ takes the form $A = U^t(-E)U$, where $U \in \text{SL}_s(\mathbb{Z})$ and E is the identity matrix. The defining relation (1) translates to

$$w^t U^t(-E)Ux \equiv x^t U^t(-E)Ux \pmod{2} \quad \text{for all } x \in H_2(P(\Gamma); \mathbb{Z}),$$

or, equivalently,

$$(Uw)^t(-E)y \equiv y^t(-E)y \pmod{2} \quad \text{for all } y \in H_2(P(\Gamma); \mathbb{Z}).$$

Therefore, Uw is characteristic for $-E$, in particular, all the coordinates of Uw are odd. Now, we have

$$w.w = w^t U^t(-E)Uw = -(Uw)^t(Uw),$$

which is equal to minus the square of the Euclidean length of the vector Uw . Since all the coordinates of Uw are odd, $w.w \leq -s$, and therefore $\bar{\mu}(\Sigma) \geq 0$. \square

Corollary 2. *If a plumbed homology sphere Σ is an algebraic link and $\bar{\mu}(\Sigma) < 0$, then Σ has infinite order in the group $\Theta_{\mathbb{Z}}^3$.*

Proof. Let $m\Sigma = \Sigma \# \dots \# \Sigma$ (m times). Since $\bar{\mu}$ is additive with respect to connected sums, $\bar{\mu}(m\Sigma) = m\bar{\mu}(\Sigma) < 0$, therefore, $m\Sigma$ is not homology cobordant to zero for any m . \square

Example. For any relatively prime integers $p, q > 0$, one can easily see that $\bar{\mu}(\Sigma(p, q, pq - 1)) < 0$. Therefore, all these homology spheres are of infinite order in $\Theta_{\mathbb{Z}}^3$. In particular, the Poincaré homology sphere $\Sigma(2, 3, 5)$ is of infinite order.

Example. Homology spheres $\Sigma(p, q, pqk - 1)$ have infinite order in $\Theta_{\mathbb{Z}}^3$ for any $k \geq 1$. If k is odd, this holds since $\bar{\mu}(\Sigma(p, q, pqk - 1))$ equals $\bar{\mu}(\Sigma(p, q, pq - 1))$ and is therefore negative. If k is even, then $\bar{\mu}(\Sigma(p, q, pqk - 1)) = 0$, and the argument of Theorem 1 cannot be applied directly. Let A be the intersection form of a negative definite plumbing X bounding $\Sigma(p, q, pqk - 1)$, k even. We perform a (-1) -surgery

on the singular fiber of degree $pqk - 1$ to get $\Sigma(p, q, pq(k + 1) - 1)$. The union of the cobordism X with the trace of this surgery is a plumbed manifold with the intersection form $A \oplus (-1)$. Now, if $\Sigma(p, q, pqk - 1)$ bounded an acyclic manifold, A would be diagonalizable, and so would be $A \oplus (-1)$. This contradicts the fact that $\bar{\mu}(\Sigma(p, q, pq(k + 1) - 1)) < 0$. A similar argument shows that none of the multiples of $\Sigma(p, q, pqk - 1)$ is homology cobordant to zero.

Many Seifert homology spheres $\Sigma(a_1, \dots, a_n)$ having infinite order in the group $\Theta_{\mathbb{Z}}^3$ by Corollary 2 can also be detected by the R -invariant of R. Fintushel and R. Stern,

$$R(a_1, \dots, a_n) = \frac{2}{a} - 3 + n + \sum_{i=1}^n \frac{2}{a_i} \sum_{k=1}^{a_i-1} \cot\left(\frac{\pi ak}{a_i^2}\right) \cot\left(\frac{\pi k}{a_i}\right) \sin^2\left(\frac{\pi k}{a_i}\right),$$

where $a = a_1 \cdot \dots \cdot a_n$. A theorem in [FS] implies that if $\Sigma(a_1, \dots, a_n)$ bounds a homology ball, then $R(a_1, \dots, a_n) < 0$. There are however Seifert fibered homology spheres which are not homology cobordant to zero and which can be detected by $\bar{\mu}$ and not by R , and vice versa.

Example. Both Seifert spheres $\Sigma(2, 11, 19)$ and $\Sigma(3, 5, 7)$ are not homology cobordant to zero. As to the former one, this follows from the fact that $\bar{\mu}(\Sigma(2, 11, 19)) = -1$ is negative (though $R(2, 11, 19) = -1$); the latter one has $\bar{\mu}(\Sigma(3, 5, 7)) = 0$ and the result follows from $R(3, 5, 7) = 1$.

The $\bar{\mu}$ -invariant also detects some plumbed homology spheres which have infinite order in $\Theta_{\mathbb{Z}}^3$ but are not Seifert fibered.

Example. Let Σ be the splice of $\Sigma(4, 7, 9)$ and $\Sigma(2, 3, 25)$ along the singular fibers of degrees 9 and 25. This manifold is an algebraic link. By using the additivity of the $\bar{\mu}$ -invariant proven in [Du] and [S2], we find that

$$\bar{\mu}(\Sigma) = \bar{\mu}(\Sigma(4, 7, 9)) + \bar{\mu}(\Sigma(2, 3, 25)) = -2 + 0 < 0.$$

Therefore, Σ has infinite order in $\Theta_{\mathbb{Z}}^3$ (although the Rohlin invariant $\mu(\Sigma)$ equals 0 modulo 2).

3. MORE CONSTRAINTS FROM EVEN PLUMBING

In 1995, M. Furuta [F] used the Seiberg-Witten invariants to prove the so-called 10/8-conjecture, which says that if A is the intersection form of a smooth closed spin 4-manifold, then $\text{rank } A / |\text{sign } A| > 10/8$. In this section we use this result to prove the following theorem.


Theorem 2. *Let Σ be a plumbed homology sphere and assume that $\bar{\mu}(\Sigma) \neq 0$. If Σ bounds a plumbed 4-manifold with even intersection form of rank not exceeding $10|\bar{\mu}(\Sigma)|$, then Σ has infinite order in the group $\Theta_{\mathbb{Z}}^3$.*

Proof. Let $P(\Gamma)$ be a plumbed manifold with boundary $\partial P(\Gamma) = \Sigma$, such that its intersection form $A = A(\Gamma)$ is even and has rank not exceeding $10|\bar{\mu}(\Sigma)|$. Since A is even, its Wu class w vanishes. Therefore, $8 \cdot \bar{\mu}(\Sigma) = \text{sign } A$, and moreover A is isomorphic over the integers to $|\bar{\mu}(\Sigma)| \cdot E_8 \oplus b \cdot H$. Since $\text{rank}(A) \leq 10|\bar{\mu}(\Sigma)|$, we get that $\text{rank } A / |\text{sign } A| \leq 10/8$.


Now, suppose that Σ bounds a smooth homology ball M , and consider the manifold $W = P(\Gamma) \cup_{\Sigma} (-M)$. This is a smooth closed oriented spin 4-manifold whose intersection form Q is isomorphic to $|\bar{\mu}(\Sigma)| \cdot E_8 \oplus b \cdot H$. Since $\text{rank } Q / |\text{sign } Q| \leq 10/8$, we get a contradiction with the 10/8 conjecture.

The argument can be repeated with Σ replaced by any of its multiples, which proves the theorem. \square


For a complete proof of Corollary 1 formulated in the introduction we refer the reader to [S1], and only prove it here in the simplest case $p = 2$. If $q \equiv 3 \pmod 4$, then $\Sigma(2, q, 2qk + 1)$, k odd, is the boundary of the 4-manifold obtained by plumbing according to the following diagram:



The intersection form of this manifold is isomorphic to $\frac{q+1}{4} \cdot E_8 \oplus H$. If $q \equiv 1 \pmod 4$, the homology sphere $\Sigma(2, q, 2qk + 1)$ bounds plumbing



with intersection form $\frac{q-1}{4} \cdot E_8 \oplus 3 \cdot H$. To deal with this case, we need to reduce the rank by surgering out two “hyperbolics” H . This can be done as follows. First, redraw the plumbing diagram:



After two obvious blow-downs we get the manifold in Figure 1 with the intersection form $\frac{q-1}{4} \cdot E_8 \oplus 2 \cdot H$. An equivalent link description is shown in Figure 2. After two more blow-downs we get rid of another “hyperbolic” H . The final link description is shown in Figure 3.




FIGURE 1




FIGURE 2




FIGURE 3

The corresponding intersection form is isomorphic to $\frac{q-1}{4} \cdot E_8 \oplus H$, and an argument similar to that in Theorem 2 can be used to show that $\Sigma(2, q, 2qk + 1)$, k odd, has infinite order in $\Theta_{\mathbb{Z}}^3$.

Example. In addition to those listed in Corollary 1, there are of course many homology spheres having infinite order in $\Theta_{\mathbb{Z}}^3$ due to Theorem 2. For example, $\Sigma(8, 13, 21)$ is of this sort because it has $\bar{\mu}$ -invariant equal 4 and can be obtained by plumbing on an even weighted graph of rank 34.

REFERENCES

- [A] S. Akbulut, *Lectures on Seiberg–Witten Invariants*, Turkish J. Math. **20** (1996), 95–118. MR **97g**:57038
- [D] S. Donaldson, *The orientation of Yang–Mills moduli spaces and 4-manifold topology*, J. Diff. Geom. **26** (1987), 397–428. MR **88j**:57020
- [Du] N. Duchon, *Involutions of plumbed manifolds*, Ph.D. Thesis, University of Maryland, College Park, 1982.
- [EN] D. Eisenbud, W. Neumann, *Three-dimensional Link Theory and Invariants of Plane Curve Singularities*, Annals of Math. Studies **110**, Princeton, 1985. MR **87g**:57007
- [FS] R. Fintushel, R. Stern, *Pseudofree orbifolds*, Ann. Math. **122** (1985), 335–364. MR **87a**:57027
- [F] M. Furuta, *Monopole equation and the 11/8 conjecture*, Preprint.
- [N] W. Neumann, *An invariant of plumbed homology spheres*, Lecture Notes in Math. **788**, Springer-Verlag, Berlin and New-York, 1980, 125–144. MR **82j**:57033
- [NR] W. Neumann, F. Raymond, *Seifert manifolds, plumbing, μ -invariant and orientation reversing maps*, Lecture Notes in Math. 664, Springer-Verlag, Berlin and New-York, 1978, 163–196. MR **80e**:57008
- [NW] W. Neumann, J. Wahl, *Casson invariant of links of singularities*, Comm. Math. Helv. **65** (1990), 58–78. MR **91c**:57022
- [S1] N. Saveliev, *Dehn surgery along torus knots*, Topology and Its Appl. (to appear).

- [S2] N. Saveliev, *Floer homology and 3-manifold invariants*, Ph.D. Thesis, University of Oklahoma, Norman, 1995.
- [S] L. Siebenmann, *On vanishing of the Rohlin invariant and nonfinitely amphicheiral homology 3-spheres*, Lecture Notes in Math. **788**, Springer-Verlag, Berlin and New-York, 1980, 172–222. MR **81k**:57011

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48109
E-mail address: `saveliev@math.lsa.umich.edu`