

THE BAIRE CATEGORY THEOREM AND THE EVASION NUMBER

MASARU KADA

(Communicated by Carl G. Jockusch)

ABSTRACT. In this paper we prove that $\epsilon \leq \text{cov}(\mathcal{M})$ where ϵ is the evasion number defined by Blass. This answers negatively a question asked by Brendle and Shelah.

1. INTRODUCTION

Let $\text{cov}(\mathcal{M})$ denote the smallest size of a family of meager sets whose union covers the real line. The combinatorial characterization for $\text{cov}(\mathcal{M})$ has been studied by Miller and Bartoszyński, and the following result is established. For $g \in \omega^\omega$, let $\mathcal{S}^g = \prod_{n < \omega} [\omega]^{\leq g(n)}$. Each element of \mathcal{S}^g is called a *slalom*.

Theorem 1.1 ([1, Lemma 2.4.2]). *The following cardinalities are the same:*

1. $\text{cov}(\mathcal{M})$.
2. the smallest size of $F \subseteq \omega^\omega$ such that for every $h \in \omega^\omega$ there exists $f \in F$ with $f(n) \neq h(n)$ for all but finitely many $n < \omega$.
3. the smallest cardinality κ satisfying the following: for every $g \in \omega^\omega$ there exists $F \subseteq \omega^\omega$ of size κ such that, for all $\varphi \in \mathcal{S}^g$, there exists $f \in F$ with $f(n) \notin \varphi(n)$ for all but finitely many $n < \omega$. \square

Blass [2] introduced a combinatorial concept called ‘predicting and evading’, and using this he defined the following cardinal invariant. Let \mathcal{P} be the collection of functions π from $\omega^{<\omega}$ to ω . Here we call each such π a *predictor*.

Definition 1.2 ([2]). The *evasion number* ϵ is the smallest size of $F \subseteq \omega^\omega$ such that, for every $\pi \in \mathcal{P}$ and $X \in [\omega]^\omega$, there exists $f \in F$ with $f(n) \neq \pi(f \upharpoonright n)$ for infinitely many $n \in X$.

Brendle and Shelah [3], [4] studied the relations between ϵ and other cardinal invariants, and asked whether $\epsilon > \text{cov}(\mathcal{M})$ is consistent [4, Subsection 3.4]. Here we give a negative answer to this question.

Received by the editors February 10, 1997 and, in revised form, April 7, 1997.

1991 *Mathematics Subject Classification*. Primary 03E05.

The author was supported by JSPS Research Fellowships for Young Scientists. The author was also supported by Grant-in-Aid for Scientific Research (Encouragement for Research Fellow, No. 97-03909), Ministry of Education, Science and Culture.

©1998 American Mathematical Society

2. THE MAIN RESULT

We introduce a different form of evasion number by modifying the original definition due to Blass.

Definition 2.1. \mathfrak{e}^* is the smallest size of $F \subseteq \omega^\omega$ such that, for every $\pi \in \mathcal{P}$ there exists $f \in F$ with $f(n) \neq \pi(f \upharpoonright n)$ for all but finitely many $n < \omega$.

Clearly $\mathfrak{e} \leq \mathfrak{e}^*$ holds, and it is easily seen from Theorem 1.1 that $\text{cov}(\mathcal{M}) \leq \mathfrak{e}^*$.

We show that \mathfrak{e}^* gives another combinatorial characterization for $\text{cov}(\mathcal{M})$. We prove the following theorem by modifying the proof for $\mathfrak{e} \leq \mathfrak{d}$, which is due to Blass [2, Theorem 13].

Theorem 2.2. $\mathfrak{e}^* = \text{cov}(\mathcal{M})$.

Proof. For a function $h \in \omega^{\omega \times \omega}$, define $x_h \in \omega^\omega$ recursively so that $x_h(n) = h(n, 1 + \max\{x_h(i) : i < n\})$. Next, for a predictor $\pi \in \mathcal{P}$, define a function φ_π from $\omega \times \omega$ to $[\omega]^{<\omega}$ by letting $\varphi_\pi(n, k) = \{\pi(t) : t \in k^n\}$. By identifying $\omega \times \omega$ with ω , we can regard φ_π as a slalom in \mathcal{S}^g for a suitable $g \in \omega^\omega$ which does not depend on π .

Now we prove the following: for $n < \omega$, if $h(n, k) \notin \varphi_\pi(n, k)$ for all k , then $x_h(n) \neq \pi(x_h \upharpoonright n)$. Suppose that $h(n, k) \notin \varphi_\pi(n, k)$ for all k . Let $k = 1 + \max\{x_h(i) : i < n\}$. Then $x_h \upharpoonright n \in k^n$ and hence $\pi(x_h \upharpoonright n) \in \varphi_\pi(n, k)$. On the other hand, $x_h(n) = h(n, k) \notin \varphi_\pi(n, k)$. Thus, $x_h(n) \neq \pi(x_h \upharpoonright n)$.

By Theorem 1.1, we can choose $F \subseteq \omega^{\omega \times \omega}$ of size $\text{cov}(\mathcal{M})$ so that, for each predictor $\pi \in \mathcal{P}$, there is $f \in F$ with $f(n, k) \notin \varphi_\pi(n, k)$ for all but finitely many $(n, k) \in \omega \times \omega$. Then the set $\{x_f : f \in F\}$ witnesses $\mathfrak{e}^* \leq \text{cov}(\mathcal{M})$, and hence $\mathfrak{e}^* = \text{cov}(\mathcal{M})$. \square

Corollary 2.3. $\mathfrak{e} \leq \text{cov}(\mathcal{M})$. \square

Let $\text{non}(\mathcal{M})$ denote the smallest size of a nonmeager set of reals. We can characterize $\text{non}(\mathcal{M})$ in a dual fashion, using [1, Lemma 2.4.8] instead of Theorem 1.1.

Theorem 2.4. $\text{non}(\mathcal{M})$ is the smallest size of $\Pi \subseteq \mathcal{P}$ satisfying the following: for every $f \in \omega^\omega$ there exists $\pi \in \Pi$ such that $f(n) = \pi(f \upharpoonright n)$ for infinitely many $n < \omega$. \square

3. A REMARK ON THE EVASION IDEAL

Brendle [3, Subsection 3.5] introduced the notion of the evasion ideal, that is, the σ -ideal generated by the sets of the form

$$\{f \in \omega^\omega : f(n) = \pi(f \upharpoonright n) \text{ for all but finitely many } n \in X\}$$

for $\pi \in \mathcal{P}$ and $X \in [\omega]^\omega$. He considered the smallest size of a subset of ω^ω which does not belong to this ideal.

Definition 3.1 ([4, Definition 3.1]). $\mathfrak{e}(\omega)$, the uniformity of the evasion ideal, is the smallest size of $F \subseteq \omega^\omega$ satisfying the following: for any countable family of pairs $\{\langle \pi_i, X_i \rangle : i < \omega\} \subseteq \mathcal{P} \times [\omega]^\omega$ there is $f \in F$ such that for each $i < \omega$ we have $f(n) \neq \pi_i(f \upharpoonright n)$ for infinitely many $n \in X_i$.

Clearly $\mathfrak{e} \leq \mathfrak{e}(\omega)$ holds. Brendle and Shelah asked whether $\mathfrak{e} = \mathfrak{e}(\omega)$ can be proved in ZFC, and they presented the following partial answer.

Theorem 3.2 ([4, Theorem 3.3]). $\mathfrak{e} \geq \min\{\mathfrak{e}(\omega), \text{cov}(\mathcal{M})\}$. Thus either $\mathfrak{e} < \text{cov}(\mathcal{M})$ or $\mathfrak{e}(\omega) \leq \text{cov}(\mathcal{M})$ implies $\mathfrak{e} = \mathfrak{e}(\omega)$. \square

We show that the latter assumption of the above theorem holds in ZFC.

Theorem 3.3. $\mathfrak{e}(\omega) \leq \mathfrak{e}^*$.

Proof. Fix $F \subseteq \omega^\omega$ of size less than $\mathfrak{e}(\omega)$ arbitrarily. Then we can choose a countable set of pairs $\{\langle \pi_i, X_i \rangle : i < \omega\} \subseteq \mathcal{P} \times [\omega]^\omega$ so that, for every $f \in F$, there is $i < \omega$ such that $f(n) = \pi_i(f \upharpoonright n)$ for all but finitely many $n \in X_i$. By shrinking X_i 's if necessary, we can assume that X_i 's are pairwise disjoint. Now define $\pi \in \mathcal{P}$ as follows: for $t \in \omega^{<\omega}$ if $|t| \in X_i$ for some $i < \omega$ then $\pi(t) = \pi_i(t)$; otherwise $\pi(t)$ is arbitrary. Then for all $f \in F$ we have $f(n) = \pi(f \upharpoonright n)$ for infinitely many $n < \omega$. \square

Corollary 3.4. $\mathfrak{e} = \mathfrak{e}(\omega)$.

Proof. By Theorems 2.2, 3.2 and 3.3. \square

ACKNOWLEDGEMENT

I would like to thank the referee for his helpful comment, which gives another view of the main theorem I had not noticed.

REFERENCES

- [1] T. Bartoszyński and H. Judah, *Set theory: On the structure of the real line*, A. K. Peters, Wellesley, Massachusetts, 1995. MR **96k**:03002
- [2] A. Blass, *Cardinal characteristics and the product of countably many infinite cyclic groups*, J. Algebra **169** (1994), 512–540. MR **95h**:20069
- [3] J. Brendle, *Evasion and prediction — the Specker phenomenon and Gross spaces*, Forum Math. **7** (1995), 513–541. MR **96i**:03042
- [4] J. Brendle and S. Shelah, *Evasion and prediction II*, J. London Math. Soc. **53** (1996), 19–27. MR **97d**:03061

OSAKA PREFECTURE UNIVERSITY, SAKAI, OSAKA, 599-8531 JAPAN
E-mail address: kada@mi.cias.osakafu-u.ac.jp