

## $C^1$ APPROXIMATIONS OF INERTIAL MANIFOLDS VIA FINITE DIFFERENCES

KAZUO KOBAYASI

(Communicated by David R. Larson)

ABSTRACT. We construct an inertial manifold for the evolution equation as a limit of the inertial manifolds for the difference approximations of the Trotter-Kato type and show that this limit is taken in a  $C^1$  topology.

### 1. INTRODUCTION

We shall study a class of nonlinear dissipative partial differential equations (PDE for short) that have inertial manifolds (IM for short). The theory of IMs allows us to reduce the long-time behavior of the PDE to that of a finite-dimensional dynamical system. In order to implement the reduced finite dynamical system computationally, one would need to know the explicit form of the IM. However, even when existence of an IM can be established, the theory does not provide it in an explicit form. Thus, a number of approximate IMs have been considered in the literature. See, e.g., [2], [3], [4], [6], [9], [10], [11], [12].

In this paper, from the point of finite differences we shall construct an IM for the PDE. Indeed, the IM is constructed as a limit of IMs for the associated finite difference equations and the limit is taken in a  $C^1$  topology. This means that on one hand the existence of the IMs for the finite difference equations assures the existence of an IM for the PDE, and on the other the IMs for the finite differences can be viewed as a small  $C^1$  perturbation of that for the PDE. The  $C^1$  closeness of the IMs would be a necessary and important step toward establishing a relationship between the dynamics of the PDE and its approximation (see [8], [14]).

Each of the PDEs can be viewed as an evolution equation in a Banach space  $Y$

$$(1.1) \quad du(t)/dt = Au(t) + Fu(t), \quad t \in \mathbb{R}^+ \equiv [0, \infty),$$

with a closed linear operator  $A$  in  $Y$  and  $F \in C^1(X, Y)$ , where  $X$  is a Banach space continuously embedded in  $Y$ .

We approximate (1.1) by the following discrete scheme:

$$(1.2) \quad x_\ell^n = C(\lambda_\ell)x_\ell^{n-1} + \lambda_\ell K_\ell F_\ell(x_\ell^{n-1}), \quad n, \ell \in \mathbb{N},$$

---

Received by the editors July 29, 1997.

1991 *Mathematics Subject Classification.* Primary 47H20; Secondary 35K55.

*Key words and phrases.* Inertial manifold, long-time behavior, finite dynamical system, evolution equation.

This research was partially supported by Waseda University Grant for special Research Projects 97A-81.

in a space  $Y_\ell$  approximating  $Y$  in some sense, where  $\lambda_\ell \downarrow 0$  as  $\ell \rightarrow \infty$ ,  $C(\lambda_\ell)$  and  $K_\ell$  are given operators in  $B(Y_\ell, Y_\ell)$  and  $F_\ell$  is a given nonlinear operator in  $Y_\ell$  stated below. We denote by  $B(W, Z)$  the space of bounded linear operators from a Banach space  $W$  into a Banach space  $Z$ . The norm in  $B(W, Z)$  will be denoted by  $\|\cdot\|_{W,Z}$ .

## 2. ASSUMPTIONS AND RESULT

We shall make the following assumptions.

(C1) Let  $X$  and  $Y$  be reflexive Banach spaces such that  $X$  is densely and continuously embedded in  $Y$  and such that  $Y = Y_1 \oplus Y_2$ , the direct sum of a finite dimensional subspace  $Y_1$  and a closed subspace  $Y_2$ .

(C2) For each  $\ell \in \mathbb{N}$  let  $X_\ell$  and  $Y_\ell$  be Banach spaces with norms  $\|\cdot\|_\ell$  and  $|\cdot|_\ell$ , respectively, such that  $X_\ell$  is continuously embedded in  $Y_\ell$ . Moreover, there exist  $V_\ell \in B(Y, Y_\ell) \cap B(X, X_\ell)$  and  $W_\ell \in B(Y_\ell, Y) \cap B(X_\ell, X)$  such that  $\lim_{\ell \rightarrow \infty} |V_\ell y|_\ell = |y|$ ,  $\lim_{\ell \rightarrow \infty} \|V_\ell x\|_\ell = \|x\|$ ,  $\lim_{\ell \rightarrow \infty} |W_\ell V_\ell y - y| = 0$  and  $V_\ell W_\ell z = z$  for  $x \in X, y \in Y, z \in Y_\ell$  and such that both  $\|W_\ell\|_{Y_\ell, Y}$  and  $\|W_\ell\|_{X_\ell, X}$  are bounded in  $\ell$ .

(C3) There exist closed subspaces  $Y_{\ell 1}$  and  $Y_{\ell 2}$  such that  $Y_\ell = Y_{\ell 1} \oplus Y_{\ell 2}$ ,  $V_\ell P_i = P_{\ell i} V_\ell$  and  $W_\ell P_{\ell i} = P_i W_\ell$  for  $i = 1, 2$ , where  $P_i$  (resp.  $P_{\ell i}$ ) denotes a projection from  $Y$  onto  $Y_i$  (resp.  $Y_{\ell i}$ ).

(C4) The linear operators  $C(\lambda_\ell)$  and  $K_\ell$  satisfy: (i) there exist  $M \geq 0$  and  $\omega \geq 0$  such that  $|C(\lambda_\ell)^n y|_\ell \leq M e^{\omega n \lambda_\ell} |y|_\ell$  and  $|K_\ell y|_\ell \leq M e^{\omega \lambda_\ell} |y|_\ell$  for  $\ell, n \in \mathbb{N}, y \in Y_\ell$ ; (ii)  $\lim_{\ell \rightarrow \infty} |(K_\ell - I)V_\ell y|_\ell = 0$  for  $y \in Y$ ; (iii) for each  $\ell, \ell' \in \mathbb{N}$  and  $i = 1, 2$ ,  $C(\lambda_\ell)$  commutes with  $P_{\ell i}$ ,  $C(\lambda_\ell)$  with  $K_\ell$ ,  $K_\ell$  with  $P_{\ell i}$ ,  $\tilde{C}(\lambda_\ell)$  with  $\tilde{C}(\lambda_{\ell'})$  and  $\tilde{C}(\lambda_\ell)$  with  $\tilde{K}_{\ell'}$ , respectively, where  $\tilde{C}(\lambda_\ell) = W_\ell C(\lambda_\ell) V_\ell$  and  $\tilde{K}_\ell = W_\ell K_\ell V_\ell$ .

(C5)  $A$  is a densely defined linear operator in  $Y$  such that  $Y_1 \subset D(A)$ , the range of  $I - \lambda_0 A$  is dense in  $Y$  for some  $\lambda_0 > 0$  and

$$\lim_{\ell \rightarrow \infty} |\lambda_\ell^{-1} (C(\lambda_\ell) - I)V_\ell y - V_\ell A y|_\ell = 0 \quad \text{for } y \in D(A).$$

(C6) The inverse of  $C(\lambda_\ell)P_{\ell 1}$  exists in  $B(Y_{\ell 1})$  and there exist constants  $\alpha, \beta > 0$ ,  $\gamma \in [0, 1)$ ,  $\eta < -\max\{\alpha, \beta\}$  and  $M_1, \dots, M_5 \geq 0$  such that

$$(2.1) \quad \|P_{\ell 1} y\|_\ell \leq M_1 |P_{\ell 1} y|_\ell,$$

$$(2.2) \quad |[C(\lambda_\ell)P_{\ell 1}]^{-n} P_{\ell 1} y|_\ell \leq M_2 e^{-(\alpha+\eta)n\lambda_\ell} |y|_\ell,$$

$$(2.3) \quad \|C(\lambda_\ell)^n P_{\ell 2} x\|_\ell \leq M_3 e^{(\eta-\beta)n\lambda_\ell} \|x\|_\ell,$$

$$(2.4) \quad \|C(\lambda_\ell)^n P_{\ell 2} K_\ell y\|_\ell \leq \{M_4((n+1)\lambda_\ell)^{-\gamma} + M_5\} e^{(\eta-\beta)n\lambda_\ell} |y|_\ell$$

for  $n \geq 0, \ell \geq 1, x \in X_\ell, y \in Y_\ell$ .

(C7)  $F_\ell \in C^1(X_\ell, Y_\ell)$  and there exists a constant  $L_F \geq 0$  satisfying

$$|F_\ell(\xi_1) - F_\ell(\xi_2)|_\ell \leq L_F \|\xi_1 - \xi_2\|_\ell \quad \text{for } \ell \in \mathbb{N}, \xi_1, \xi_2 \in X_\ell.$$

(C8) For each  $x, z \in X$  and each positive sequence  $\{\nu_\ell\}$  convergent to 0 we have

$$\begin{aligned} \lim_{\ell \rightarrow \infty} |F_\ell(V_\ell x) - V_\ell F(x)|_\ell &= 0, \\ \lim_{\ell \rightarrow \infty} |DF_\ell(V_\ell x)V_\ell z - V_\ell DF(x)z|_\ell &= 0, \quad \text{and} \\ \lim_{\ell \rightarrow \infty} \left( \sup_{\|\xi\|_\ell \leq \nu_\ell} |(DF_\ell(V_\ell x + \xi) - DF_\ell(V_\ell x))V_\ell z|_\ell \right) &= 0. \end{aligned}$$

Then we have

**Theorem.** *Let (C1)-(C8) be satisfied and  $F \in C^1(X, Y)$ . In addition we assume*

$$(G) \quad K(\alpha, \beta)L_F < 1 \quad \text{and} \quad \frac{M_2M'_3K(\alpha, \beta)L_F}{1 - K(\alpha, \beta)L_F} < 1$$

where

$$(2.5) \quad k(\alpha, \beta) = M\{M_1M_2\alpha^{-1} + M'_4\Gamma(1 - \gamma)\beta^{\gamma-1} + M'_5\beta^{-1}\},$$

$M'_i = M_i \max\{1, \liminf_{\ell \rightarrow \infty} \|W_\ell\|_{X_\ell, X}\}$  for  $i = 3, 4, 5$  and  $\Gamma$  denotes the gamma function. Then, (1.1) (resp. (1.2)) has an inertial manifold  $\mathcal{M}$  (resp.  $\mathcal{M}_\ell$ ) represented as a graph of a function  $h \in C^1(Y_1, P_2X)$  (resp.  $h_\ell \in C^1(Y_{\ell 1}, P_{\ell 2}X_\ell)$ ). (See, e.g., [2], [5], [12] for the definitions of the IMs.) Moreover, it holds that for every bounded set  $B \subset Y_1$

$$(2.6) \quad \limsup_{\ell \rightarrow \infty} \sup_{y \in B} \|h_\ell(V_\ell y) - V_\ell h(y)\|_\ell = 0, \quad \text{and}$$

$$(2.7) \quad \limsup_{\ell \rightarrow \infty} \sup_{y \in B} \|Dh_\ell(V_\ell y) - V_\ell Dh(y)\|_{Y_1, X_\ell} = 0.$$

### 3. PROOF

**Existence:** To prove the existence of IMs we use the results of [11] and [12] (also see [1]). By (C4) and (C5) the discrete version of the Trotter-Kato theorem (see [13, Theorem 6.7]) shows that  $\bar{A}$ , the closure of  $A$  in  $Y$ , generates a  $C_0$ -semigroup  $\{S(t); t \geq 0\}$  on  $Y$  satisfying

$$(3.1) \quad \lim_{k_\ell \lambda_\ell \rightarrow t} |C(\lambda_\ell)^{k_\ell} V_\ell y - V_\ell S(t)y|_\ell = 0 \quad \text{for } y \in Y, t \geq 0.$$

In particular, (3.1) together with (C3) and (C4) implies that  $|P_i S(t)y - S(t)P_i y| = \lim_{\ell \rightarrow \infty} |V_\ell(P_i S(t)y - S(t)P_i y)|_\ell = 0$ , which shows that  $P_i S(t) = S(t)P_i$  for  $i = 1, 2$  and  $t \geq 0$ . Set  $A_1 = \bar{A}|_{Y_1}$ . Then,  $D(A_1) = Y_1$  by (C5), so that  $A_1 \in B(Y_1)$  by the closed graph theorem. The family  $\{S_1(t)\}$  defined by  $S_1(t) = S(t)|_{Y_1}$  forms a uniformly continuous group on  $Y_1$  with the infinitesimal generator  $A_1$ . This proves conditions (S2) and (S3) in [11]. To show condition (S4) in [11] we fix  $y \in Y$  and  $t \geq 0$ . By (2.2) and (3.1) we have one of the inequalities in (S4):

$$\begin{aligned} |S_1(-t)P_1 y| &= \lim_{k_\ell \lambda_\ell \rightarrow t} |V_\ell S_1(-t)P_1 y|_\ell \\ &\leq M_2 \lim_{k_\ell \lambda_\ell \rightarrow t} e^{-(\alpha+\eta)k_\ell \lambda_\ell} |V_\ell y|_\ell = M_2 e^{-(\alpha+\eta)t} |y|. \end{aligned}$$

Moreover, we have by (2.4)

$$(3.2) \quad \begin{aligned} &\|W_\ell C(\lambda_\ell)^{k_\ell} P_{\ell 2} K_\ell V_\ell y\| \\ &\leq \|W_\ell\|_{X_\ell, X} \{M_4((k_\ell + 1)\lambda_\ell)^{-\gamma} + M_5\} e^{(\eta-\beta)k_\ell \lambda_\ell} |V_\ell y|_\ell, \end{aligned}$$

which implies that  $\|W_\ell C(\lambda_\ell)^{k_\ell} P_{\ell 2} K_\ell V_\ell y\|$  is bounded as  $k_\ell \lambda_\ell \rightarrow t > 0$ . Note that by (3.1)  $W_\ell C(\lambda_\ell)^{k_\ell} P_{\ell 2} K_\ell V_\ell y$  converges as  $\ell \rightarrow \infty$  to  $S(t)P_2 y$  in  $Y$ . Since  $X$  is reflexive and  $Y^*$  is dense in  $X^*$  by assumption, it also converges weakly in  $X$  to  $S(t)P_2 y$ . Then, passing to the limit in (3.2) yields

$$\|S(t)P_2 y\| \leq \liminf_{\ell \rightarrow \infty} \|W_\ell\|_{X_\ell, X} \{M_4 t^{-\gamma} + M_5\} e^{(\eta-\beta)t} |y|,$$

the second inequality in (S4). Likewise, the remainder inequalities in (S4) will be proved.

Finally, let us prove condition (S1) in [11]. Note that  $S(t)Y = P_1S(t)Y + S(t)P_2Y \subset X$  for  $t > 0$  and  $S(\cdot)x \in C(\mathbb{R}^+; Y)$  for  $x \in X$ . Since  $\|S(t)x\| \leq C\|x\|$  by (S4), we find that  $S(\cdot)x$  is weakly continuous in  $X$  and hence  $S(\cdot)x \in C(\mathbb{R}^+; X)$  (see [7, Theorem 10.6.5]). Thus (S1) is proved. Consequently, we can conclude from [11] and [1] that (1.1) has an IM, provided (G) is satisfied. The existence of an IM for (1.2) is already proved in [12] under the same assumptions as above.

**Convergence:** Next we prove (2.6) and (2.7). To this end recall ([12]) that the IM  $\mathcal{M}$  for (1.1) is constructed as the graph of the function  $h : Y_1 \rightarrow X$  defined by  $h(y) = f(y, 0) - y$ , where  $f(y, t)$  is the unique function in  $C^1(Y_1, C_{\eta+\varepsilon}(\mathbb{R}^+, X))$  for all  $\varepsilon \in [0, \alpha)$  satisfying

$$(3.3) \quad f(y, s) = S(-t)y - \int_0^t S(s-t)P_1F(f(y, s))ds + \int_t^\infty S(s-t)P_2F(f(y, s))ds$$

for  $y \in Y_1$  and  $t \geq 0$ . Here  $C_\eta(\mathbb{R}^+, X)$  denotes the Banach space of continuous functions  $u : \mathbb{R}^+ \rightarrow X$  with the norm  $\|u\|^{(\eta)} = \sup_{t \geq 0} e^{\eta t} \|u(t)\|$ . Similarly, the IM  $\mathcal{M}_\ell$  for (1.2) is constructed as the graph of the function  $h_\ell : Y_{\ell 1} \rightarrow X_\ell$  defined by  $h_\ell(\xi) = \varphi_\ell(\xi, 0) - \xi$ , where  $\varphi_\ell(\xi, n)$  is the unique function in  $C^1(Y_{\ell 1}, c_{\eta+\varepsilon}(\mathbb{N}, X_\ell))$  for all  $\varepsilon \in [0, \alpha)$  satisfying

$$(3.4) \quad \begin{aligned} \varphi_\ell(\xi, n) = R_\ell^n \xi - \lambda_\ell \sum_{i=1}^n R_\ell^{n-i+1} P_{\ell 1} K_\ell F_\ell(\varphi_\ell(\xi, i)) \\ + \lambda_\ell \sum_{i=n+1}^\infty Q_\ell^{i-n-1} P_{\ell 2} K_\ell F_\ell(\varphi_\ell(\xi, i)) \end{aligned}$$

for  $\xi \in Y_{\ell 1}$  and  $n \geq 0$ . Here  $R_\ell = [C(\lambda_\ell)P_{\ell 1}]^{-1}$ ,  $Q_\ell = C(\lambda_\ell)P_{\ell 2}$  and  $c_\eta(\mathbb{N}, X_\ell)$  denotes the Banach space of bounded sequences  $\tilde{x} = \{x_n\}_{n \geq 0}$  in  $X_\ell$  with the norm  $\|\tilde{x}\|_\ell^{(\eta)} = \sup_{n \geq 0} e^{\eta n \lambda_\ell} \|x_n\|_\ell$ .

Since (2.6) is proved in [12], we shall show (2.7) only. Fix an arbitrary bounded set  $B \subset Y_1$  and set for  $y \in B$

$$(3.5) \quad \Omega_\ell(y) = \sup_{n \geq 0} \{ \sup_{t \in G_{\ell n}} e^{\eta n \lambda_\ell} \|D\varphi_\ell(V_\ell y, n)V_\ell - V_\ell Df(y, t)\|_{Y_1, X_\ell} \}$$

where  $G_{\ell n} = ((n-1)\lambda_\ell, n\lambda_\ell] \cap \mathbb{R}^+$ . Here  $Df(y, \cdot)$  denotes the Fréchet derivative of  $f$  at  $y \in Y_1$ , and so  $Df(y, t) \in B(Y_1, X)$ . Likewise,  $D\varphi_\ell(\xi, n) \in B(Y_{\ell 1}, X_\ell)$  for  $\xi \in Y_{\ell 1}$ . To prove (2.7) it suffices to show

$$(3.6) \quad \lim_{\ell \rightarrow \infty} \sup_{y \in B} \Omega_\ell(y) = 0.$$

For  $n \in \mathbb{N}, t \in G_{\ell n}$  and  $y \in B$  we write

$$D\varphi_\ell(V_\ell y, n)V_\ell - V_\ell Df(y, t) = \sum_{i=1}^8 H_i$$

with

$$\begin{aligned}
 H_1 &= R_\ell^n V_\ell - V_\ell S(-n\lambda_\ell) P_1, \\
 H_2 &= - \sum_{i=1}^n \int_{G_{\ell i}} R_\ell^{n-i+1} K_\ell DF_\ell(\varphi_\ell(V_\ell y, i)) \{D\varphi_\ell(V_\ell y, i) V_\ell - V_\ell Df(y, s)\} ds, \\
 H_3 &= - \sum_{i=1}^n \int_{G_{\ell i}} R_\ell^{n-i+1} K_\ell \{DF_\ell(\varphi_\ell(V_\ell y, i)) V_\ell - V_\ell DF(f(y, s))\} Df(y, s) ds, \\
 H_4 &= - \sum_{i=1}^n \int_{G_{\ell i}} \{R_\ell^{n-i+1} K_\ell V_\ell - V_\ell S(s - n\lambda_\ell) P_1\} DF(f(y, s)) Df(y, s) ds, \\
 H_5 &= \sum_{i=n+1}^\infty \int_{G_{\ell i}} Q_\ell^{i-n-1} P_{\ell 2} K_\ell DF_\ell(\varphi_\ell(V_\ell y, i)) \{D\varphi_\ell(V_\ell y, i) V_\ell - V_\ell Df(y, s)\} ds, \\
 H_6 &= \sum_{i=n+1}^\infty \int_{G_{\ell i}} Q_\ell^{i-n-1} P_{\ell 2} K_\ell \{DF_\ell(\varphi_\ell(V_\ell y, i)) V_\ell - V_\ell DF(f(y, s))\} Df(y, s) ds, \\
 H_7 &= \sum_{i=n+1}^\infty \int_{G_{\ell i}} \{Q_\ell^{i-n-1} P_{\ell 2} K_\ell V_\ell - V_\ell S(s - n\lambda_\ell) P_2\} DF(f(y, s)) Df(y, s) ds, \\
 H_8 &= V_\ell (Df(y, n\lambda_\ell) - Df(y, t)).
 \end{aligned}$$

By [12, Lemma 3.7] we have that for  $\varepsilon > 0, \alpha + \varepsilon\eta > 0, z \in Y_1$  and  $k, n \in \mathbb{N}$

$$e^{\eta n \lambda_\ell} \|H_1 z\|_\ell \leq C_{\varepsilon, k} \rho_1(\ell, z) + C \rho_2(k, z)$$

with the functions  $\rho_j(m, y), j = 1, 2$ , such that the family  $\{\rho_j(m, y)\}_{m \geq 1}$  is equicontinuous in  $y \in Y$  and  $\lim_{m \rightarrow \infty} \rho_j(m, y) = 0$  for each  $y$ . Here and in what follows,  $C$  denotes various constants and  $C_{\varepsilon, k}$  denotes a constant depending on  $\varepsilon$  and  $k$ . Set  $B_1 = \{z \in Y_1; |z| \leq 1\}$ . Since  $B_1$  is compact in  $Y_1$ , we have  $\lim_{m \rightarrow \infty} \sup_{z \in B_1} \rho_j(m, z) = 0$ , and hence

$$(3.7) \quad \lim_{\ell \rightarrow \infty} \sup_{n \geq 0} e^{\eta n \lambda_\ell} \|H_1\|_{Y_1, X_\ell} = 0.$$

By a similar way as in [12, p.176] we can compute

$$(3.8) \quad e^{\eta n \lambda_\ell} \|H_2\|_{Y_1, X_\ell} \leq e^{(\omega - \eta)\lambda_\ell} M M_1 M_2 L_F \alpha^{-1} \Omega_\ell(y).$$

For a fixed  $T > 0$  take  $T_\ell \in \mathbb{N}$  so that  $T/\lambda_\ell - 1 < T_\ell \leq T/\lambda_\ell$ . Set

$$d_{\ell, T} = \sup |\{DF_\ell(\varphi_\ell(V_\ell y, i)) V_\ell - V_\ell DF(f(y, s))\} Df(y, s) z|_\ell,$$

where the supremum is taken with respect to  $s, i, y$  and  $z$  satisfying  $s \in G_{\ell i}, 1 \leq i \leq T_\ell, y \in B$  and  $z \in B_1$ . Observing

$$\begin{aligned}
 &|\{DF_\ell(\varphi_\ell(V_\ell y, i)) V_\ell - V_\ell DF(f(y, s))\} Df(y, s) z|_\ell \\
 &\leq |\{DF_\ell(\varphi_\ell(V_\ell y, i)) - DF_\ell(V_\ell f(y, s))\} V_\ell Df(y, s) z|_\ell \\
 &\quad + |\{DF_\ell(V_\ell f(y, s)) V_\ell - V_\ell DF(f(y, s))\} Df(y, s) z|_\ell
 \end{aligned}$$

and  $\lim_{\ell \rightarrow \infty} \|\varphi_\ell(V_\ell y, i) - V_\ell f(y, s)\|_\ell = 0$  uniformly for  $y \in B, s \in G_{\ell i}$  and  $1 \leq i \leq T_\ell$  by (2.6) (also see [12, (4.3)]), we see that  $\lim_{\ell \rightarrow \infty} d_{\ell, T} = 0$  by (C8) because the sets  $\{f(y, s); y \in B, s \in [0, T]\}$  and  $\{Df(y, s)z; y \in B, z \in B_1, s \in [0, T]\}$  are

compact in  $X$  and  $Y$ , respectively. We also observe that  $\|V_\ell\|_{X, X_\ell}$  and  $\|V_\ell\|_{Y, Y_\ell}$  are bounded in  $\ell$  by the uniform boundedness principle and that

$$(3.9) \quad \begin{aligned} \|Df(y, s)\|_{Y_1, X} &\leq e^{-(\eta+\varepsilon)i\lambda_\ell} \|Df(y)\|_{Y_1, C_{\eta+\varepsilon}(\mathbb{R}^+, X)} \\ &\leq C e^{-(\eta+\varepsilon)i\lambda_\ell} \quad \text{for } s \in G_{\ell i} \text{ and } y \in B. \end{aligned}$$

Hence, by (C3)-(C7)  $\|H_3 z\|_\ell$  for  $z \in B_1$  is estimated by

$$C e^{-(\alpha+\eta)n\lambda_\ell} \left\{ d_{\ell, T} \sum_{i=1}^{T_\ell} \lambda_\ell e^{(\alpha+\eta)(i-1)\lambda_\ell} + L_F \sum_{i=T_\ell+1}^n e^{(\alpha+\eta)(i-1)\lambda_\ell} \int_{G_{\ell i}} \|Df(y, s)z\| ds \right\},$$

and so by (3.9)

$$(3.10) \quad e^{\eta n \lambda_\ell} \|H_3\|_{Y_1, X_\ell} \leq C(d_{\ell, T} + e^{-\varepsilon T}).$$

To estimate  $H_4$  we take  $z \in B_1$  and put  $w = DF(f(y, s))Df(y, s)z$ . By [12, Lemma 3.7] again  $e^{\eta n \lambda_\ell} \|H_4 z\|_\ell$  is estimated by

$$\begin{aligned} &e^{\eta n \lambda_\ell} \sum_{i=1}^{T_\ell} \int_{G_{\ell i}} e^{-(n-i+1)(\alpha+\eta)\lambda_\ell/(1-\varepsilon)} (C_{\varepsilon, k} \rho_1(\ell, w) + C \rho_2(k, w)) ds \\ &\quad + C e^{\eta n \lambda_\ell} \sum_{i=T_\ell+1}^n \int_{G_{\ell i}} e^{-(n\lambda_\ell+s)(\alpha+\eta)} L_F \|Df(y, s)z\| ds \\ &\leq e^{-(\alpha+\eta)T/(1-\varepsilon)} (C_{\varepsilon, k} \rho_1^*(\ell) + C \rho_2^*(k) + C e^{-\varepsilon T}). \end{aligned}$$

Here we set

$$\rho_j^*(m) = \sup\{\rho_j(m, w); y \in B, z \in B_1, s \in [0, T]\}, \quad j = 1, 2.$$

Since the set  $\{w; y \in B, z \in B_1, s \in [0, T]\}$  is compact in  $Y$ , it holds that  $\lim_{m \rightarrow \infty} \rho_j^*(m) = 0$ . Hence, letting  $\ell \rightarrow \infty, k \rightarrow \infty$  and  $T \rightarrow \infty$  in this order, we get

$$(3.11) \quad \lim_{\ell \rightarrow \infty} \sup_{n \geq 1, y \in B} e^{\eta n \lambda_\ell} \|H_4\|_{Y_1, X_\ell} = 0.$$

By a similar manner as in [12, p.178] we can compute

$$(3.12) \quad e^{\eta n \lambda_\ell} \|H_5\|_{Y_1, X_\ell} \leq e^{(\beta-\eta)\lambda_\ell} L_F \{M_4 \Gamma(1-\gamma) \beta^{\gamma-1} + M_5 \beta^{-1}\} \Omega_\ell(y).$$

By using (2.4) one can estimate  $\|H_6\|_{Y_1, X_\ell}$  by

$$\begin{aligned} &C \sum_{i=n+1}^{T_\ell} \lambda_\ell \{((i-n)\lambda_\ell)^{-\gamma} + 1\} e^{(\eta-\beta)(i-n-1)\lambda_\ell} d_{\ell, T} \\ &\quad + C \sum_{i=T_\ell+1}^\infty \int_{G_{\ell i}} \{((i-n)\lambda_\ell)^{-\gamma} + 1\} e^{(\eta-\beta)(i-n-1)\lambda_\ell} L_F \|Df(y, s)\|_{Y_1, X} ds. \end{aligned}$$

Hence, by (3.9)

$$(3.13) \quad e^{\eta n \lambda_\ell} \|H_6\|_{Y_1, X_\ell} \leq C(d_{\ell, T} + e^{-\varepsilon T}).$$

Next, by using [12, Lemma 3.10] we obtain for  $z \in B_1$  and  $v = DF(f(y, s))Df(y, s)z$

$$\begin{aligned} \|H_7 z\|_\ell &\leq \int_0^{T-n\lambda_\ell} (s^{-\gamma} + 1)e^{(1-\varepsilon)(\eta-\beta)s} (C_{\varepsilon,k}\sigma_1(\ell, v) + C\sigma_2(k, v)) ds \\ &\quad + C \int_{T-n\lambda_\ell}^\infty (s^{-\gamma} + 1)e^{(\eta-\beta)s} \|Df(y, s)z\| ds \end{aligned}$$

with the functions  $\sigma_j(m, w), j = 1, 2$ , such that the family  $\{\sigma_j(m, w)\}_{m \geq 1}$  is equicontinuous in  $w \in Y$  and  $\lim_{m \rightarrow \infty} \sigma_j(m, w) = 0$  for each  $w$ . Hence, by (3.9)

$$(3.14) \quad e^{nm\lambda_\ell} \|H_7 z\|_{Y_1, X_\ell} \leq C_{\varepsilon,k}\sigma_1^*(\ell) + C\sigma_2^*(k) + Ce^{-\varepsilon T}.$$

Here we set

$$\sigma_j^*(m) = \sup\{\sigma_j(m, v); y \in B, z \in B_1, s \in [0, T]\}, \quad j = 1, 2.$$

Just as in the case of  $\rho_j^*$  we see that  $\lim_{m \rightarrow \infty} \sigma_j^*(m) = 0$ .

Finally, set

$$\delta_T(h) = \sup \|Df(y, s)z - Df(y, \hat{s})z\|$$

where the supremum is taken over all  $y \in B, z \in B_1$  and  $s, \hat{s} \in [0, 2T]$  with  $|s - \hat{s}| \leq h$ . It is easy to see that  $\lim_{h \downarrow 0} \delta_T(h) = 0$ . A similar computation as in [12, p.179] yields that for  $z \in B_1, y \in B$

$$(3.15) \quad e^{\eta n\lambda_\ell} \|H_8 z\|_\ell \leq C(\delta_T(\lambda_\ell) + e^{-\varepsilon T} \|Df(y)\|_{Y_1, C_{\eta+\varepsilon}(\mathbb{R}^+, X)}).$$

We are now in a position to prove (3.6). By virtue of (3.7), (3.8) and (3.10)-(3.15) we obtain

$$\overline{\lim}_{\ell \rightarrow \infty} \sup_{y \in B} \Omega_\ell(y) \leq K(\alpha, \beta) L_F \overline{\lim}_{\ell \rightarrow \infty} \sup_{y \in B} \Omega_\ell(y).$$

Since  $K(\alpha, \beta)L_F < 1$  by (G), we conclude that (3.6) holds.

#### 4. EXAMPLE

We briefly consider the renormalized Kuramoto-Sivashinsky equation (KSE)

$$u_t + D^4 u + D^2 u + uDu = 0, \quad (x, t) \in \mathbb{R} \times \mathbb{R}^+,$$

subject to periodic boundary condition, with period  $L$ . We refer to [4], [6], [12] in the notation and some results concerning the KSE. We view it as an evolution equation in the Hilbert space  $Y = \{u \in L^2_{per}(0, L); u \text{ is odd}\}$  with the usual  $L^2$  norm. Since the KSE in  $Y$  has a bounded absorbing set in  $X = H^2_{per}(0, L) \cap Y$ , i.e., there exists a constant  $r_0 > 0$  such that for every  $r > 0$  we can choose a time  $T^*(r) > 0$  satisfying  $\|u(t)\|_{H^2} \leq r_0$  for all  $t \geq T^*(r)$  and all  $u(0) \in Y$  with  $\|u(0)\|_{L^2} \leq r$ , we may consider the prepared equation instead of the KSE

$$(4.1) \quad du/dt = Au + Fu, \quad t \in \mathbb{R}^+,$$

where  $Au = -D^4 u$  and  $Fu = -D^2 u - \rho(\|u\|_{H^2})uDu$  with a smooth function  $\rho$  satisfying  $0 \leq \rho \leq 1$  and  $\rho(r) = 1$  if  $|r| \leq r_0, \rho(r) = 0$  if  $|r| \geq 2r_0$ .

Let  $Y_\ell$  be the Hilbert space  $S^{\ell}_{odd,per}$ , the set of vectors  $\xi = (\xi_0, \dots, \xi_{\ell-1})$  satisfying  $\xi_0 = 0$  and  $\xi_j = \xi_{\ell-j}$  for  $j = 1, \dots, \ell - 1$ , with norm  $|\xi|_\ell = (h \sum_{k=1}^{\ell-1} \xi_k^2)^{1/2}, h = L/\ell$ . For convenience  $\xi$  will be extended periodically to an infinite sequence by  $\xi_{j+\ell} = \xi_j$ . Define  $\Delta_\ell : Y_\ell \rightarrow Y_\ell$  by

$$(\Delta_\ell \xi)_k = h^{-2}(\xi_{k-1} - 2\xi_k + \xi_{k+1}) \quad \text{for } \xi \in Y_\ell,$$

where  $(\Delta_\ell \xi)_k$  denotes the  $k$ -th element of  $\Delta_\ell \xi$ . We also consider  $S_{odd,per}^\ell$  as a Hilbert space with norm  $\|\xi\|_\ell = |\Delta_\ell \xi|_\ell$ , which is denoted by  $X_\ell$ .

We approximate (4.1) by the finite difference scheme

$$(4.2) \quad \lambda_\ell^{-1}(\xi^{i+1} - \xi^i) + \Delta_\ell^2((1 - \theta)\xi^i + \theta\xi^{i+1}) - F_\ell(\xi^i) = 0, \quad \xi^i \in Y_\ell, \quad i \in \mathbb{N}$$

( $1/2 < \theta \leq 1$ ), where  $F_\ell(\xi) = -\Delta_\ell \xi - \rho(\|\xi\|_\ell^2)B^\ell(\xi)$  and  $B^\ell : Y_\ell \times Y_\ell \rightarrow Y_\ell$  is defined by

$$(B^\ell(\xi))_k = (6h)^{-1}(\xi_{k-1} + \xi_k + \xi_{k+1})(\xi_{k+1} - \xi_{k-1}) \quad \text{for } \xi \in Y_\ell.$$

Then, (4.2) is rewritten as the form (1.2) by setting

$$C(\lambda_\ell) = (I - (1 - \theta)\lambda_\ell \Delta_\ell^2)(I + \theta\lambda_\ell \Delta_\ell^2)^{-1} \text{ and } K_\ell = (I + \theta\lambda_\ell \Delta_\ell^2)^{-1}.$$

By some operational calculi and spectral theorems one finds that conditions (C1)-(C8) hold with  $M = M_2 = M_3 = 1, M_4 = 2, \gamma = 1/2, \omega = 0, M_1 = \nu_N^{1/2}, M_5 = (2\nu_{N+1})^{1/2}, \alpha = \beta = (\nu_{N+1} - \nu_N)/4, \eta = -(\nu_{N+1} + \nu_N)/2$  for  $N \in \mathbb{N}$ . Here we set  $\nu_k = (2\pi k/L)^4$ , the eigenvalues of the operator  $A$  in  $Y$ . Hence, if  $N$  is large, then condition (G) is satisfied and consequently we can apply our theorem.

#### REFERENCES

1. S.N. Chow and K. Lu, *Invariant manifolds for flows in Banach spaces*, J. Differential Equations **74** (1988), 285–317. MR **89h**:58163
2. S.N. Chow, K. Lu and G.R. Sell, *Smoothness of inertial manifolds*, J. Math. Anal. Appl. **169** (1992), 283–321. MR **94a**:58172
3. F. Demengel and J.M. Ghidaglia, *Inertial manifolds for partial differential equations under time-discretization: Existence and applications*, J. Math. Anal. Appl. **155** (1991), 177–225. MR **92e**:35079
4. C. Foias, M.S. Jolly, I.G. Kevrekidis and E.S. Titi, *Dissipativity of numerical schemes*, Nonlinearity **4** (1991), 591–613. MR **92f**:65099
5. C. Foias, G. Sell and R. Temam, *Inertial manifolds for nonlinear evolutionary equations*, J. Differential Equations **73** (1988), 309–353.
6. C. Foias and E.S. Titi, *Determining nodes, finite difference schemes and inertial manifolds*, Nonlinearity **4** (1991), 135–153. MR **92a**:65241
7. E. Hille and R.S. Phillips, *Functional Analysis and Semi-Groups*, Amer. Math. Soc., Providence, RI, 1957. MR **54**:11077; MR **19**:664d
8. M.S. Hirsch, C.C. Pugh and M. Shub, *Invariant Manifolds*, Lecture Notes in Mathematics Vol.583, Springer-Verlag, New York, 1977. MR **58**:18595
9. D.A. Jones and A.M. Stuart, *Attractive invariant manifolds under approximation. Inertial manifolds*, J. Differential Equations **123** (1995), 588–637. MR **96j**:34110
10. D.A. Jones and E.S. Titi,  *$C^1$  approximation of inertial manifolds for dissipative nonlinear equations*, J. Differential Equations **127** (1996), 54–86. MR **97g**:58154
11. K. Kobayasi, *Convergence and approximation of inertial manifolds for evolution equations*, Differential and Integral Equations **8** (1995), 1117–1134. MR **96c**:34131
12. K. Kobayasi, *Inertial manifolds for discrete approximations of evolution equations: Convergence and approximations*, Advances in Math. Sci. Appl. **3** (1993/1994), 161–189. MR **95d**:34107
13. A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, Applied Mathematical Sciences 44, Springer-Verlag, New York, 1983. MR **85g**:47061
14. V.A. Pliss and G.R. Sell, *Perturbations of attractors of differential equations*, J. Differential Equations **92** (1991), 100–124. MR **92f**:58094

DEPARTMENT OF MATHEMATICS, SCHOOL OF EDUCATION, WASEDA UNIVERSITY, 1-6-1 NISHI-WASEDA, SHINJUKU-KU, TOKYO 169-8050, JAPAN  
*E-mail address*: kzokoba@mn.waseda.ac.jp