

THE GROWTH THEOREM OF CONVEX MAPPINGS ON THE UNIT BALL IN \mathbb{C}^n

HIDETAKA HAMADA

(Communicated by Steven R. Bell)

ABSTRACT. Let $\|\cdot\|$ be an arbitrary norm on \mathbb{C}^n . Let f be a normalized biholomorphic convex mapping on the unit ball in \mathbb{C}^n with respect to the norm $\|\cdot\|$. We will give an upper bound of the growth of f .

Let Ω be a domain in \mathbb{C}^n which contains the origin in \mathbb{C}^n . A holomorphic mapping f from Ω to \mathbb{C}^n is said to be normalized, if $f(0) = 0$ and the Jacobian matrix at the origin $Df(0)$ is the identity matrix. Let \mathbb{B}^n denote the Euclidean unit ball in \mathbb{C}^n . Let $f(z)$ be a normalized biholomorphic convex mapping on \mathbb{B}^n . Then FitzGerald and Thomas [2], Liu [6] and Suffridge [7] independently used different methods to prove the following growth theorem.

$$\frac{|z|}{1+|z|} \leq |f(z)| \leq \frac{|z|}{1-|z|},$$

where $|\cdot|$ denotes the Euclidean norm. Let

$$B_p = \left\{ z \in \mathbb{C}^n; \|z\|_p = \left(\sum_{i=1}^n |z_i|^p \right)^{1/p} < 1 \right\}$$

for $p \geq 1$. Gong and Liu [4] gave the following upper bound of the growth of normalized biholomorphic convex mappings on B_p .

$$\|f(z)\|_p \leq \frac{\|z\|_p}{1 - \|z\|_p}.$$

They also obtained an upper bound of the growth of normalized biholomorphic convex mappings on the convex complex ellipsoid

$$D(p_1, \dots, p_n) = \{z \in \mathbb{C}^n; |z_1|^{p_1} + \dots + |z_n|^{p_n} < 1\}$$

with $p_1, \dots, p_n \geq 1$.

Let $\|\cdot\|$ be an arbitrary norm on \mathbb{C}^n and let \mathbb{B} denote the unit ball in \mathbb{C}^n with respect to the norm $\|\cdot\|$. Using the idea of FitzGerald and Thomas [2], we obtain the following theorem.

Received by the editors June 3, 1997 and, in revised form, July 16, 1997.

1991 *Mathematics Subject Classification*. Primary 32H02; Secondary 30C45.

©1999 American Mathematical Society

Theorem. Let $f(z)$ be a normalized biholomorphic convex mapping from \mathbb{B} to \mathbb{C}^n . Then

$$\|f(z)\| \leq \frac{\|z\|}{1 - \|z\|}.$$

Proof. Let Δ be the unit disc in \mathbb{C} . For any fixed $w \in \partial\mathbb{B}$ and $\zeta \in \Delta$, let

$$f(\zeta w) = \zeta w + \sum_{i=2}^{\infty} d_i \zeta^i.$$

Since f is a holomorphic mapping into \mathbb{C}^n , $d_i \in \mathbb{C}^n$. Let $m \geq 2$, $m \in \mathbb{Z}$ be fixed. Let $\varepsilon = \exp(2\pi i/m)$. Then

$$\sum_{k=0}^{m-1} f(\zeta^{1/m} \varepsilon^k w) = m \sum_{i=1}^{\infty} d_{mi} \zeta^i$$

is holomorphic with respect to $\zeta \in \Delta$. Since $f(\mathbb{B})$ is convex,

$$h(\zeta) = f^{-1} \left(\frac{1}{m} \sum_{k=0}^{m-1} f(\zeta^{1/m} \varepsilon^k w) \right)$$

is well-defined and holomorphic on Δ . Since f is normalized,

$$f^{-1}(z) = z + O(|z|^2).$$

Therefore, $h(\zeta) = d_m \zeta + O(|\zeta|^2)$. Since $h(\Delta) \subset \mathbb{B}$, we obtain $\|d_m\| \leq (1 - \delta)^{-1}$ by applying the maximum modulus theorem with values in a complex Banach space (cf. Dunford and Schwartz [1]) to the holomorphic mapping $h(\zeta)/\zeta$ on $|\zeta| < 1 - \delta$. Letting δ tend to 0, we have $\|d_m\| \leq 1$. Then we have

$$\|f(\zeta w)\| \leq |\zeta| + \sum_{i=2}^{\infty} |\zeta|^i = \frac{|\zeta|}{1 - |\zeta|} = \frac{\|\zeta w\|}{1 - \|\zeta w\|}.$$

□

Let D be a bounded convex balanced domain in \mathbb{C}^n . Then the Minkowski function of D is a norm on \mathbb{C}^n and D is the unit ball with respect to the norm (cf. Jarnicki and Pflug [5]). Then the above theorem holds for D . In particular, the theorem gives another growth theorem of convex mappings on $D(p_1, \dots, p_n)$ with $p_1, \dots, p_n \geq 1$.

The author would like to thank the referee for his helpful comments and suggestions.

REFERENCES

1. N. Dunford and J. Schwartz, *Linear operators*, vol. 1, Interscience, New York, 1958. MR **22**:8302
2. C. H. FitzGerald and C. R. Thomas, *Some bounds on convex mappings in several complex variables*, Pacific J. Math. **165** (1994), 295–320. MR **95k**:32021
3. S. Gong, *Biholomorphic mappings in several complex variables*, Contemporary Math. **142** (1993), 15–48. MR **94a**:32035
4. S. Gong and T. Liu, *The growth theorem of biholomorphic convex mappings on \mathbb{B}^p* , Chin. Quar. Jour. Math. **6** (1991), 78–82.
5. M. Jarnicki and P. Pflug, *Invariant distances and metrics in complex analysis*, de Gruyter, Berlin-New York, 1993. MR **94k**:32039

6. T. Liu, *The growth theorems, covering theorems and distortion theorems for biholomorphic mappings on classical domains*, University of Science and Technology of China Thesis (1989).
7. T. J. Suffridge, *Biholomorphic mappings of the ball onto convex domains*, Abstracts of papers presented to American Mathematical Society **11(66)** (1990), 46.

FACULTY OF ENGINEERING, KYUSHU KYORITSU UNIVERSITY, 1-8, JIYUGAOKA, YAHATANISHI-KU,
KITAKYUSHU 807, JAPAN

E-mail address: hamada@kyukyo-u.ac.jp