

TORIC DEGENERATIONS AND VECTOR BUNDLES

JOSEPH GUBELADZE

(Communicated by Wolmer V. Vasconcelos)

ABSTRACT. There are many affine subalgebras of polynomial rings with highly non-trivial projective modules, whose initial algebras (toric degenerations) are still finitely generated and have all projective modules free.

Let $k[X_1, \dots, X_n]$ be a polynomial algebra (k a field, $n \in \mathbb{N}$) and A an affine k -subalgebra. Let \prec denote a term order on the multiplicative semigroup of monomials in the X_i and let $\text{in}_{\prec}(A)$ denote the monomial subalgebra of $k[X_1, \dots, X_n]$, generated by the leading monomials of elements $f \in A$ with respect to \prec . In case the *initial algebra* $\text{in}_{\prec}(A)$ is finitely generated, one can obtain many properties of A by checking them for $\text{in}_{\prec}(A)$ (called sometimes a *toric degeneration* of A) (see [CHV], [RS]). However, this is not the case for the property ‘all projective modules are free’ – thanks to Bernd Sturmfels for asking me this question.

Theorem 1. *Let $A = k[X, Y, Z^2, Z^3 - XYZ]$ and \prec be the lexicographic term order corresponding to $Z \prec Y \prec X$. Then $SK_0(A) = \text{Ker}(\tilde{K}_0(A) \xrightarrow{\det} \text{Pic}(A))$ is not trivial (equivalently, there are projective A -modules which are not even stably of type $\text{free} \oplus \text{rank } 1$), while $\text{in}_{\prec}(A)$ is finitely generated and all projective $\text{in}_{\prec}(A)$ -modules are free.*

(Here ‘projective’ includes ‘finitely generated’.)

Proof. Since $(Z^2 - XY)k[X, Y, Z] \subset A$, the following diagram with the upper horizontal identity embedding is a pull-back diagram (all the letters refer to variables)

$$\begin{array}{ccc} A & \longrightarrow & k[X, Y, Z] \\ \downarrow & & \downarrow \\ k[U, V] & \longrightarrow & k[S^2, ST, T^2], \end{array}$$

where $X \mapsto S^2$, $Y \mapsto T^2$, $Z \mapsto ST$, $U \mapsto S^2$, $V \mapsto T^2$. It is similarly easy to show that $\text{in}_{\prec}(A) = k[X, Y, Z^2, XYZ]$ – a seminormal monomial algebra (as a k -vector space it is generated by the normal monomial subalgebras $k[X, Y]$, $k[X, Z^2]$, $k[Y, Z^2]$ and $k[\{X^a Y^b Z^c \mid a > 0, b > 0, c > 0\}]$). So by [Gu1] projective modules over $\text{in}_{\prec}(A)$ are free, while the Mayer-Vietoris sequence (see [Bass], p.490), applied to the diagram above, implies $SK_0(A) = SK_1(k[S^2, ST, T^2])$. Hence, by [Gu2] $SK_0(A) \neq 0$. \square

Received by the editors February 20, 1998.

1991 *Mathematics Subject Classification.* Primary 13D15, 19A49.

This research was supported in part by the Alexander von Humboldt Foundation and CRDF grant #GM1-115.

Remark 2. Actually, by the (U – Pic)–Mayer-Vietoris sequence $\text{Pic}(A) = 0$ for the ring A in Theorem 1.

By [Gu2] essentially all (and conjecturely all) monomial algebras, not isomorphic to polynomial rings, have non-trivial SK_1 -groups. So changing the lower right algebra in the diagram above by other monomial algebras and changing correspondingly the ‘side’ polynomial algebras, one can produce a big class of rings having this property. One uses the general observation that if some powers of all variables belong to A , both A itself and its initial algebra $\text{in}_{\prec}(A)$ are finitely generated because they are intermediate algebras of the module finite extensions

$$k[X_1^{a_1}, \dots, X_n^{a_n}] \subset k[X_1, \dots, X_n].$$

Nontriviality of SK_1 for the special monomial ring $k[S^2, ST, T^2]$ (over a characteristic zero field k) was also shown in [Sri].

REFERENCES

- [Bass] H. Bass, *Algebraic K-theory*, W. A. Benjamin, Inc., 1968. MR **40**:2736
- [CHV] A. Conca, J. Herzog, G. Valla, *SAGBI bases with applications to blow-up algebras*, J. Reine Angew. Math. **474** (1996), 113-138. MR **97h**:13023
- [Gu1] J. Gubeladze, *Anderson’s conjecture and the maximal monoid class, over which projective modules are free*, Math. USSR Sbornik **63** (1989), 165–180. MR **89d**:13010
- [Gu2] J. Gubeladze, *Nontriviality of $SK_1(R[M])$* , J. Pure Appl. Alg. **104** (1995), 169-190. MR **96j**:19002
- [RS] L. Robbiano and M. Sweedler, *Subalgebra bases*, in Commutative algebra, Proc. Workshop, Salvador/Brasil 1988, Lect. Notes Math. **1430** (1990), 61-87. MR **91f**:13027
- [Sri] V. Srinivas, *K_1 of the cone over a curve*, J. Reine Angew. Math. **381** (1987), 37-50. MR **89e**:14008

A. RAZMADZE MATHEMATICAL INSTITUTE, ALEXIDZE ST. 1, 380093 TBILISI, GEORGIA
E-mail address: gubel@rmi.acnet.ge