

TYPE II_∞ FACTORS GENERATED BY PURELY INFINITE SIMPLE C^* -ALGEBRAS ASSOCIATED WITH FREE GROUPS

WOJCIECH SZYMAŃSKI AND SHUANG ZHANG

(Communicated by David R. Larson)

ABSTRACT. Let $\Gamma = G_1 * G_2 * \dots * G_n * \dots$ be a free product of at least two but at most countably many cyclic groups. With each such group Γ we associate a family of C^* -algebras, denoted $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ and generated by the reduced group C^* -algebra $C_r^*\Gamma$ and a collection \mathcal{P}_Λ of projections onto the ℓ^2 -spaces over certain subsets of Γ . We determine $W^*(\Gamma, \mathcal{P}_\Lambda)$, the weak closure of $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ in $\mathcal{L}(\ell^2(\Gamma))$, and use this result to show that many of the C^* -algebras in question are non-nuclear.

0. INTRODUCTION

In recent years a great deal of progress has been made towards the classification of separable, simple C^* -algebras. This program, initiated by George Elliott [6] and Mikael Rørdam [14], has been advanced significantly through combined efforts of a number of researchers. The progress has been particularly visible in the subcategory of unital, purely infinite and nuclear C^* -algebras; e.g. see [7], [9], [13]. And yet a number of important related problems still remain open. One such outstanding problem is whether there exists a simple C^* -algebra containing both finite and infinite projections. Partially motivated by the desire to find an example of such an algebra the second named author initiated investigations of a class of C^* -algebras related to free-product groups [17], [18].

Let $\Gamma = G_1 * G_2 * \dots * G_n * \dots$ be a free product of finitely or countably many cyclic groups. For each i we fix a generator g_i of G_i . For any $h \in \Gamma \setminus \{e\}$ let $\Gamma(h)$ be the set of reduced words, in the fixed set $\{g_i\}$ of generators of Γ , whose initial segments coincide with h . Let P_h denote the orthogonal projection from $\ell^2(\Gamma)$ onto $\ell^2(\Gamma(h))$, and let $C_r^*(\Gamma, P_h)$ be the C^* -subalgebra of $\mathcal{L}(\ell^2(\Gamma))$ generated by the reduced group C^* -algebra $C_r^*\Gamma$ and the projection P_h [17], [18]. This construction is somewhat similar, and actually related to those of [2], [12], [15].

Against the hope of finding a counterexample to the outstanding problem mentioned above, the second named author showed [17], [18] that all C^* -algebras of the form $C_r^*(\Gamma, P_h)$ are either purely infinite (see [5] for the definition) and simple, or extensions of purely infinite, simple C^* -algebras by the compacts. This result brought several natural questions about the finer structure of these algebras. The

Received by the editors April 27, 1998.

1991 *Mathematics Subject Classification*. Primary 46L05.

This research was partially supported by NSF grant DMS - 9225076.

most important one is whether the algebras are nuclear, and thus fit into the classifiable category, or not. In this article we provide a partial answer to this question. Our investigations are carried out for a larger class of C^* -algebras, containing those considered in [17], [18].

Let Λ be a subset of Γ and denote $\mathcal{P}_\Lambda = \{P_h : h \in \Lambda\}$. We define $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ as the C^* -subalgebra of $\mathcal{L}(\ell^2(\Gamma))$ generated by $C_r^*\Gamma$ and \mathcal{P}_Λ . Let Γ_Λ be the subgroup of Γ generated by those of $\{g_i^{\pm 1}\}$ which do not occur as final letters of reduced forms of elements in Λ . We denote by $W^*(\Gamma, \mathcal{P}_\Lambda)$ the weak closure of $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ in $\mathcal{L}(\ell^2(\Gamma))$.

The main result of this article, Theorem I, states that there is an isomorphism

$$W^*(\Gamma, \mathcal{P}_\Lambda) \cong W^*(\Gamma_\Lambda) \otimes \mathcal{L}(\mathcal{H}),$$

where \mathcal{H} is an infinite dimensional, separable Hilbert space, and $W^*(\Gamma_\Lambda)$ denotes the standard group von Neumann algebra. Since the subgroups Γ_Λ themselves are isomorphic to free-products of cyclic groups, all of them, with the exceptions of $\Gamma_\Lambda \cong \mathbb{Z}, \mathbb{Z}_k, \mathbb{Z}_2 * \mathbb{Z}_2$, are non-amenable. We are then able to conclude that for non-amenable Γ_Λ the von Neumann algebras $W^*(\Gamma, \mathcal{P}_\Lambda)$ are not injective and, consequently, the corresponding C^* -algebras $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ are non-nuclear (Theorem II). Thus, our construction yields a large class of non-nuclear, purely infinite, simple C^* -algebras, which are not classifiable by the existing K -theoretic invariants. It seems plausible that these algebras may become helpful examples when the classification program extends beyond the class of nuclear C^* -algebras. Having this ultimate objective in mind we carry out a detailed investigation, in a separate paper [16], of the structure of the algebras $C_r^*(\Gamma, \mathcal{P}_\Lambda)$.

1. MAIN RESULTS

1.0. Throughout this article Γ denotes a free product of finitely or countably many cyclic groups G_i . Let e denote the unit of Γ and let

$$\mathcal{G} = \{g_1, g_2, \dots, g_n, \dots\}$$

be a fixed set of generators of Γ , where g_i is a generator of G_i .

If g_i has finite order m , then we make a convention that elements of G_i be written as $e, g_i, g_i^2, \dots, g_i^{m-1}$. Then each element of Γ can be uniquely written as a reduced word

$$g_{i_1}^{\epsilon_1} g_{i_2}^{\epsilon_2} \dots g_{i_k}^{\epsilon_k}, \quad \epsilon_j = +1 \text{ or } -1.$$

A word is called *reduced* if it does not contain a subword of the form hh^{-1} , $h \in \Gamma$.

Let $\{\xi_h : h \in \Gamma\}$ be the standard basis of the Hilbert space $\ell^2(\Gamma)$, where $\xi_h(s) = 1$ if $h = s$ and $\xi_h(s) = 0$ if $h \neq s$. Let $L : \Gamma \rightarrow \mathcal{L}(\ell^2(\Gamma))$ be the left regular representation, i.e. $L_h(\xi_s) = \xi_{hs}$ for $h, s \in \Gamma$. The reduced group C^* -algebra $C_r^*\Gamma$ is generated by $\{L_h : h \in \Gamma\}$, and the group von Neumann algebra $W^*(\Gamma)$ is the weak closure of $C_r^*\Gamma$ [8, Vol. II, 6.7].

For a reduced word $h \in \Gamma \setminus \{e\}$ let $\Gamma(h)$ be the set of all reduced words in Γ whose initial segments coincide with h . Let P_h be the orthogonal projection from $\ell^2(\Gamma)$ onto the subspace $\ell^2(\Gamma(h))$. For a subset Λ of Γ let \mathcal{P}_Λ denote the set of projections $\{P_h : h \in \Lambda\}$. We define $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ as the C^* -subalgebra of $\mathcal{L}(\ell^2(\Gamma))$ generated by $C_r^*\Gamma$ and \mathcal{P}_Λ . We denote by $W^*(\Gamma, \mathcal{P}_\Lambda)$ the weak closure of $C_r^*(\Gamma, \mathcal{P}_\Lambda)$.

It is shown in [17, 1.1] that if h is a reduced word ending with $g_i^{\pm 1}$, then

$$C_r^*(\Gamma, P_h) = C_r^*(\Gamma, P_{g_i}) = C_r^*(\Gamma, P_{g_i^{-1}}).$$

Similarly, $P_h \in C_r^*(\Gamma, \mathcal{P}_\Lambda)$ if and only if $P_{g_i} \in C_r^*(\Gamma, \mathcal{P}_\Lambda)$. Thus, it suffices to consider the case when Λ is a subset of \mathcal{G} , and from now on we assume that this is indeed the case.

Let $\Lambda \subseteq \mathcal{G}$. We define Γ_Λ as the subgroup of Γ generated by $\mathcal{G} \setminus \Lambda$. It turns out that the subgroup Γ_Λ plays an important role in determining the structures of $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ and $W^*(\Gamma, \mathcal{P}_\Lambda)$. We denote by \mathcal{H} an infinite dimensional, separable Hilbert space.

1.1. Theorem I. *There exists a $*$ -isomorphism*

$$W^*(\Gamma, \mathcal{P}_\Lambda) \cong W^*(\Gamma_\Lambda) \otimes \mathcal{L}(\mathcal{H}).$$

Furthermore, the commutant $W^*(\Gamma, \mathcal{P}_\Lambda)'$ is $*$ -isomorphic to $W^*(\Gamma_\Lambda)$.

In order to prove Theorem I we need two lemmas. Let $R : \Gamma \rightarrow \mathcal{L}(\ell^2(\Gamma))$ be the right regular representation, i.e. $R_h(\xi_s) = \xi_{sh^{-1}}$ for $h, s \in \Gamma$. If $f \in \ell^2(\Gamma)$ and $R_f \in \mathcal{L}(\ell^2(\Gamma))$, then

$$R_f(\xi_s) = \sum_{h \in \Gamma} f(h)\xi_{sh^{-1}} \quad \forall s \in \Gamma.$$

1.2. Lemma. *Let $f \in \ell^2(\Gamma)$ and $R_f \in \mathcal{L}(\ell^2(\Gamma))$ be self-adjoint. Let $g \in \mathcal{G}$ (i.e. there exists i such that $g = g_i$). Then $R_f P_g = P_g R_f$ if and only if $\{s \in \Gamma : f(s) \neq 0\} \subseteq \Gamma_{\{g\}}$.*

Proof. Clearly, $R_f = R_f^*$ if and only if $f(s) = \overline{f(s^{-1})}$ for all $s \in \Gamma$. Also, $R_f P_g = P_g R_f$ if and only if the subspace $\ell^2(\Gamma(g))$ is R_f -invariant (i.e. $R_f(\ell^2(\Gamma(g))) \subseteq \ell^2(\Gamma(g))$), i.e. $R_f(\xi_h) \in \ell^2(\Gamma(g))$ for all $h \in \Gamma(g)$. This is equivalent to

$$\{\Gamma(g)s^{-1}\} \cup \{\Gamma(g)s\} \subseteq \Gamma(g) \quad \text{whenever } f(s) \neq 0.$$

Notice that $f(s) \neq 0$ if and only if $f(s^{-1}) \neq 0$. Thus, if $R_f P_g = P_g R_f$, then the function f must satisfy

$$\{s \in \Gamma : f(s) \neq 0\} \subseteq \{s \in \Gamma : \{\Gamma(h)s^{-1}\} \cup \{\Gamma(g)s\} \subseteq \Gamma(g)\}.$$

We claim that $\{\Gamma(g)s^{-1}\} \cup \{\Gamma(g)s\} \subseteq \Gamma(g)$ if and only if neither g nor g^{-1} is contained as a factor in the reduced word s . In order to prove this claim we consider separately two cases:

- (1) g has infinite order,
- (2) g has finite order m .

Case (1). Suppose that s contains a factor g or g^{-1} , i.e. in reduced form

$$s = g_{i_1}^{\epsilon_1} \dots g_{i_r}^{\epsilon_r} g_{i_{r+1}}^{\epsilon_{r+1}} \dots g_{i_k}^{\epsilon_k},$$

where $\epsilon, \epsilon_i = 1$ or -1 . If $\epsilon = 1$, set $h = g g_{i_{r+1}}^{\epsilon_{r+1}} \dots g_{i_k}^{\epsilon_k}$. Since $g_{i_{r+1}}^{\epsilon_{r+1}} \neq g^{-1}$, we have $h \in \Gamma(g)$. Then $hs^{-1} \notin \Gamma(g)$, because $g_{i_r}^{\epsilon_r} \neq g^{-1}$. If $\epsilon = -1$, set $h = g(g_{i_r}^{\epsilon_r})^{-1}(g_{i_{r-1}}^{\epsilon_{r-1}})^{-1} \dots (g_{i_1}^{\epsilon_1})^{-1}$, again an element of $\Gamma(g)$. Also $hs \notin \Gamma(g)$, due to reasoning as above. Thus, $\{\Gamma(g)s\} \cup \{\Gamma(g)s^{-1}\} \not\subseteq \Gamma(g)$.

Conversely, suppose that the reduced form of s contains neither g nor g^{-1} as a factor, and let $h \in \Gamma(g)$. Then the initial letter g of h survives all cancellations in hs (or hs^{-1}) and, thus, $\{\Gamma(g)s\} \cup \{\Gamma(g)s^{-1}\} \subseteq \Gamma(g)$.

Case (2). By our convention, all elements of the subgroup G_i ($g = g_i$) are written as $e, g, g^2, \dots, g^{m-1}$. Suppose first that the reduced form of s contains g as a factor, i.e.

$$s = g_{i_1}^{\epsilon_1} \dots g_{i_r}^{\epsilon_r} g^n g_{i_{r+1}}^{\epsilon_{r+1}} \dots g_{i_k}^{\epsilon_k},$$

where $1 \leq n \leq m-1$, and g_{i_r} and $g_{i_{r+1}}$ are different from g . Set

$$h = g^{m-n} (g_{i_r}^{\epsilon_r})^{-1} (g_{i_{r-1}}^{\epsilon_{r-1}})^{-1} \dots (g_{i_1}^{\epsilon_1})^{-1}.$$

Then $hs \in \Gamma(g_{i_{r+1}}^{\epsilon_{r+1}})$ and, hence, $hs \notin \Gamma(g)$.

Conversely, if s does not contain a factor g , then hs, hs^{-1} are in $\Gamma(g)$ for any $h \in \Gamma(g)$, i.e. $\{\Gamma(g)s\} \cup \{\Gamma(g)s^{-1}\} \subseteq \Gamma(g)$.

Consequently, $R_f P_g = P_g R_f$ if and only if $\{s \in \Gamma : f(s) \neq 0\} \subseteq \Gamma_{\{g\}}$, as desired. \square

1.3. Lemma. $W^*(\Gamma, \mathcal{P}_\Lambda)' \cong W^*(\Gamma_\Lambda)$.

Proof. It is obvious that $W^*(\Gamma, \mathcal{P}_\Lambda)' = W^*(\Gamma)' \cap (\bigcap_{h \in \Lambda} \{P_h\}')$. By the classical theory [8, Vol. II, 6.7.2], the commutant $W^*(\Gamma)'$ in $\mathcal{L}(\ell^2(\Gamma))$ coincides with the von Neumann algebra generated by $\{R_s : s \in \Gamma\}$. Also, each element of $W^*(\Gamma)'$ can be written in the form R_f , for some $f \in \ell^2(\Gamma)$. It is immediate that

$$W^*(\Gamma, \mathcal{P}_\Lambda)' = \{R_f \in \mathcal{L}(\ell^2(\Gamma)) : R_f P_h = P_h R_f \quad \forall h \in \Lambda\}.$$

Let us consider the self-adjoint part of $W^*(\Gamma, \mathcal{P}_\Lambda)'$, which of course generates the whole $W^*(\Gamma, \mathcal{P}_\Lambda)'$. If $R_f = R_f^*$ and $R_f \in W^*(\Gamma, \mathcal{P}_\Lambda)'$, then by Lemma 1.2

$$\{s \in \Gamma : f(s) \neq 0\} \subseteq \Gamma_\Lambda.$$

It follows that $W^*(\Gamma, \mathcal{P}_\Lambda)'$ equals the von Neumann algebra generated by the unitaries $\{R_s : s \in \Gamma_\Lambda\}$. Therefore, $W^*(\Gamma, \mathcal{P}_\Lambda)'$ is $*$ -isomorphic to $W^*(\Gamma_\Lambda)$ [8, Vol. II, 6.7]. \square

1.4. Proof of Theorem I. We write $\tilde{\Gamma}$ for a fixed set of representatives of cosets Γ/Γ_Λ . Then the disjoint union $\Gamma = \bigcup_{h \in \tilde{\Gamma}} h\Gamma_\Lambda$ induces a decomposition

$$\ell^2(\Gamma) = \bigoplus_{h \in \tilde{\Gamma}} \ell^2(h\Gamma_\Lambda).$$

For any pair $h_1, h_2 \in \tilde{\Gamma}$ we define a partial isometry $V_{h_1 h_2}$ from $\ell^2(h_1\Gamma_\Lambda)$ onto $\ell^2(h_2\Gamma_\Lambda)$ by

$$V_{h_1 h_2}(\xi_{h_1 s}) = \xi_{h_2 s}, \quad \forall s \in \Gamma_\Lambda.$$

It is clear that $\{V_{h_1, h_2} : h_1, h_2 \in \tilde{\Gamma}\}$ form a system of matrix units which generates a type I factor, which is $*$ -isomorphic to $\mathcal{L}(\mathcal{H})$. Furthermore, all $V_{h_1 h_2}$ are in the double commutant $W^*(\Gamma, \mathcal{P}_\Lambda)''$. Let P_0 be the projection onto $\ell^2(\Gamma_\Lambda)$. Then it is clear from [8, Vol. II, 6.7] that $P_0 W^*(\Gamma, \mathcal{P}_\Lambda)'' P_0$ is the von Neumann algebra generated by the unitaries $\{L_h : h \in \Gamma_\Lambda\}$. Thus, $P_0 W^*(\Gamma, \mathcal{P}_\Lambda)'' P_0 \cong W^*(\Gamma_\Lambda)$. Consequently, the Double Commutant Theorem [8, Vol. I, 5.3.1] implies

$$W^*(\Gamma, \mathcal{P}_\Lambda) \cong W^*(\Gamma, \mathcal{P}_\Lambda)'' \cong W^*(\Gamma_\Lambda) \otimes \mathcal{L}(\mathcal{H}).$$

\square

1.5. Corollary. (i) If Γ_Λ contains at least two generating cyclic groups and $\Gamma_\Lambda \neq \mathbb{Z}_2 * \mathbb{Z}_2$, then $W^*(\Gamma_\Lambda)$ is a factor of type II_1 and, hence, $W^*(\Gamma, \mathcal{P}_\Lambda)$ is a factor of type II_∞ . In particular, if Γ is a free group \mathcal{F}_n on n generators and $\Lambda = \{g_{n-m}, g_{n-m+1}, \dots, g_n\}$, where $3 \leq n \leq +\infty$ and $n - m \geq 1$, then

$$W^*(\Gamma, \mathcal{P}_\Lambda) \cong \begin{cases} \mathcal{L}^\infty(S^1) \otimes \mathcal{L}(\mathcal{H}) & \text{if } n - m = 1, \\ W^*(\mathcal{F}_{n-m}) \otimes \mathcal{L}(\mathcal{H}) & \text{if } n - m > 1, \end{cases}$$

where $+\infty - 1$ is understood as $+\infty$.

(ii) If $\Gamma = \mathbb{Z} * \mathbb{Z}_m$, where $2 \leq m < +\infty$, and $\Lambda = \{g_i\}$, then

$$W^*(\Gamma, \mathcal{P}_\Lambda) \cong \begin{cases} \mathcal{L}^\infty(S^1) \otimes \mathcal{L}(\mathcal{H}) & \text{if } G_i \cong \mathbb{Z}_m, \\ \underbrace{\mathcal{L}(\mathcal{H}) \oplus \mathcal{L}(\mathcal{H}) \oplus \dots \oplus \mathcal{L}(\mathcal{H})}_{m \text{ times}} & \text{if } G_i \cong \mathbb{Z}. \end{cases}$$

(iii) If $\Gamma = \mathbb{Z}_n * \mathbb{Z}_m$, where $2 \leq m, n < +\infty$, and $\Lambda = \{g_i\}$, then

$$W^*(\Gamma, \mathcal{P}_\Lambda) \cong \underbrace{\mathcal{L}(\mathcal{H}) \oplus \mathcal{L}(\mathcal{H}) \oplus \dots \oplus \mathcal{L}(\mathcal{H})}_{m \text{ times}}, \text{ if } G_i \cong \mathbb{Z}_n.$$

As the most important application of Theorem I we determine non-nuclearity of certain C^* -algebras of the form $C_r^*(\Gamma, \mathcal{P}_\Lambda)$. At first we recall some relevant known results.

- 1.6.** (a) Assume that G is a discrete group. Then C_r^*G is amenable (nuclear) if and only if the group G is amenable ([1], [11]).
- (b) Assume that G is a countable discrete i.c.c. group. Then $W^*(G)$ is $*$ -isomorphic to the unique hyperfinite type II_1 factor \mathcal{R} if and only if G is amenable [4]. Also, $W^*(G) \otimes \mathcal{L}(\mathcal{H})$ is $*$ -isomorphic to the unique injective type II_∞ factor $\mathcal{R} \otimes \mathcal{L}(\mathcal{H})$ if and only if G is amenable [4].
- (c) A C^* -algebra \mathcal{A} is amenable if and only if \mathcal{A} is nuclear, and again if and only if the enveloping von Neumann algebra \mathcal{A}^{**} is injective [3], [4], [10].

1.7. Theorem II. If Γ_Λ contains at least two different generators of Γ but $\Gamma_\Lambda \neq \mathbb{Z}_2 * \mathbb{Z}_2$, then the C^* -algebra $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ is non-nuclear.

Proof. It is well known that our assumptions imply that the group Γ_Λ is non-amenable (e.g. cf. [12, Theorem 1.1]). Consequently, Theorem I and (b) above imply that $W^*(\Gamma, \mathcal{P}_\Lambda)$, the weak closure of $C_r^*(\Gamma, \mathcal{P}_\Lambda)$, is not injective. Thus, the enveloping von Neumann algebra $C_r^*(\Gamma, \mathcal{P}_\Lambda)^{**}$ is not injective either, and (c) above implies that the C^* -algebra $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ is non-nuclear, as desired. \square

1.8. Remark. The problem of nuclearity of those algebras $C_r^*(\Gamma, \mathcal{P}_\Lambda)$ for which the group Γ_Λ is amenable (i.e. Γ_Λ is either cyclic or $\mathbb{Z}_2 * \mathbb{Z}_2$) has already been answered in the affirmative in [16].

ACKNOWLEDGEMENTS

The second named author wishes to thank Professor George Elliott for inviting him to the Fields Institute of Canada in the Summer of 1994. Both authors are grateful to Professor Zhong-Jin Ruan for providing some very helpful references.

REFERENCES

1. J. Bunce, *Finite operators and amenable C^* -algebras*, Proc. Amer. Math. Soc. **56** (1976), 145–151. MR **53**:6333
2. M. D. Choi, *A simple C^* -algebra generated by two finite-order unitaries*, Canad. J. Math. **31** (1979), 867–880. MR **80j**:46092
3. M. D. Choi and E. G. Effros, *Nuclear C^* -algebras and injectivity: the general case*, Indiana Univ. Math. J. **26** (1977), 443–446. MR **55**:3799
4. A. Connes, *Classification of injective factors*, Ann. of Math. **104** (1976), 73–115. MR **56**:12908
5. J. Cuntz, *K -theory for certain C^* -algebras*, Ann. of Math. **113** (1981), 181–197. MR **84c**:46058
6. G. A. Elliott, *A classification of certain simple C^* -algebras*, J. reine angew. Math. **443** (1993), 179–219. MR **94i**:46074
7. G. A. Elliott and M. Rørdam, *Classification of certain infinite simple C^* -algebras, II*, Comment. Math. Helv. **70** (1995), 615–638. MR **96e**:46080b
8. R. V. Kadison and J. R. Ringrose, *Fundamentals of the theory of operator algebras, Vol. I, II*, Pure and Applied Mathematics, A Series of Monographs and Textbooks, Academic Press, Inc. (1986). MR **98f**:46001a; MR **96f**:46001b
9. E. Kirchberg, *The classification of purely infinite C^* -algebras using Kasparov's theory (preprint)*.
10. C. Lance, *Tensor products and nuclear C^* -algebras*, Operator Algebras and Applications (ed. R. V. Kadison), Proc. Symp. Pure Math., Part I **38** (1981), 379–399.
11. C. Lance, *On nuclear C^* -algebras*, J. Funct. Anal. **12** (1973), 157–176. MR **49**:9640
12. W. L. Paschke and N. Salinas, *C^* -algebras associated with free products of groups*, Pac. J. Math. **82** (1979), 211–221. MR **82c**:22010
13. N. C. Phillips, *Classification of purely infinite simple C^* -algebras*, preprint, 1994.
14. M. Rørdam, *Classification of inductive limits of Cuntz algebras*, J. reine angew. Math. **440** (1993), 175–200. MR **94k**:46120
15. J. Spielberg, *Free-product groups, Cuntz-Krieger algebras, and covariant maps*, Internat. J. Math. **2** (1991), 457–476. MR **92j**:46120
16. W. Szymański and S. Zhang, *Infinite simple C^* -algebras and reduced cross products of abelian C^* -algebras and free groups*, Manuscripta Math. **92** (1997), 487–514. MR **98a**:46073
17. S. Zhang, *Purely infinite simple C^* -algebras arising from reduced group C^* -algebras*, Contemp. Math. **228** (1998), 365–389. CMP 99:07
18. S. Zhang, *C^* -algebras generated by a projection and the reduced group C^* -algebras $C_r^*\mathbb{Z} * \mathbb{Z}_n$ and $C_r^*\mathbb{Z}_m * \mathbb{Z}_n$* , preprint, 1994.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF NEWCASTLE, NEWCASTLE, NEW SOUTH WALES 2308, AUSTRALIA

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF CINCINNATI, CINCINNATI, OHIO 45221-0025

E-mail address: zhangs@math.uc.edu