

SMALL INDUCTIVE DIMENSION OF COMPLETIONS OF METRIC SPACES. II

S. MRÓWKA

(Communicated by Alan Dow)

ABSTRACT. Extending the results of a previous paper under the same title we show that, under $\mathbf{S}(\aleph_0)$, $\text{ind}_c \nu\mu_0^2 = 2$.

1. INTRODUCTION

In this paper we continue the study of the small inductive dimension, ind , of completions of esoteric spaces (= metric spaces with $\text{ind} \neq \dim$) initiated in a previous paper under the same title [M2]. In [M2] we have described the metric space $\nu\mu_0$ such that $\text{ind} \nu\mu_0 = 0$, but, under the following condition every completion of $\nu\mu_0$ contains an interval and therefore $\text{ind}_c \nu\mu_0 = 1$, where, for a metric space M , $\text{ind}_c M$ stands for $\min\{\text{ind} \tilde{M} : \tilde{M} \text{ is a completion of } M\}$:

$\mathbf{S}(\aleph_0)$: *If A is a set of cardinality 2^{\aleph_0} , then the product A^{\aleph_0} cannot be written as $A^{\aleph_0} = F_1 \cup F_2 \cup \dots$, where each F_n is an \mathbf{F}_σ -set in the product topology of A^{\aleph_0} (A – discrete) and it is countable on all lines parallel to the n -th axis.¹*

$\mathbf{S}(\aleph_0)$ disagrees very strongly with the continuum hypothesis, but fortunately its consistency with ZFC has been recently shown by Dougherty [Dou], who, in this way, terminated a long and unenviable period during which I was at the mercy of the logicians. In this paper we give a very natural continuation of [M2]; our main result is

1.1. Main Theorem. *Under $\mathbf{S}(\aleph_0)$, every completion of $\nu\mu_0^2$ contains the square $[0, 1]^2$ and therefore $\text{ind}_c \nu\mu_0^2 = \dim \nu\mu_0^2 = 2$.*

The result concerning $\text{ind}_c \nu\mu_0^2$ appears to be very natural. The problem of the equality $\text{ind}_c = \text{ind}$ (which, for spaces with $\text{ind} = 0$, can be stated without any reference to dimension theory: *does every metric space having a base consisting of clopen sets have a completion with such a base?*) is one of the simplest and most

Received by the editors March 9, 1998 and, in revised form, June 4, 1998.

1991 *Mathematics Subject Classification.* Primary 54F45; Secondary 54A35, 54E35, 54H05.

Key words and phrases. Inductive and covering dimension, metric spaces, completion, Bernstein sets, scattered sets.

¹If \tilde{a} is an element of the product $\prod_{\eta \in H} A_\eta$ and $\xi \in H$, then the line passing through \tilde{a} and parallel to the ξ -th axis, $L_\xi^{\tilde{a}}$, is the set of all elements of $\prod_{\eta} A_\eta$ which differ from \tilde{a} only in the ξ -th coordinate. We say that a subset F of $\prod_{\eta} A_\eta$ has a property \mathbf{P} on all lines parallel to the ξ -th axis provided that, for every $\tilde{a} \in \prod_{\eta} A_\eta$, $F \cap L_\xi^{\tilde{a}}$ has \mathbf{P} . More generally, if $\Xi \subset H$, then we can define hyperplanes parallel to the Ξ -th hyperaxis and speak about a set having a given property on all such hyperplanes.

basic questions concerning concepts that were in existence for over 75 years. Before [M2], nothing was known about the answer to this question (for non-separable spaces). Now the first step has been made; but only partial – not validity, only consistency. This is not unusual; it happens quite often that consistency results are obtained first because they are easier – or much easier – than validity or because validity cannot be attained at all and the final solution is the independence. But once the first step has been made, the next is following very quickly. In fact, as far as consistency is concerned, all the main questions concerning ind_c (or even $\text{ind}_c^*(M) = \min\{\text{ind}(\tilde{M} \setminus M) : \tilde{M} \text{ is a completion of } M\}$; see part b) of 4.3) are now answered. However, the matters become more unusual if we look at dim . This definitely is *not* the first step in this direction; indeed, the first step, existence of metric spaces with $\text{d-spread} = \text{dim} - \text{ind} = 1$, had been made about 35 years ago [R]. So far all the efforts to raise the value of d-spread from 1 to 2 have failed. Now, in a way, this has been done but not as well as for $\text{d-spread} = 1$. The reason why there is such a chasm between the first two values of d-spread is not yet understood.

A further consequence of Theorem 1.1 is the existence, in ZFC, of metric spaces for which dim cannot be determined in ZFC; indeed, under the continuum hypothesis, $\text{dim } \nu\mu_0^2 = 1$ (the proof from [M3] applies). Concerning ind_c , this was already achieved in [M2]: $\text{ind}_c \nu\mu_0$ cannot be determined in ZFC (but note that, in ZFC, $\text{dim } \nu\mu_0 = 1$). Since the existence of spaces with undeterminable dim and/or ind_c is an entirely new phenomenon, worthy of further development, an additional discussion of these matters is given in the problem section of this paper (sect. 5); here we will summarize what is known at the present moment:

1.2. *The values of ind , ind_c , and dim of $\nu\mu_0$ and $\nu\mu_0^2$ are as follows:*

- (i) *in ZFC:* $\text{ind } \nu\mu_0 = \text{ind } \nu\mu_0^2 = 0;$ $\text{dim } \nu\mu_0 = 1;$
- (ii) *under CH:* $\text{ind}_c \nu\mu_0 = \text{ind}_c \nu\mu_0^2 = 0;$ $\text{dim } \nu\mu_0^2 = 1;$
- (iii) *under $S(\aleph_0)$:* $\text{ind}_c \nu\mu_0 = 1;$ $\text{ind}_c \nu\mu_0^2 = \text{dim } \nu\mu_0^2 = 2.$

To keep the paper short we just show what is new rather than try to get the strongest version of the results; e.g., we work only with $\nu\mu_0^2$ although it is reasonably certain that better results could be obtained through the use of higher powers of $\nu\mu_0$.

Note. The author is pleased to report substantial progress in the area of esoteric spaces due to Kulesza [Ku1], [Ku2] and also to Ostaszewski [Ost]. Kulesza has produced numerous modifications of Δ and μ_0 and in this way he was able to vary properties of esoteric spaces to a large degree. In particular, he has a *complete* \aleph -compact esoteric space. He also studies *irreducible* esoteric spaces – in the sense that they do not contain esoteric subspaces of weight smaller than that of the space itself. This seems to be a promising direction of investigations; indeed, the concept of irreducibility can be modified in many ways and this leads to a number of very natural but probably very difficult problems. Again, further comments on this matter are in sect. 5.

Notation. Let t be a number from $[0, 1]$. If t can be written in the form $t = (2k + 1)2^{-n}$, where k is an integer and n is a non-negative integer, then (t is called a dyadic rational and) we let $\text{ord } t = n$; if t is a dyadic irrational, then, obviously, we let $\text{ord } t = \infty$. The letters \mathcal{R} , \mathcal{N} , \mathcal{J} , \mathcal{P} , \mathcal{C} will stand, respectively, for the space

of the reals, the positive integers, the closed interval $[0, 1]$, the dyadic irrationals in $[0, 1]$ and the Cantor set.

Concluding this introduction I wish to express my sincere gratitude to the referee for suggesting several changes, correcting several errors and for completing the report in a remarkably short time.

2. TOOLS

The only item of substance in this section is 2.5. In addition, a variation of the concept of Bernstein sets (2.4 and above) is of interest.

A. σ -non-archimedean bases. A σ -non-archimedean base \mathfrak{B} in a space X is a sequence $\mathfrak{B}^{(1)}, \mathfrak{B}^{(2)}, \dots$ such that each $\mathfrak{B}^{(k)}$ covers X , $U \in \mathfrak{B}^{(k)}, V \in \mathfrak{B}^{(j)}$ and $U \cap V \neq \emptyset$ imply $V \subset U$ for every $k, j = 1, 2, \dots, k \leq j$ and $\bigcup_k \mathfrak{B}^{(ki)}$ is a base for X . We shall frequently identify \mathfrak{B} with $\bigcup_k \mathfrak{B}^{(k)}$ (i.e., we will write $\mathfrak{B} = \bigcup_k \mathfrak{B}^{(k)}$). Note that the members $\mathfrak{B}^{(k)}$ are mutually disjoint, hence $\mathfrak{B}^{(k)}$ is a discrete collection of clopen sets. \mathfrak{B} is called *complete* provided that for every descending sequence U_n with $\emptyset \neq U_n \in \mathfrak{B}^{(i_n)}$, where $i_n \rightarrow \infty$, we have $\bigcap_n U_n \neq \emptyset$.

The following can be regarded as known.

2.1. *A space X has a σ -non-archimedean base iff it is strongly 0-dimensional and metrizable. A space X has a complete σ -non-archimedean base iff it is strongly 0-dimensional and metrizable in a complete way.*

Historical note. The second sentence of 2.1 follows from a strongly 0-dimensional version of a result of Kuratowski:

2.2. *If X is a metric space with $\dim X = 0$, then for every \mathbf{G}_δ -subset G of X we have $G \subset_{\text{cl}} X \times N^{\aleph_0}$.*

The original Kuratowski result asserts: *for every \mathbf{G}_δ -subset G of a metric space X we have $G \subset_{\text{cl}} X \times \mathcal{R}^{\aleph_0}$.* Both versions are particular cases of 2.1 in [M1] (the Embedding Theorem) or – *vice versa* – the Embedding Theorem can be viewed as a natural continuation of the Kuratowski result. In turn, the Kuratowski result is a natural continuation of a very early result of Kuratowski and Sierpiński: *if a subset L of a metric space X is the difference of two closed sets, then $L \subset_{\text{cl}} X \times \mathcal{R}$.* As far as I know, the above is the first result on closed embedding into products. If the reader knows earlier results, I would appreciate a note to this effect.

B. Bernstein sets. A Bernstein subset of a space X is a set $S \subset X$ which intersects every non-empty perfect subset of X . It is known that Bernstein subsets of \mathcal{J}^2 are connected.

It is possible that the following is well-known.

2.3. *If S is a subset of \mathcal{J}^2 so that S and $\mathcal{J}^2 \setminus S$ are both Bernstein, then for every \mathbf{G}_δ set G with $S \subset G \subset \mathcal{J}^2$, $G \setminus S$ is also Bernstein. Consequently, for every completion \tilde{S} of S , $\tilde{S} \setminus S$ is connected.*

Proof. Let $C \subset \mathcal{J}^2$ be a perfect set. Then $S \cap C$ is dense in C ; thus $G \cap C$ is dense in C . Consequently, $G \cap C$ contains a perfect set C_1 . Now, $\mathcal{J}^2 \setminus S$ intersects C_1 ; thus G intersects $\mathcal{J}^2 \setminus S$, and hence $G \setminus S$ intersects C_1 . \square

A subset K of a product $X \times Y$ will be called *oblique* provided that for every $(x_1, y_1), (x_2, y_2) \in K, (x_1, y_1) \neq (x_2, y_2)$ imply $x_1 \neq x_2$ and $y_1 \neq y_2$; i.e., both projection maps of K are one-to-one. A *2-Bernstein* subset of a space X is a set $S \subset X$ such that S^2 intersects every non-empty oblique perfect subset of X^2 . The standard construction of disjoint Bernstein sets (in perfect metric spaces) can be easily adapted to yield that the Cantor set \mathcal{C} contains two disjoint 2-Bernstein sets.

For one of the results in sect. 4 we need a further refinement. Let \mathcal{D} be a collection of subsets of Y . A subset S of X is called *\mathcal{D} -Bernstein* provided that for every $D \in \mathcal{D}$, $S \times D$ intersects every non-empty oblique perfect subset of $X \times Y$. We have

2.4. *If X and Y are complete dense-in-itself separable metric spaces and \mathcal{D} is a collection of Bernstein subsets of Y with $\text{card } \mathcal{D} \leq 2^{\aleph_0}$, then X contains two disjoint \mathcal{D} -2-Bernstein sets $B^{(1)}$ and $B^{(2)}$.*

A \mathcal{D} -2-Bernstein set is, obviously, a set which is 2-Bernstein and \mathcal{D} -Bernstein. The above is not the best result: the number of \mathcal{D} -2-Bernstein sets can be increased and the result can be extended to \mathcal{D} - n -Bernstein sets. But we need even less than 2.4.

C. Scattered sets. Let \mathbf{Q} be downward monotone property of closed sets (i.e., F has \mathbf{Q} and $F' \subset F$ imply F' has \mathbf{Q}). We say that F is *\mathbf{Q} -scattered* provided that for every non-empty closed $F' \subset F$ there is an open U so that $U \cap F'$ is closed, non-empty and has \mathbf{Q} .

2.5. *Let X be a space with a σ -non-archimedean base $\mathfrak{B} = \mathfrak{B}^{(1)} \cup \mathfrak{B}^{(2)} \cup \dots$. Let $\mathbf{Q}_1, \dots, \mathbf{Q}_l$ be downward monotone properties of closed sets. If a closed set F is $\mathbf{Q}_1 \vee \dots \vee \mathbf{Q}_l$ -scattered, then there are closed sets $F_{k,i}$ and classes $\mathfrak{F}_{k,i} \subset \mathfrak{B}^{(k)}$, $k = 1, 2, \dots, i = 1, \dots, l$, such that (i) $F = \bigcup_{k,i} F_{k,i}$, (ii) $F_{k,i} \subset \bigcup \mathfrak{F}_{k,i}$ and (iii) for every $U \in \mathfrak{F}_{k,i}$, $U \cap F_{k,i}$ has \mathbf{Q}_i .*

Proof. Let λ be an ordinal with $\bar{\lambda} > \text{card } F$. We shall show that there exists a ξ_0 with $0 < \xi_0 < \lambda$ and such that for every $\eta < \xi_0$ we can define $F^{(\eta)}$ and $\mathfrak{F}_{k,i}^{(\eta)} \subset \mathfrak{B}^{(k)}$ so that (a) $F^{(\eta)}$ form a strictly decreasing sequence of closed sets, (b) $F^{(\eta)} \cap \bigcup \left\{ \bigcup \mathfrak{F}_{k,i}^{(\eta')} : \eta' < \eta, k = 1, 2, \dots, i = 1, \dots, l \right\} = \emptyset$, (c) for every $U \in \mathfrak{F}_{k,i}^{(\eta)}$, $U \cap F^{(\eta)}$ has \mathbf{Q}_i , and (d) $\bigcap_{\eta < \xi_0} F^{(\eta)} = \emptyset$.

Let $F^{(0)} = F$ and $\mathfrak{F}_{k,i}^{(0)} = \emptyset$. Assume that for some ξ , $0 < \xi < \lambda$, $F^{(\eta)}$ and $\mathfrak{F}_{k,i}^{(\eta)}$ are already defined for every $\eta < \xi$ so that (a), (b) and (c) are satisfied (this was so for $\xi = 1$). If $\bigcap_{\eta < \xi} F^{(\eta)} = \emptyset$, then we terminate the induction. Otherwise, we continue letting $F^{(\xi)} = \bigcap_{\eta < \xi} F^{(\eta)} \setminus \bigcup \left\{ \bigcup \mathfrak{F}_{k,i}^{(\eta)} : \eta < \xi, k = 1, 2, \dots, i = 1, \dots, l \right\}$ and $\mathfrak{F}_{k,i}^{(\xi)} = \left\{ U \in \mathfrak{B}^{(k)} : U \cap F^{(\xi)} \neq \emptyset, U \cap F^{(\xi)} \text{ has } \mathbf{Q}_i \right\}$ (if there are several i 's so that $U \cap F^{(\xi)}$ has \mathbf{Q}_i , then use just one of them). Since $\bar{\lambda} > \text{card } F$, the induction must terminate before reaching λ ; i.e., the $\xi_0 < \lambda$ exists. Observe that for $\xi < \xi' < \xi_0$, $\mathfrak{F}_{k,i}^{(\xi)}$ and $\mathfrak{F}_{k,i}^{(\xi')}$ are disjoint, indeed, $U \in \mathfrak{F}_{k,i}^{(\xi')}$ implies $U \cap F^{(\xi')} \neq \emptyset$ and $F^{(\xi')}$ is disjoint from $\bigcup \mathfrak{F}_{k,i}^{(\xi)}$. It follows that letting $\mathfrak{F}_{k,i} = \bigcup \left\{ \mathfrak{F}_{k,i}^{(\eta)} : \eta < \xi_0 \right\}$, we have, for a $U \in \mathfrak{F}_{k,i}$, $U \cap F = U \cap F^\eta$ for the unique η with $U \in \mathfrak{F}_{k,i}^{(\eta)}$. Thus, letting $F_{k,i} = F \cap \bigcup \mathfrak{F}_{k,i}$ we complete the proof. \square

3. PROOF OF THE MAIN THEOREM

The proof of 1.1 will come after a sequence of intermediate results. We start with the description of objects used in the proof.

Recall that the construction of $\nu\mu$ and $\nu\mu_0$ involved the Cantor set \mathcal{C} and its subset A . As in [M2] we need that $\text{card } A = 2^{\aleph_0}$ but this time we also need that $B = \mathcal{C} \setminus A$ is 2-Bernstein. In addition, for 4.1 (the case concerning Bernstein set) we will need B to be \mathcal{D} -2-Bernstein, where \mathcal{D} will be specified in the proof of 4.1. Infinite sequences of elements of \mathcal{C} (i.e., members of \mathcal{C}^{\aleph_0}) will be denoted as $\tilde{x}, \tilde{y}, \tilde{z}, \dots$, possibly with subscripts, and furthermore $\tilde{x}(n)$ (and not \tilde{x}_n) will denote the n -th term of \tilde{x} . The letter \mathbf{a} will be reserved for members of A^{\aleph_0} ; it will be used mainly in the case when such a member is built up by extracting terms from an $\tilde{x} \in \mathcal{C}^{\aleph_0}$ (see the definition of the maps Φ^{m_1, m_2} and Φ_i^m below). In the set $(A^{\aleph_0})^2$ of all pairs $(\mathbf{a}_1, \mathbf{a}_2)$, $\mathbf{a}_i \in A^{\aleph_0}$; $\mathbf{a}_i(m)$ will be called the (m, i) -coordinate of $(\mathbf{a}_1, \mathbf{a}_2)$.

We need some maps. The map $\Phi^{m_1, m_2} : (A^{\aleph_0})^2 \rightarrow A^{\aleph_0} \times \mathcal{C} \times \mathcal{C}$ carries $(\tilde{x}_1, \tilde{x}_2)$ onto the triple (\mathbf{a}, c_1, c_2) , where $\mathbf{a} \in A^{\aleph_0}$ is a sequence built out of terms of \tilde{x}_1 and \tilde{x}_2 , omitting the terms $\tilde{x}_1(m_1)$ and $\tilde{x}_2(m_2)$ and $c_i = \tilde{x}_i(m_i)$. The map $\Phi_i^m : (A^{\aleph_0})^2 \rightarrow A^{\aleph_0} \times \mathcal{C}$ carries an $(\tilde{x}_1, \tilde{x}_2) \in (A^{\aleph_0})^2$ onto the pair (\mathbf{a}, c) , where \mathbf{a} is the sequence build out of terms of \tilde{x}_1 and \tilde{x}_2 , omitting the term $\tilde{x}_i(m)$ and c is $\tilde{x}_i(m)$. Φ^{m_1, m_2} and Φ_i^m are continuous and one-to-one.

Recall that the set of points of $\nu\mu$ is the set of all pairs (\tilde{x}, t) where $t \in \mathbb{J}$ and $\tilde{x}(n) \in A$ for every $n \neq \text{ord } t$. The topology is defined by the neighborhoods $U_n(\tilde{x}, t)$:

$$U_n(\tilde{x}, t) = \{(\tilde{y}, s) : s \in I_n(t) \text{ and } \tilde{y}(i) = \tilde{x}(i) \text{ for } i \leq n\},$$

if $n < \text{ord } t$, and

$$U_n(\tilde{x}, t) = \{(\tilde{y}, s) : s \in I_n(t) \text{ and } \tilde{y}(i) = \tilde{x}(i) \text{ for } i \leq n + 1, i \neq \text{ord } t; \tilde{y}(i)|n = \tilde{x}(i)|n \text{ for } i = \text{ord } t\},$$

if $n \geq \text{ord } t$.

In the above, $I_n(t)$ is an open interval (a, b) containing t and so that a and b are the dyadic rationals of the smallest possible orders with $b - a = 2^{-n}$. $\nu\mu_0$ consists of all those $(\tilde{x}, t) \in \nu\mu$ for which $\tilde{x}(n) \in B$ for $n = \text{ord } t$.

For $\tilde{x}, \tilde{y} \in \mathcal{C}^{\aleph_0}$ the $\tilde{x}\tilde{y}$ -plank (or a horizontal plank) of $\nu\mu^2$ is the set $H_{\tilde{x}, \tilde{y}}$ of all the $((\tilde{x}, t), (\tilde{y}, s)) \in \nu\mu^2$ with the fixed \tilde{x}, \tilde{y} . For a subspace κ of $\nu\mu$, the $\tilde{x}\tilde{y}$ -plank of κ is the set $H_{\tilde{x}, \tilde{y}} \cap \kappa$. Let $\hat{h} = h \times h$ where $h : \nu\mu \rightarrow I$ is defined by $h(\tilde{x}, t) = t$ (i.e., h is the function from 2.3 in [M2]). By 2.3 in [M2], \hat{h} is a homeomorphism on each of the horizontal planks.

Let t, t_1, t_2 be dyadic rationals of orders m, m_1, m_2 , respectively; we let $\mathcal{P}_1^t = \{t\} \times \mathcal{P}$, $\mathcal{P}_2^t = \mathcal{P} \times \{t\}$ (\mathcal{P} is the set of all dyadic irrationals in \mathbb{J}), $\mathbb{P}^{t_1, t_2} = \hat{h}^{-1}(t_1, t_2)$, $\mathbb{P}_i^t = \hat{h}^{-1}[\mathcal{P}_i^t]$.

The map

$$\hat{\Phi}^{t_1, t_2}((\tilde{x}_1, t_1), (\tilde{x}_2, t_2)) = \Phi^{m_1, m_2}(\tilde{x}_1, \tilde{x}_2)$$

is a homeomorphism of \mathbb{P}^{t_1, t_2} onto $A^{\aleph_0} \times \mathcal{C} \times \mathcal{C}$. Similarly, the map

$$\hat{\Phi}_i^t((\tilde{x}_1, t), (\tilde{x}_2, u)) = (\Phi_i^m(\tilde{x}_1, \tilde{x}_2), u)$$

is a homeomorphism of \mathbb{P}_i^t onto $A^{\aleph_0} \times \mathcal{C} \times \mathcal{P}$.

We are ready to start the argument. Since every completion of $\nu\mu_0^2$ contains a \mathbf{G}_δ -subset F of $\nu\mu^2$ with $\nu\mu_0^2 \subset F$, it suffices to show that F contains a copy of \mathcal{J}^2 . Let such an F be given and let $\nu\mu^2 \setminus F = \bigcup_n F_n$, where F_n are closed in $\nu\mu^2$.

Select complete σ -non-archimedean bases $\mathfrak{B}_l = (\mathfrak{B}_l^{(1)}, \mathfrak{B}_l^{(2)}, \dots)$, $l = 1, 2, 3$, in the spaces A^{\aleph_0} , \mathcal{C} and \mathcal{P} , respectively. \mathfrak{B}_1 and \mathfrak{B}_2 are countable; we let $\mathfrak{B}_1 = \{J_1, J_2, \dots\}$, $\mathfrak{B}_2 = \{I_1, I_2, \dots\}$. Letting $\mathfrak{B}^{(k)} = \{U \times J \times I : U \in \mathfrak{B}_1^{(k)}, J \in \mathfrak{B}_2^{(k)}, I \in \mathfrak{B}_3^{(k)}\}$, we obtain a (complete) σ -non-archimedean base in $A^{\aleph_0} \times \mathcal{C} \times \mathcal{P}$.

Let $K_n^{t_1, t_2} = \Phi^{t_1, t_2}[F_n \cap \mathbb{P}^{t_1, t_2}]$. Let Q_i be the property of closed subsets T of $A^{\aleph_0} \times \mathcal{C} \times \mathcal{C}$: “all $(\mathbf{a}, c^1, c^2) \in T$ have the same c^i ”.

3.1. $K_n^{t_1, t_2}$ is $Q_1 \vee Q_2$ -scattered.

$K_n^{t_1, t_2}$ is a closed subset of $A^{\aleph_0} \times \mathcal{C} \times \mathcal{C}$ and since F_n are disjoint from $\nu\mu_0^2$, $K_n^{t_1, t_2}$ is disjoint from $A^{\aleph_0} \times B \times B$. The rest of the proof is the standard procedure of producing the homeomorph of the Cantor set in complete dense-in-itself metric spaces. If 3.1 fails, then there is a closed subset M of $K_n^{t_1, t_2}$ such that for every $\mathbf{U} \in \mathfrak{B}$ with $\mathbf{U} \cap M \neq \emptyset$ there are points $p, q \in \mathbf{U} \cap M$ for which both c^1 's and c^2 's are distinct. (Strictly speaking, we get the existence of points p_1, p_2, q_1, q_2 – not necessarily distinct – so that p_1, p_2 have distinct c^1 's and q_1, q_2 have distinct c^2 's. Patiently examining possible cases we get the stated conclusion.) Starting with an arbitrary $\mathbf{U} \in \mathfrak{B}$ with $\mathbf{U} \cap M \neq \emptyset$, we denote the above-mentioned points by p_0 and p_1 . Now we select $\mathbf{U}_0, \mathbf{U}_1 \in \mathfrak{B}$ such that $p_i \in \mathbf{U}_i \subset \mathbf{U}$ and so that, writing $\mathbf{U}_i = U_i \times J_i \times I_i$, we have $J_0 \cap J_1 = \emptyset = I_0 \cap I_1$. Now we select points $p_{i,0}, p_{i,1} \in \mathbf{U}_i \cap M$ for which both c^1 's and c^2 's are distinct. Repeating the above – in a standard way – infinitely many times we obtain a system of members of \mathfrak{B} and of points of M indexed by all finite sequences of 0's and 1's. Passing to the limit, we obtain the set of points $(\mathbf{a}_i, c_i^1, c_i^2)$, – the index i runs through all infinite sequences of 0's and 1's – such that $c_i^1 \neq c_{i'}^1$ and $c_i^2 \neq c_{i'}^2$ for every two distinct i and i' and the set $C = \{(c_i^1, c_i^2)\}_i$ is closed in $\mathcal{C} \times \mathcal{C}$. Since M is closed, all the $(\mathbf{a}_i, c_i^1, c_i^2)$ belong to M . But C is an oblique perfect set, therefore there is an i_0 with $(c_{i_0}^1, c_{i_0}^2) \in B^2$. But then $(\mathbf{a}_{i_0}, c_{i_0}^1, c_{i_0}^2) \in A^{\aleph_0} \times B \times B$. Contradiction.

Now apply 2.5 to get the classes $\mathfrak{F}_{n,i,k}^{t_1, t_2}$ and the sets $K_{n,i,k}^{t_1, t_2}$. Let $\hat{K}_{n,i,k}^{t_1, t_2} = (\Phi^{m_1, m_2})^{-1}[K_{n,i,k}^{t_1, t_2}]$, where $m_i = \text{ord } t_i$. $\hat{K}_{n,i,k}^{t_1, t_2}$ is closed subset of $(A^{\aleph_0})^2$ and since all $(\mathbf{a}, c^1, c^2) \in K_{n,i,k}^{t_1, t_2}$ have the same c^i , $\hat{K}_{n,i,k}^{t_1, t_2}$ has at most one element on each line parallel to the (m_i, i) -axis.

Let $K_n^{*t} = \Phi_i^t[F_n \cap \mathbb{P}_i^t]$; the handling of these sets is more intricate than that of $K_n^{t_1, t_2}$. Let Q be the property of closed subsets T of $A^{\aleph_0} \times \mathcal{C} \times \mathcal{P}$: “all $(\mathbf{a}, c, u) \in T$ have the same c ”.

3.2. $K_{n,i}^{*t}$ is Q -scattered.

The proof is similar to that of 3.1 but now, assuming that 3.2 fails, we can produce a set of points $(\mathbf{a}_i, c_i, u_i) \in K_{n,i}^{*t}$ in which c_i form a perfect set and u_i are dyadic irrational but not necessarily distinct. Taking i_0 with $c_{i_0} \in B$, we have $(\mathbf{a}_{i_0}, c_{i_0}, u_{i_0}) \in \nu\mu_0$ and we have the contradiction.

Now apply 2.5 to get the classes $\mathfrak{F}_{n,i,k}^t$ and the sets $K_{n,i,k}^{*t}$. Further, let

$$\mathfrak{F}_{n,i,k,l_1,l_2}^t = \{U \times J \times I \in \mathfrak{F}_{n,i,k}^t : J = J_{l_1}, I = I_{l_2}\},$$

$$K_{n,i,k,l_1,l_2}^{*t} = K_{n,i,k}^{*t} \cap \bigcup \mathfrak{F}_{n,i,k,l_1,l_2}^t.$$

Let $\overline{K_{n,i,k,l_1,l_2}^{*t}}$ be the closure of K_{n,i,k,l_1,l_2}^{*t} in $A^{\aleph_0} \times \mathcal{C} \times \mathcal{J}$ (under the natural embedding of \mathcal{P} in \mathcal{J}). Since $\mathfrak{F}_{n,i,k,l_1,l_2}^t$ is discrete,

$$\overline{K_{n,i,k,l_1,l_2}^{*t}} = \bigcup \{ \overline{K_{n,i,k}^{*t}} \cap (U \times J_{l_1} \times I_{l_2}) : U \times J_{l_1} \times I_{l_2} \in \mathfrak{F}_{n,k,j}^{t,*} \};$$

therefore all $(\mathbf{a}, c, u) \in \overline{K_{n,i,k,l_1,l_2}^{*t}}$ still have the same c . Finally, let K_{n,i,k,l_1,l_2}^t be the projection of $\overline{K_{n,i,k,l_1,l_2}^{*t}}$ onto $A^{\aleph_0} \times \mathcal{C}$. Since this is a projection along a compact axis, K_{n,i,k,l_1,l_2}^t is closed in $A^{\aleph_0} \times \mathcal{C}$ and all $(\mathbf{a}, c) \in \overline{K_{n,i,k,l_1,l_2}^t}$ still have the same c . Now, letting $\hat{K}_{n,i,k,l_1,l_2}^t = (\Phi_i^m)^{-1}[K_{n,i,k,l_1,l_2}^t]$, where $m = \text{ord } t$, we obtain a closed subset of $(A^{\aleph_0})^2$ which has at most one element on each line parallel to the (m, i) -axis.

We thus have a countable collection of closed subsets

$$(1) \quad \hat{K}_{n,i,k}^{t_1,t_2} \quad \text{and} \quad \hat{K}_{n,i,k,l_1,l_2}^t$$

of $(A^{\aleph_0})^2$ each of which has at most one element on all lines parallel to some axis. Grouping these sets properly we obtain \mathbf{F}_σ -sets to which $S(\aleph_0)$ applies. Hence there is an $(\tilde{x}_0, \tilde{y}_0) \in (A^{\aleph_0})^2$ which does not belong to any of the sets in (1). But then, verifying the definitions of these sets we find that $((\tilde{x}_0, t), (\tilde{y}_0, s)) \in F$ for every $t_1; t_2 \in \mathcal{J}$. In other words, \hat{h} maps (homeomorphically) the $\tilde{x}_0\tilde{y}_0$ -plank of F onto \mathcal{J}^2 ; thus F has a copy of \mathcal{J}^2 .

4. EXTENSION OF 1.1: COMPLETIONS OF SUBSPACES $\nu\mu_0^2$

Given a subset S of \mathcal{J}^2 and a subspace κ of $\nu\mu^2$, the κ -cylinder over S is the set $\kappa(S) = \tilde{h}^{-1}[S] \cap \kappa$ (explicitly $\kappa(S) = \{(\tilde{x}, t), (\tilde{y}, s) \in \kappa : (t, s) \in S\}$).

4.1. *If $S \subset \mathcal{J}^2$ is either \mathbf{G}_δ -set or a Bernstein set in \mathcal{J}^2 , then every completion of $\nu\mu_0(S)$ has a horizontal plank homeomorphic to a completion of S .²*

Proof. The proof is the repetition of the arguments of the preceding section with some changes. 3.1 is applied only to those dyadic rationals t_1, t_2 for which $(t_1, t_2) \in S$. 3.2 requires more attention. \mathbb{P}_i^t are replaced by $\mathbb{P}_i^t(S)$ and $K_{n,i}^{*t}$ are replaced by $K_{n,i}^{*t}(S) = \Phi_i^t[F_n \cap \mathbb{P}_i^t(S)]$. Proof is again by contradiction; i.e., we assume that $K_{n,i}^{*t}(S)$ is not Q -scattered. If S is \mathbf{G}_δ -set, then we work in the space $A^{\aleph_0} \times \mathcal{C} \times (S \cap \mathcal{J}_i^t)$. \mathfrak{B}_3 will now stand for complete σ -non-archimedean base in $S \cap \mathcal{J}_i^t$ (use 2.1 to get such a base) and the rest of the argument is as before.

In the case when S is a Bernstein set, we work under the assumption (recall the remarks at the beginning of sect. 3) that B is a \mathcal{D} -2-Bernstein, with $\mathcal{D} = \{D_i^t : t\text{-dyadic rational, } i = 1, 2\}$, where D_i^t is the intersection $\mathcal{P}_i^t \cap S$ treated as a subset of \mathcal{P} . As before, we work in the space $A^{\aleph_0} \times \mathcal{C} \times \mathcal{P}$, but we distinguish two cases. Let \mathbf{Q}' be the property of closed subsets T of $A^{\aleph_0} \times \mathcal{C} \times \mathcal{P}$: “all $(\mathbf{a}, c, u) \in T$ have the same u ”. If $K_{n,i}^{*t}(S)$ is \mathbf{Q}' -scattered, then we select a $\mathbf{U}_0 = U_0 \times J_0 \times I_0 \in \mathfrak{B}$ so that all $(\mathbf{a}, c, u) \in K_{n,i}^{*t}(S) \cap \mathbf{U}_0$ have the same u , say u_0 . We can get a set of points $(\mathbf{a}_i, c_i, u_0) \in K_{n,i}^{*t}(S)$ so that c_i form a perfect set – contradiction (same as

²In case S is a Bernstein set, the set A used in the definition of $\nu\mu_0$ depends on S . Thus, strictly speaking, this part of the theorem should be phrased as: *if S is a Bernstein set, then there exists a $\nu\mu_0$ so that every completion ...*

in sect. 3). If $K_{n,i}^{*t}(S)$ is not \mathbf{Q}' -scattered, then $K_{n,i}^{*t}(S)$ is not $\mathbf{Q} \vee \mathbf{Q}'$ -scattered; hence, working as in the proof of 3.1, we get a set of points $(\mathfrak{a}_i, c_i, u_i) \in K_{n,i}^{*t}(S)$ so that (c_i, u_i) form an oblique perfect set (u_i are dyadic irrationals). Since B is \mathcal{D} -2-Bernstein, there is a $(c_{i_0}, u_{i_0}) \in B \times D_i^t$ and the contradiction is obtained by looking at the point $(\mathfrak{a}_{i_0}, c_{i_0}, u_{i_0})$. \square

4.2. *Each $H_{\bar{x}, \bar{y}}$ is an intersection of clopen subsets of $\nu\mu$. Consequently, if F is a subspace of $\nu\mu$, then the quasicomponents of F are contained in its horizontal planks.*

Let \mathcal{P}_i , $i = 0, 1, \dots, 4$, stand, respectively, for $\text{ind} = 0$, totally disconnected (i.e., quasicomponents are one-point sets), hereditarily disconnected (i.e., components are one-point sets), pointlike (or punctiform – i.e., no non-trivial subcontinua), and “no non-trivial arcs”. It is known that for $i = 0, 1, 2, 3$ there exists a \mathbf{G}_δ -subset of the square \mathcal{J}^2 having $\neg\mathcal{P}_i$ and \mathcal{P}_{i+1} .

4.3. Theorem. a). *For $i = 0, 1, 2, 3$ there exists a metric space M with $\text{ind} = 0$ and such that every completion of M has $\neg\mathcal{P}_i$, but there exists a completion of M which has \mathcal{P}_{i+1} .*

b). *There exists a metric space M with $\text{ind} = 0$ and such that for every completion \tilde{M} of M , $\tilde{M} \setminus M$ has non-trivial components and hence $\text{ind}(\tilde{M} \setminus M) > 0$.*

Proof. In both cases we let $M = \nu\mu_0^2(S)$ for a suitable $S \subset \mathcal{J}^2$. For part a) we take a \mathbf{G}_δ -subset S of \mathcal{J}^2 with $\neg\mathcal{P}_i$ and \mathcal{P}_{i+1} . By 4.1, every completion of M has $\neg\mathcal{P}_i$, and, by 4.2, $\nu\mu^2(S)$ is a completion of M which has \mathcal{P}_{i+1} . For b) we take S so that both S and $\mathcal{J}^2 \setminus S$ are Bernstein in \mathcal{J}^2 and apply 2.3 and 4.1. \square

5. PROBLEMS

This is a very casual preview of problems arising from the present stage of development.

We now know that there exist, in ZFC, metric spaces for which neither dim nor ind_c can be determined in ZFC.³ It is interesting to explore the *degree* of indeterminacy of the dimension functions dim and ind_c as well as relations between

³As far as I know this phenomenon has not yet been observed within non-metrizable (completely regular) spaces – of course, for non-metrizable spaces we are concerned only with dim . This is somewhat curious inasmuch as pathology of dimension functions is far better documented within arbitrary spaces than within the metric ones. On the other hand, it is known that it is very easy to define spaces for which dim has not yet been determined and it is probably very difficult to determine or perhaps cannot be determined at all. For instance, given two topologies τ_1 and τ_2 for a set X , $\tau_1 \subset \tau_2$, and an τ_2 -closed set $A \subset X$, define the *intermediate* topology $\tau = (\tau_1|\tau_2, A)$ by the closure operation $\text{cl}_\tau(S) = (\text{cl}_{\tau_1} S \cap (X \setminus A)) \cup \text{cl}_{\tau_2} S$ for every $S \subset X$. The simplest instance of this procedure is, for a given space X and an arbitrary $A \subset X$, to consider $\tau = (\tau_1|\tau_2, A)$, where τ_1 is the original topology of X and τ_2 is the discrete one; X with this τ is denoted by X_A . Even such a primitive operation leads to serious problems concerning dim . Let A_1, A_2, \dots , be subsets of the Cantor set \mathcal{C} and consider the product space $\prod_n \mathcal{C}_{A_n}$. One would expect that $\prod_{n=1}^m \mathcal{C}_{A_n}$ is hereditarily strongly 0-dimensional; however, the best I can do at the present moment is to prove that $\text{dim} \mathcal{C}_A^2 = 0$ assuming that A is not too pathological. The possibility is that $\text{dim} \mathcal{C}_A^2$ might depend upon set-theoretic assumptions, but at that time I did not pursue the matters further. Now it looks more interesting; in any case, I would like to challenge the masters of covering dimension to do better than I did. The infinite product $\prod_n \mathcal{C}_{A_n}$ is probably strongly 0-dimensional but not hereditarily. (Incidentally, the intermediate topology, as well as σ -non-archimedean bases, are discussed in the Ph.D. dissertation of my former student H. P. Tan.)

these functions. One would expect that this investigation would bring formulation of new set-theoretic principles possibly far more intricate than $\mathbf{S}(\aleph_0)$. The most obvious (and probably hopelessly difficult) question is: *does there exist a 0-dimensional metric space X such that for every $m, n = 0, 1, \dots, \infty$, $n \leq M$ the statement $(\text{ind}_c X = n)$ and $(\text{dim } X = m)$ is consistent with ZFC?*

$\nu\mu_0$ itself raises further consistency questions. According to 1.2, both of the systems of equalities

$$(1) \quad \text{ind}_c \nu\mu_0 = 0, \quad \text{dim } \nu\mu_0^2 = 1;$$

$$(2) \quad \text{ind}_c \nu\mu_0 = 1, \quad \text{dim } \nu\mu_0^2 = 2.$$

are consistent with ZFC. Are the remaining systems of equalities

$$(1') \quad \text{ind}_c \nu\mu_0 = 1, \quad \text{dim } \nu\mu_0^2 = 1;$$

$$(2') \quad \text{ind}_c \nu\mu_0 = 0, \quad \text{dim } \nu\mu_0^2 = 2$$

consistent with ZFC?

Irreducibility (for ease of formulation we will consider only spaces with $\text{ind} = 0$): Kulesza [Ku1] and Ostaszewski [Ost] produced (in ZFC) esoteric spaces of weight \aleph_1 ; such spaces are automatically irreducible. Kulesza [Ku2] has also produced (in ZFC) an irreducible esoteric space of weight 2^{\aleph_0} and obtained consistency results concerning spaces of higher weight. But it is not known whether for every \mathfrak{m} with $\aleph_1 < \mathfrak{m} < 2^{\aleph_0}$ there is, in ZFC, an irreducible esoteric space of weight \mathfrak{m} .

One of the possible modifications of the irreducibility is the condition⁴ that every closed subset F of a space X with $\text{weight } F < \text{weight } X$ has a base of clopen sets (i.e., $\text{Ind}_A X = 0$). Some of the results of Kulesza [Ku2] refer to this type of irreducibility, and it is my understanding that he has further results in this direction.

Irreducibility can also be applied to ind_c rather than to dim (or Ind). Here an obvious question is: *does there exist a space X with $\text{ind}_c > \text{ind}$ (of weight, say, 2^{\aleph_0}) which is irreducible relative to ind_c but reducible relative to dim – i.e., for every subspace X' of X with $\text{weight } X' < \text{weight } X$ we have $\text{ind}_c X' = 0$, but there exists a subspace X_0 with $\text{weight } X_0 < \text{weight } X$ and $\text{dim } X_0 > 0$?* It is possible that the statement that $\nu\mu_0$ has this property is consistent, but this, of course, cannot be settled overnight.

REFERENCES

- [Dou] R. Dougherty, *Narrow coverings of ω -ary product spaces*, Annals of Pure and Applied Logic **88** (1997), 47-91. CMP 98:03
- [Ku1] John Kulesza, *An example in the dimension theory of metrizable spaces*, Topology and its appl. **35** (1990), 109 - 120. MR **91g**:54045
- [Ku2] John Kulesza, *Metrizable spaces where the inductive dimensions disagree*, Trans. A.M.S. **318**, no. **2** . (1990), 763–781. MR **90g**:54030
- [M1] S. Mrówka, *Further results on E -compact spaces*, Acta Math. **120** (1968), 161 - 185. MR **37**:2165
- [M2] ———, *Small inductive dimension of completions of metric spaces*, Proc. Amer. Math. Soc. **125** (5) (1997), 1545 - 1554. MR **97i**:54043
- [M3] ———, *N -compactness, metrizability and covering dimension*, Rings of continuous functions, Marcell Dekker, Inc., New York and Basel, 1985, pp. 248 - 275. MR **86i**:54034

⁴This could be called *-irreducibility, but since other modifications of irreducibility are likely to appear, I am reluctant to introduce firm terminology at this time.

- [Ost] A. Ostaszewski, *A note on the Prabir Roy space*, Topology Appl. **35**, no. **2-3**, (1990), 95–107. MR **91j**:54060
- [R] P. Roy, *Failure of equivalence of dimension concepts for metric spaces*, Bull. A.M.S. **68** (1962), 609 - 613. MR **25**:5495

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK AT BUFFALO, 134 DE-
FENDORF HALL, BUFFALO, NEW YORK 14224

E-mail address: `mrowka@acsu.buffalo.edu`