

AN OBSTRUCTION TO THE CONFORMAL COMPACTIFICATION OF RIEMANNIAN MANIFOLDS

SEONGTAG KIM

(Communicated by Christopher Croke)

ABSTRACT. In this paper, we study noncompact complete Riemannian n -manifolds with $n \geq 3$ which are not pointwise conformal to subdomains of any compact Riemannian n -manifold. For this, we compare the Sobolev Quotient at infinity of a noncompact complete Riemannian manifold with that of the singular set in a compact Riemannian manifold using the method for the Yamabe problem.

1. INTRODUCTION

In this paper, we study noncompact complete Riemannian n -manifolds with $n \geq 3$ which are not pointwise conformal to subdomains of any compact Riemannian n -manifold. We show that the conformal structure at infinity provides an obstruction to the conformal compactification of certain noncompact complete Riemannian manifolds. This obstruction is interesting because Nomizu and Ozeki [6] showed that any Riemannian manifold admits a complete metric by the conformal change of metric.

First, we calculate the Sobolev Quotient of a singular set in a given compact Riemannian manifold using the method which was developed for the Yamabe problem. Second, we compare the Sobolev Quotient at infinity of a given noncompact complete Riemannian manifold with the Sobolev Quotient of a singular set in a compact Riemannian manifold. To do this we introduce some notation:

$$Q(M, g) \equiv \inf_{u \in C_0^\infty(M)} \frac{\int_M |\nabla u|^2 + C_n S u^2 dV_g}{\left(\int_M u^{2n/(n-2)} dV_g \right)^{(n-2)/n}}$$

and

$$\overline{Q(M, g)} \equiv \lim_{r \rightarrow \infty} \inf_{u \in C_0^\infty(M-B_r)} \frac{\int_{M-B_r} |\nabla u|^2 + C_n S u^2 dV_g}{\left(\int_{M-B_r} u^{2n/(n-2)} dV_g \right)^{(n-2)/n}},$$

Received by the editors July 22, 1998.

1991 *Mathematics Subject Classification*. Primary 53C21; Secondary 58G30.

Key words and phrases. Scalar curvature, Yamabe problem, conformal metric.

The author was supported in part by KOSEF96070102013 and BSRI 97-1419 Ministry of Education.

where $C_n = \frac{n-2}{4(n-1)}$, S is the scalar curvature of (M, g) and B_r is the ball of radius r with center at a fixed point x_0 .

We also use the notation $Q(M)$ for $Q(M, g)$ and $Q(S^n)$ for $Q(S^n, g_0)$, where g_0 is the standard metric. Note that $\overline{Q(M, g)}$ is well defined since

$$\begin{aligned} & \inf_{u \in C_0^\infty(M-B_r)} \frac{\int_{M-B_r} |\nabla u|^2 + C_n S u^2 dV_g}{\left(\int_{M-B_r} u^{2n/(n-2)} dV_g \right)^{(n-2)/n}} \\ & \leq \inf_{u \in C_0^\infty(M-B_{r+1})} \frac{\int_{M-B_{r+1}} |\nabla u|^2 + C_n S u^2 dV_g}{\left(\int_{M-B_{r+1}} u^{2n/(n-2)} dV_g \right)^{(n-2)/n}}. \end{aligned}$$

$\overline{Q(M, g)}$ was used for a study of the Yamabe problem on noncompact complete Riemannian manifolds by Kim [4]. Obstructions for the existence of a complete conformal metric with prescribed scalar curvature on a subdomain $K - \Gamma$ of a compact Riemannian manifold (K, h) were studied by Aviles and McOwen [2], Schoen and Yau [8], Delanoe [3] and McOwen [5] using the dimension of Γ . In this paper, we have an obstruction for the conformal compactification of a noncompact complete Riemannian manifold using the conformal structure at infinity.

2. THE SOBOLEV QUOTIENT OF A SINGULAR SET

In this section, we show that the Sobolev Quotient of a singular set is the same as that of the standard sphere. Let us denote the n dimensional Riemannian volume of a subset Ω of (M, g) by $|\Omega|$. We state our theorem.

Theorem 2.1. *Let (M, g) be a Riemannian manifold of dimension $n \geq 3$. Assume that there exists a sequence $\{\Gamma_i\}$ of smooth bounded open subsets of (M, g) with $|\Gamma_i| \rightarrow 0$ and $\Gamma_i \subset \Gamma_1$ for $i = 1, 2, \dots$. Then we have $\lim_{i \rightarrow \infty} Q(\Gamma_i) = Q(S^n)$.*

Proof. Assume that there exists a subsequence $\{\Gamma_{i_k}\}$ of $\{\Gamma_i\}$ such that

$$(2.1) \quad \lim_{i \rightarrow \infty} Q(\Gamma_{i_k}) < Q(S^n).$$

We use the same notation $\{\Gamma_i\}$ for this subsequence. By the work on the Yamabe problem in the compact case (see [9], [1], [7], and [8]), there exists a positive solution $u_i \in C^\infty(\Gamma_i)$ with

$$(2.2) \quad -\Delta u_i + C_n S u_i = q_i u_i^{(n+2)/(n-2)} \text{ on } \Gamma_i,$$

$$\int_{\Gamma_i} u_i^{2n/(n-2)} dV_g = 1 \text{ and } u_i = 0 \text{ on } \partial\Gamma_i,$$

where $q_i = Q(\Gamma_i)$. We extend the domain of u_i by defining $u_i = 0$ on the outside of Γ_i . By multiplying (2.2) by u_i^{1+2b} where $b > 0$ and integrating over Γ_i ,

$$(2.3) \quad q_i \int_{\Gamma_i} u_i^{2n/(n-2)+2b} dV_g = \int_{\Gamma_i} \frac{1+2b}{(1+b)^2} |\nabla u_i^{1+b}|^2 + C_n S u_i^{2+2b} dV_g.$$

Using Hölder’s inequality,

$$\begin{aligned}
 (2.4) \quad & \int_{\Gamma_i} u_i^{2n/(n-2)+2b} dV_g \\
 & \leq \left(\int_{\Gamma_i} u_i^{(1+b)2n/(n-2)} dV_g \right)^{\frac{n-2}{n}} \left(\int_{\Gamma_i} u_i^{(2n/(n-2)+2b-(1+b)2)\frac{n}{2}} dV_g \right)^{\frac{2}{n}} \\
 & \leq \left(\int_{\Gamma_i} u_i^{(1+b)2n/(n-2)} dV_g \right)^{\frac{n-2}{n}}.
 \end{aligned}$$

By the Sobolev Embedding Theorem on Riemannian manifolds for u_i on Γ_i , $\Gamma_i \subset \Gamma_1 \subset M$ (note $\bar{\Gamma}_1$ is compact), and (2.3), for any given $\epsilon > 0$, there exists $C(\epsilon)$ with

$$\begin{aligned}
 (2.5) \quad & \left(\int_{\Gamma_i} u_i^{(1+b)2n/(n-2)} dV_g \right)^{\frac{n-2}{n}} \\
 & \leq (1 + \epsilon) \frac{1}{Q(S^n)} \int_{\Gamma_i} |\nabla u_i^{1+b}|^2 dV_g + C(\epsilon) \int_{\Gamma_i} u_i^{2+2b} dV_g \\
 & \leq (1 + \epsilon) \frac{1}{Q(S^n)} \frac{(1 + b)^2}{(1 + 2b)} \left(\int_{\Gamma_i} q_i u_i^{2n/(n-2)+2b} - C_n \int_{\Gamma_i} S u_i^{2+2b} dV_g \right) \\
 & \quad + C(\epsilon) \int_{\Gamma_i} u_i^{2+2b} dV_g.
 \end{aligned}$$

From (2.4),

$$\begin{aligned}
 (2.6) \quad & \left(\int_{\Gamma_i} u_i^{(1+b)2n/(n-2)} dV_g \right)^{\frac{n-2}{n}} \\
 & \leq (1 + \epsilon) \frac{1}{Q(S^n)} \frac{(1 + b)^2}{(1 + 2b)} \left(q_i \left(\int_{\Gamma_i} u_i^{(1+b)2n/(n-2)} dV_g \right)^{\frac{n-2}{n}} \right. \\
 & \quad \left. - C_n \int_{\Gamma_i} S u_i^{2+2b} dV_g \right) + C(\epsilon) \int_{\Gamma_i} u_i^{2+2b} dV_g.
 \end{aligned}$$

Since $q_i < c < Q(S^n)$ for some c , we can take $\epsilon > 0$ and $0 < b < \frac{2}{n-2}$ so that

$$(2.7) \quad (1 + \epsilon) \frac{q_i}{Q(S^n)} \frac{(1 + b)^2}{(1 + 2b)} < c_0 < 1$$

for some constant c_0 . Therefore, we have

$$\begin{aligned}
 (2.8) \quad & \left(1 - (1 + \epsilon) \frac{q_i}{Q(S^n)} \frac{(1 + b)^2}{(1 + 2b)} \right) \left(\int_{\Gamma_i} u_i^{(1+b)2n/(n-2)} dV_g \right)^{\frac{n-2}{n}} \\
 & + C \int_{\Gamma_i} S u_i^{2+2b} dV_g \\
 & \leq C(\epsilon) \int_{\Gamma_i} u_i^{2+2b} dV_g,
 \end{aligned}$$

where C is a general positive constant independent of i . We may assume that $|\Gamma_i| < 1$. Using Hölder’s inequality,

$$\begin{aligned} \int_{\Gamma_i} u_i^{2+2b} dV_g &\leq \left(\int_{\Gamma_i} u_i^{2n/(n-2)} dV_g \right)^{t_1} |\Gamma_i|^{1-t_1} \\ &\leq |\Gamma_i|^{1-t_1} < 1, \end{aligned}$$

where $t_1 = \frac{(1+b)(n-2)}{n}$.

Finally, we have

$$(2.9) \quad \left(\int_M u_i^{(1+b)2n/(n-2)} dV_g \right)^{\frac{n-2}{n}} = \left(\int_{\Gamma_i} u_i^{(1+b)2n/(n-2)} dV_g \right)^{\frac{n-2}{n}} \leq A_1,$$

where A_1 does not depend on i .

From (2.9), we have

$$(2.10) \quad \begin{aligned} 1 = \int_{\Gamma_i} u_i^{2n/(n-2)} dV_g &\leq \left(\int_{\Gamma_i} u_i^{(1+b)2n/(n-2)} dV_g \right)^t |\Gamma_i|^s \\ &\leq A_1^{nt/(n-2)} |\Gamma_i|^s, \end{aligned}$$

where $t = \frac{1}{1+b}$ and $s = \frac{b}{1+b}$.

Since $|\Gamma_i| \rightarrow 0$, (2.10) does not hold. This contradiction comes from (2.1). Thus Theorem 2.1 is proved. □

3. APPLICATIONS ON NONCOMPACT COMPLETE RIEMANNIAN MANIFOLDS

Using Theorem 2.1, we show that there exists an obstruction to the conformal compactification of certain noncompact complete Riemannian manifolds.

Theorem 3.1. *Let (M, g) be a noncompact complete Riemannian manifold of dimension $n \geq 3$ with $Q(M, g) < Q(S^n)$. Then (M, g) is not pointwise conformal to a subdomain of any compact Riemannian n -manifold.*

Proof. Assume that (M, g) is pointwise conformal to a subdomain $(M, u^{4/(n-2)}g)$ in (K, h) , where (K, h) is a compact Riemannian n -manifold. Take smooth compact domain X_i in M with $X_i \subset \overline{X_i} \subset X_{i+1}$ so that $\text{Vol}(M - X_i, u^{4/(n-2)}g) \rightarrow 0$ as $i \rightarrow \infty$, where $\text{Vol}(M - X_i, u^{4/(n-2)}g)$ is the volume of $M - X_i$ with the metric $u^{4/(n-2)}g$. We let $|\cdot|_h$ ($|\cdot|_g$) be the length of the tangent vector and S_h (S_g) be the scalar curvature with the metric h (respectively g).

Since we assume that $Q(M, g) < Q(S^n)$, there exists a smooth function $f_i \in C_0^\infty(M - \overline{X_i})$ with

$$\frac{\int_{M-\overline{X_i}} |\nabla f_i|_g^2 + C_n S_g f_i^2 dV_g}{\left(\int_{M-\overline{X_i}} f_i^{2n/(n-2)} dV_g \right)^{(n-2)/n}} < Q(S^n) - c_0$$

for a positive constant c_0 and $i = 1, 2, \dots$.

We take a smooth bounded open subset Y_i of M with $X_i \subset \overline{X_i} \subset Y_i$ and

$$Z_i \subset Y_i - X_i \subset M - X_i,$$

where Z_i is the support of f_i . By the conformal invariance of $Q(Y_i - \overline{X_i}, h)$,

$$\begin{aligned}
 (3.1) \quad Q(Y_i - \overline{X_i}, h) &= Q(Y_i - \overline{X_i}, u^{4/(n-2)}g) \\
 &\leq \frac{\int_{Y_i - \overline{X_i}} |\nabla(f_i/u)|_h^2 + C_n S_h (f_i/u)^2 dV_h}{\left(\int_{Y_i - \overline{X_i}} (f_i/u)^{2n/(n-2)} dV_h\right)^{(n-2)/n}} \\
 &\leq \frac{\int_{Y_i - \overline{X_i}} |\nabla f_i|_g^2 + C_n S_g f_i^2 dV_g}{\left(\int_{Y_i - \overline{X_i}} f_i^{2n/(n-2)} dV_g\right)^{(n-2)/n}} \\
 &\leq Q(S^n) - c_0.
 \end{aligned}$$

Since the $\text{Vol}(Y_i - \overline{X_i}, u^{4/(n-2)}g) \rightarrow 0$, (3.1) contradicts Theorem 2.1. Theorem 3.1 is proved. \square

Using Theorem 3.1, we have the following.

Corollary 3.2. *Let (M, g) be a noncompact complete Riemannian manifold of dimension $n \geq 3$. Assume that there exists a positive number c_0 and a sequence $\{\Omega_i\}$ of smooth open sets where $\Omega_i \subset M - B_i(x_0)$ for $i = 1, 2, \dots$ and $Q(\Omega_i) < Q(S^n) - c_0$. Then (M, g) is not pointwise conformal to a subdomain of any compact Riemannian n -manifold.*

Example 3.3. Let $B(p, r)$ be the Euclidean ball of radius r with center p in (R^n, δ_{ij}) with $n \geq 3$. Take any smooth metric h on $B(p, 2)$ so that $h = \delta_{ij}$ on $B(p, 2) - B(p, 1)$ and h is not conformally flat at the center p . Then $Q(B(p, 2), h) < Q(S^n)$ by the resolution of the Yamabe problem. Consider (R^n, g) with $n \geq 3$, where $g = h$ on each Euclidean balls $B((5n, 0), 2)$ with center $(5n, 0) \in R \times R^{n-1}$ and radius 2 in the Euclidean norm, for $n = 1, 2, 3, \dots$ and $g = \delta_{ij}$ on $R^n - \bigcup B((5n, 0), 2)$. Then (R^n, g) is not pointwise conformal to a subdomain of any compact Riemannian n -manifold, since $Q(R^n, g) < Q(S^n)$.

ACKNOWLEDGEMENTS

The author thanks Patricio Aviles for his encouragement on this work.

REFERENCES

1. T. Aubin, *Equations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire*, J. Math. Pures Appl. n **55** (1976) 269-296. MR **55**:4288
2. P. Aviles and R. McOwen, *Complete conformal metrics with negative scalar curvature in compact Riemannian manifolds*, Duke Math. J. **56** (1988) 225-239. MR **89b**:58224
3. P. Delanoe, *Generalized stereographic projections with prescribed scalar curvature*, Geometry and nonlinear partial differential equations, (Contemp. Math., vol 127 pp. 17-25) Providence, A.M.S. 1992 MR **93e**:53045
4. S. Kim, *Scalar curvature on noncompact complete Riemannian manifolds*, Nonlinear Analysis **26** (1996) 1985-1993. MR **97a**:53056
5. R. McOwen, *Singularities and the conformal scalar curvature equation*, Geometry and nonlinear partial differential equations (Contemp. Math., vol 127 pp. 221-233) Providence, A.M.S. 1992 MR **94b**:53076
6. K. Nomizu and H. Ozeki, *The existence of complete Riemannian metrics*, Proc. Amer. Math. Soc. **12** (1961) 889-891. MR **24**:A3610
7. R. Schoen, *Conformal deformation of a Riemannian metric to constant scalar curvature*, J. Diff. Geom. **20** (1984) 479-495. MR **86i**:58137

8. R. Schoen and S. Yau, *Conformally flat manifolds, Kleinian groups and scalar curvature*, Invent. Math. **92** (1988) 47-71. MR **89c**:58139
9. H. Yamabe, *On a deformation of Riemannian structures on compact manifolds*, Osaka Math. J. **12** (1960) 21-37. MR **23**:A2847

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON 440-746, SOUTH KOREA
E-mail address: `stkim@yurim.skku.ac.kr`