

CLASS NUMBERS OF IMAGINARY ABELIAN NUMBER FIELDS

KU-YOUNG CHANG AND SOUN-HI KWON

(Communicated by David E. Rohrlich)

ABSTRACT. Let N be an imaginary abelian number field. We know that h_N^- , the relative class number of N , goes to infinity as f_N , the conductor of N , approaches infinity, so that there are only finitely many imaginary abelian number fields with given relative class number. First of all, we have found all imaginary abelian number fields with relative class number one: there are exactly 302 such fields. It is known that there are only finitely many CM-fields N with cyclic ideal class groups of 2-power orders such that the complex conjugation is the square of some automorphism of N . Second, we have proved in this paper that there are exactly 48 such fields.

1. INTRODUCTION

Let N be an imaginary abelian number field of conductor f_N , maximal real subfield N^+ , and relative class number h_N^- . It is known that h_N^- goes to infinity as f_N approaches infinity, so that there exist only finitely many number fields N with $h_N^- = 1$ ([S1, Theorem 2]). It is interesting to find all imaginary number fields with relative class number one. Yamamura has determined all imaginary abelian number fields with class number one. One of the purposes of this paper is to prove the following:

Theorem 1. *There are exactly 302 imaginary abelian number fields with relative class number one: 243 out of them are non-cyclic number fields. Their degrees are less than or equal to 24 and their conductors are less than or equal to 65689. These fields are given in Table I.*

Louboutin [Lou1] has proved that there are only finitely many CM-fields N with cyclic ideal class groups of 2-power orders such that the complex conjugation is the square of some automorphism of N and he also proved that the relative class numbers of such fields are equal to 1 or 2. Furthermore he determined the non-quadratic imaginary cyclic number fields of 2-power degrees with cyclic ideal class groups of 2-power orders. Our second goal of this paper is to determine all imaginary abelian number fields with relative class numbers less than or equal to 4 such that the complex conjugation is the square of some automorphism of N :

Received by the editors May 1, 1998.

1991 *Mathematics Subject Classification.* Primary 11R29; Secondary 11R20.

This research was supported by Grant BSRI-97-1408 from the Ministry of Education of Korea.

TABLE I. The imaginary non-cyclic abelian number fields with relative class number one. $\chi_7(3) = e^{2\pi i/6}$, $\chi_{13}(2) = e^{2\pi i/12}$, $\psi_9(2) = e^{2\pi i/3}$

Type	Field N (h_{N^+}, Q_N)
$(2^*, 3, 3)$	$\langle \chi_3, \chi_7^2, \psi_9 \rangle (1, 1)$
$(2^*, 2^*)$	$\langle f_{k_1}, f_{k_2} \rangle : k_1 = \mathbb{Q}(\sqrt{-m_1}), k_2 = \mathbb{Q}(\sqrt{-m_2})$ $\langle 3, 4 \rangle(1, 2), \langle 3, 7 \rangle(1, 2), \langle 3, 8 \rangle(1, 2), \langle 3, 11 \rangle(1, 2)$ $\langle 3, 19 \rangle(1, 2), \langle 3, 43 \rangle(1, 2), \langle 3, 67 \rangle(1, 2), \langle 3, 163 \rangle(1, 2)$ $\langle 4, 7 \rangle(1, 2), \langle 4, 8 \rangle(1, 1), \langle 4, 11 \rangle(1, 2), \langle 4, 19 \rangle(1, 2)$ $\langle 4, 43 \rangle(1, 2), \langle 4, 67 \rangle(1, 2), \langle 4, 163 \rangle(1, 2), \langle 7, 8 \rangle(1, 2)$ $\langle 7, 11 \rangle(1, 2), \langle 7, 19 \rangle(1, 2), \langle 7, 43 \rangle(1, 2), \langle 7, 67 \rangle(3, 2)$ $\langle 7, 163 \rangle(1, 2), \langle 8, 11 \rangle(1, 2), \langle 8, 19 \rangle(1, 2), \langle 8, 43 \rangle(1, 2)$ $\langle 8, 67 \rangle(1, 2), \langle 8, 163 \rangle(3, 2), \langle 11, 19 \rangle(1, 2), \langle 11, 43 \rangle(3, 2)$ $\langle 11, 67 \rangle(1, 2), \langle 11, 163 \rangle(1, 2), \langle 19, 43 \rangle(5, 2), \langle 19, 67 \rangle(1, 2)$ $\langle 19, 163 \rangle(1, 2), \langle 43, 67 \rangle(1, 2), \langle 43, 163 \rangle(1, 2)$ $\langle 67, 163 \rangle(1, 2), \langle 3, 15 \rangle(1, 1), \langle 3, 20 \rangle(2, 1), \langle 3, 24 \rangle(1, 1)$ $\langle 3, 35 \rangle(2, 1), \langle 3, 40 \rangle(2, 1), \langle 3, 51 \rangle(1, 1), \langle 3, 88 \rangle(2, 1)$ $\langle 3, 115 \rangle(2, 1), \langle 3, 123 \rangle(1, 1), \langle 3, 187 \rangle(2, 1), \langle 3, 232 \rangle(2, 1)$ $\langle 3, 235 \rangle(2, 1), \langle 3, 267 \rangle(1, 1), \langle 4, 15 \rangle(2, 1), \langle 4, 20 \rangle(1, 1)$ $\langle 4, 35 \rangle(2, 1), \langle 4, 40 \rangle(2, 1), \langle 4, 52 \rangle(1, 1), \langle 4, 91 \rangle(2, 1)$ $\langle 4, 115 \rangle(2, 1), \langle 4, 148 \rangle(1, 1), \langle 4, 232 \rangle(2, 1), \langle 4, 235 \rangle(6, 1)$ $\langle 4, 403 \rangle(2, 1), \langle 4, 427 \rangle(6, 1), \langle 7, 15 \rangle(2, 1), \langle 7, 20 \rangle(2, 1)$ $\langle 7, 35 \rangle(1, 1), \langle 7, 40 \rangle(2, 1), \langle 7, 51 \rangle(2, 1), \langle 7, 52 \rangle(2, 1)$ $\langle 7, 91 \rangle(1, 1), \langle 7, 115 \rangle(2, 1), \langle 7, 123 \rangle(2, 1), \langle 7, 187 \rangle(2, 1)$ $\langle 7, 235 \rangle(2, 1), \langle 7, 267 \rangle(2, 1), \langle 7, 403 \rangle(2, 1), \langle 7, 427 \rangle(1, 1)$ $\langle 8, 15 \rangle(2, 1), \langle 8, 20 \rangle(2, 1), \langle 8, 35 \rangle(2, 1), \langle 8, 40 \rangle(1, 1)$ $\langle 8, 52 \rangle(2, 1), \langle 8, 91 \rangle(2, 1), \langle 8, 115 \rangle(2, 1), \langle 8, 148 \rangle(2, 1)$ $\langle 8, 232 \rangle(1, 1), \langle 8, 235 \rangle(2, 1), \langle 8, 403 \rangle(2, 1), \langle 8, 427 \rangle(2, 1)$ $\langle 11, 24 \rangle(2, 1), \langle 11, 51 \rangle(2, 1), \langle 11, 52 \rangle(2, 1), \langle 11, 88 \rangle(1, 1)$ $\langle 11, 91 \rangle(2, 1), \langle 11, 123 \rangle(2, 1), \langle 11, 187 \rangle(1, 1)$ $\langle 11, 232 \rangle(2, 1), \langle 11, 403 \rangle(2, 1), \langle 11, 427 \rangle(2, 1)$ $\langle 19, 24 \rangle(2, 1), \langle 19, 52 \rangle(2, 1), \langle 19, 88 \rangle(2, 1), \langle 19, 91 \rangle(2, 1)$ $\langle 19, 123 \rangle(2, 1), \langle 19, 148 \rangle(2, 1), \langle 19, 232 \rangle(2, 1)$ $\langle 19, 267 \rangle(6, 1), \langle 19, 403 \rangle(2, 1), \langle 43, 15 \rangle(2, 1)$ $\langle 43, 20 \rangle(2, 1), \langle 43, 24 \rangle(2, 1), \langle 43, 35 \rangle(2, 1)$ $\langle 43, 40 \rangle(2, 1), \langle 43, 88 \rangle(2, 1), \langle 43, 115 \rangle(2, 1)$ $\langle 43, 148 \rangle(2, 1), \langle 43, 232 \rangle(2, 1), \langle 43, 235 \rangle(2, 1)$ $\langle 43, 267 \rangle(2, 1), \langle 43, 427 \rangle(2, 1), \langle 67, 15 \rangle(2, 1)$ $\langle 67, 20 \rangle(2, 1), \langle 67, 24 \rangle(2, 1), \langle 67, 35 \rangle(2, 1)$ $\langle 67, 40 \rangle(2, 1), \langle 67, 52 \rangle(2, 1), \langle 67, 88 \rangle(2, 1)$ $\langle 67, 91 \rangle(14, 1), \langle 67, 115 \rangle(10, 1), \langle 67, 123 \rangle(2, 1)$ $\langle 67, 235 \rangle(2, 1), \langle 67, 403 \rangle(2, 1), \langle 67, 427 \rangle(14, 1)$ $\langle 163, 15 \rangle(2, 1), \langle 163, 20 \rangle(2, 1), \langle 163, 24 \rangle(2, 1)$ $\langle 163, 35 \rangle(2, 1), \langle 163, 40 \rangle(2, 1), \langle 163, 51 \rangle(2, 1)$ $\langle 163, 52 \rangle(2, 1), \langle 163, 88 \rangle(2, 1), \langle 163, 91 \rangle(2, 1)$ $\langle 163, 115 \rangle(2, 1), \langle 163, 148 \rangle(2, 1), \langle 163, 187 \rangle(2, 1)$ $\langle 163, 232 \rangle(2, 1), \langle 163, 235 \rangle(2, 1), \langle 163, 267 \rangle(2, 1)$ $\langle 163, 403 \rangle(2, 1)$

Type	Field N (h_{N^+}, Q_N)
$(2^*, 2^*, 3)$	$\langle \chi_7^3, \chi_3, \chi_7^2 \rangle (1, 2), \langle \chi_7^3, \chi_4, \chi_7^2 \rangle (1, 2)$ $\langle \chi_7^3, \chi_5^2 \chi_7^3, \chi_7^2 \rangle (1, 1), \langle \chi_7^3, \chi_4 \psi_8, \chi_7^2 \rangle (1, 2)$ $\langle \chi_7^3, \chi_{11}^5, \chi_7^2 \rangle (1, 2), \langle \chi_7^3, \chi_3 \chi_5^2, \chi_7^2 \rangle (2, 1)$ $\langle \chi_3, \chi_4, \chi_7^2 \rangle (1, 2), \langle \chi_3, \chi_5^2 \chi_7^3, \chi_7^2 \rangle (2, 1)$ $\langle \chi_3, \chi_4 \psi_8, \chi_7^2 \rangle (1, 2), \langle \chi_3, \chi_{11}^5, \chi_7^2 \rangle (1, 2)$ $\langle \chi_3, \chi_3 \chi_5^2, \chi_7^2 \rangle (1, 1), \langle \chi_4, \chi_5^2 \chi_7^3, \chi_7^2 \rangle (2, 1)$ $\langle \chi_4, \chi_4 \psi_8, \chi_7^2 \rangle (1, 1), \langle \chi_4, \chi_{11}^5, \chi_7^2 \rangle (1, 2)$ $\langle \chi_4, \chi_3 \chi_5^2, \chi_7^2 \rangle (2, 1), \langle \chi_5^2 \chi_7^3, \chi_4 \psi_8, \chi_7^2 \rangle (2, 1)$ $\langle \chi_4 \psi_8, \chi_{11}^5, \chi_7^2 \rangle (3, 2), \langle \chi_4 \psi_8, \chi_3 \chi_5^2, \chi_7^2 \rangle (2, 1)$ $\langle \chi_3, \chi_4, \psi_9 \rangle (1, 2), \langle \chi_3, \chi_3 \chi_5^2, \psi_9 \rangle (1, 1)$ $\langle \chi_3, \chi_7^3, \psi_9 \rangle (1, 2), \langle \chi_3, \chi_3 \psi_8, \psi_9 \rangle (1, 1)$ $\langle \chi_4, \chi_3 \chi_5^2, \psi_9 \rangle (2, 1), \langle \chi_4, \chi_7^3, \psi_9 \rangle (1, 2)$ $\langle \chi_3 \chi_5^2, \chi_7^3, \psi_9 \rangle (2, 1), \langle \chi_3, \chi_7^3, \chi_{13}^4 \rangle (1, 2)$ $\langle \chi_3, \chi_4 \psi_8, \chi_{13}^4 \rangle (1, 2), \langle \chi_4 \chi_{13}^6, \chi_7^3, \chi_{13}^4 \rangle (2, 1)$ $\langle \chi_4 \chi_{13}^6, \chi_4 \psi_8, \chi_{13}^4 \rangle (2, 1), \langle \chi_7^3, \chi_4 \psi_8, \chi_{13}^4 \rangle (1, 2)$ $\langle \chi_4, \chi_{19}^9, \chi_{19}^6 \rangle (1, 2), \langle \chi_3, \chi_{43}^{21}, \chi_{43}^{14} \rangle (1, 2)$ $\langle \chi_3, \chi_7^3, \chi_7^4 \psi_9 \rangle (3, 2), \langle \chi_7^3, \chi_7^3 \chi_{13}^6, \chi_7^4 \chi_{13}^4 \rangle (3, 1)$
$(2^*, 2^*, 5)$	$\langle \chi_3, \chi_{11}^5, \chi_{11}^2 \rangle (1, 2), \langle \chi_3, \chi_4, \chi_{11}^2 \rangle (1, 2), \langle \chi_4, \chi_{11}^5, \chi_{11}^2 \rangle (1, 2)$
$(4^*, 2^*)$	$\langle \chi_5, \chi_3 \rangle (1, 2), \langle \chi_5, \chi_4 \rangle (1, 2), \langle \chi_5, \chi_7^3 \rangle (1, 2)$ $\langle \chi_5, \chi_4 \psi_8 \rangle (1, 2), \langle \chi_{13}^3, \chi_4 \rangle (1, 2), \langle \chi_{13}^3, \chi_7^3 \rangle (1, 2)$ $\langle \chi_4 \psi_{16}, \chi_3 \rangle (1, 2), \langle \chi_4 \psi_{16}, \chi_4 \rangle (1, 1), \langle \chi_4 \psi_{16}, \chi_{11}^5 \rangle (1, 2)$ $\langle \chi_4 \psi_{16}, \chi_4 \chi_5^2 \rangle (2, 1), \langle \chi_{37}^9, \chi_4 \rangle (1, 2), \langle \chi_{29}^7, \chi_4 \psi_8 \rangle (1, 2)$ $\langle \chi_4 \psi_{16} \chi_5^2, \chi_4 \rangle (2, 1), \langle \chi_3 \psi_{16}, \chi_3 \rangle (1, 1)$ $\langle \chi_3 \psi_{16}, \chi_{11}^5 \rangle (2, 1), \langle \chi_7^3 \chi_{17}^4, \chi_3 \rangle (2, 1)$ $\langle \chi_7^3 \chi_{17}^4, \chi_{11}^5 \rangle (2, 1), \langle \chi_{61}^{15}, \chi_7^3 \rangle (5, 2)$
$(4^*, 2^*, 3)$	$\langle \chi_5, \chi_7^3, \chi_7^2 \rangle (1, 2), \langle \chi_5, \chi_3, \psi_9 \rangle (1, 2), \langle \chi_5, \chi_3, \chi_7^2 \rangle (1, 2)$
$(8^*, 2^*)$	$\langle \chi_4 \psi_{32}, \chi_4 \rangle (1, 1), \langle \chi_4 \psi_{32}, \chi_3 \rangle (1, 2), \langle \chi_4 \psi_{32}, \chi_4 \chi_5^2 \rangle (2, 1)$
$(2^*, 2^*, 2^*)$	$(f_{k_1}, f_{k_2}, f_{k_3}) : k_1 = \mathbb{Q}(\sqrt{-m_1}), k_2 = \mathbb{Q}(\sqrt{-m_2}), k_3 = \mathbb{Q}(\sqrt{-m_3})$ $\langle 3, 4, 7 \rangle (1, 2), \langle 3, 4, 8 \rangle (1, 2), \langle 3, 4, 11 \rangle (1, 2)$ $\langle 3, 4, 15 \rangle (1, 2), \langle 3, 4, 19 \rangle (1, 2), \langle 3, 7, 8 \rangle (1, 2)$ $\langle 3, 7, 15 \rangle (1, 2), \langle 3, 8, 15 \rangle (1, 2), \langle 3, 11, 19 \rangle (1, 2)$ $\langle 3, 11, 24 \rangle (1, 2), \langle 3, 11, 51 \rangle (1, 2), \langle 4, 7, 19 \rangle (1, 2)$ $\langle 4, 7, 20 \rangle (1, 2), \langle 4, 7, 52 \rangle (1, 2), \langle 4, 8, 11 \rangle (1, 2)$ $\langle 4, 8, 20 \rangle (1, 1), \langle 7, 8, 35 \rangle (1, 2)$
$(2^*, 2^*, 2^*, 3)$	$\langle \chi_3, \chi_4, \chi_7^3, \chi_7^2 \rangle (1, 2), \langle \chi_3, \chi_7^3, \chi_3 \chi_5^2, \chi_7^2 \rangle (1, 2)$
$(4^*, 2^*, 2^*)$	$\langle \chi_5, \chi_4, \chi_4 \psi_8 \rangle (1, 2), \langle \chi_5, \chi_3, \chi_4 \rangle (1, 2), \langle \chi_5, \chi_3, \chi_7^3 \rangle (1, 2)$ $\langle \chi_5, \chi_3, \chi_4 \psi_8 \rangle (1, 2), \langle \chi_5, \chi_4, \chi_7^3 \rangle (1, 2)$ $\langle \chi_4 \psi_{16}, \chi_3, \chi_4 \rangle (1, 2), \langle \chi_4 \psi_{16}, \chi_4, \chi_4 \chi_5^2 \rangle (1, 1)$
$(4^*, 2)$	$\langle \chi_5, \psi_8 \rangle (1, 1), \langle \chi_5, \chi_{13}^6 \rangle (1, 1), \langle \chi_5, \chi_{17}^8 \rangle (1, 1)$ $\langle \chi_{13}^3, \chi_5^2 \rangle (1, 1), \langle \chi_{13}^3, \psi_8 \rangle (1, 1), \langle \chi_4 \psi_{16}, \chi_5^2 \rangle (1, 1)$
$(4^*, 4^*)$	$\langle \chi_5, \chi_{13}^3 \rangle (1, 2), \langle \chi_5, \chi_4 \psi_{16} \rangle (1, 2)$

Theorem 2. *There are exactly 107 imaginary abelian number fields with relative class numbers ≤ 4 such that the complex conjugation is the square of some automorphism of N : 58 out of them are non-cyclic number fields. These fields are given in Table II.*

TABLE II. The imaginary non-cyclic abelian number fields with relative class number 2 and 4. The fields with non-trivial cyclic ideal class group of 2-power orders are mentioned with \langle, \rangle^c .

Type	h_N^-	Field N (h_{N^+}, Q_N)
$(4^*, 2)$	2	$\langle \chi_4 \psi_{16}, \chi_3 \chi_4 \rangle^c(1, 2)$, $\langle \chi_5, \chi_3 \chi_4 \rangle^c(1, 1)$ $\langle \chi_5, \chi_3 \chi_7^3 \rangle^c(1, 1)$, $\langle \chi_5, \chi_3 \chi_4 \psi_8 \rangle^c(1, 1)$ $\langle \chi_5, \chi_4 \chi_7^3 \rangle^c(1, 1)$, $\langle \chi_5, \chi_{29}^{14} \rangle^c(2, 1)$ $\langle \chi_{13}^3, \chi_{17}^8 \rangle^c(1, 1)$, $\langle \chi_{29}^7, \chi_5^2 \rangle^c(2, 1)$ $\langle \chi_5 \psi_8, \psi_8 \chi_{13}^6 \rangle^c(4, 1)$, $\langle \chi_5 \psi_8, \psi_8 \chi_{17}^8 \rangle^c(2, 1)$ $\langle \chi_5 \chi_{13}^6, \chi_{13}^6 \chi_{17}^8 \rangle^c(2, 1)$, $\langle \chi_5^2 \chi_{13}^3, \chi_5^2 \psi_8 \rangle^c(4, 1)$ $\langle \chi_3 \psi_{16}, \chi_3 \chi_4 \chi_5^2 \rangle(2, 1)$
	4	$\langle \chi_4 \psi_{16}, \chi_4 \chi_7^3 \rangle(1, 2)$, $\langle \chi_4 \psi_{16}, \chi_{17}^8 \rangle(1, 1)$ $\langle \chi_3 \psi_{16}, \chi_3 \chi_7^3 \rangle(1, 1)$, $\langle \chi_4 \psi_{16} \chi_5^2, \chi_4 \chi_5^2 \chi_7^3 \rangle(2, 1)$ $\langle \chi_5, \chi_3 \chi_{11}^5 \rangle(1, 1)$, $\langle \chi_5, \chi_{41}^{20} \rangle(1, 1)$ $\langle \chi_5, \chi_4 \chi_{11}^5 \rangle(1, 1)$, $\langle \chi_5, \chi_3 \chi_{19}^9 \rangle(1, 1)$ $\langle \chi_5, \chi_{61}^{30} \rangle(1, 1)$, $\langle \chi_{13}^3, \chi_3 \chi_4 \rangle(1, 1)$ $\langle \chi_{13}^3, \chi_3 \chi_7^3 \rangle(1, 1)$, $\langle \chi_{13}^3, \chi_{29}^{14} \rangle(1, 1)$ $\langle \chi_5 \psi_8, \chi_3 \chi_4 \psi_8 \rangle(1, 1)$, $\langle \chi_5 \psi_8, \chi_3 \chi_7^3 \psi_8 \rangle(2, 1)$ $\langle \chi_5 \psi_8, \chi_3 \chi_4 \rangle(1, 1)$, $\langle \chi_5 \psi_8, \chi_4 \psi_8 \chi_7^3 \rangle(1, 1)$ $\langle \chi_5 \psi_8, \psi_8 \chi_{29}^{14} \rangle(4, 1)$, $\langle \chi_5 \chi_{13}^6, \chi_3 \chi_4 \chi_{13}^6 \rangle(2, 1)$ $\langle \chi_5 \chi_{13}^6, \chi_3 \chi_7^3 \chi_{13}^6 \rangle(2, 1)$, $\langle \chi_5 \chi_{13}^6, \chi_3 \chi_4 \psi_8 \chi_{13}^6 \rangle(2, 1)$ $\langle \chi_5 \chi_{13}^6, \chi_4 \chi_7^3 \chi_{13}^6 \rangle(4, 1)$, $\langle \chi_5 \chi_{13}^6, \chi_{13}^6 \chi_{29}^{14} \rangle(4, 1)$ $\langle \chi_5 \chi_{17}^8, \chi_3 \chi_4 \chi_{17}^8 \rangle(2, 1)$, $\langle \chi_5 \chi_{17}^8, \chi_3 \chi_7^3 \chi_{17}^8 \rangle(8, 1)$ $\langle \chi_5 \chi_{17}^8, \chi_3 \chi_4 \psi_8 \chi_{17}^8 \rangle(2, 1)$, $\langle \chi_5 \chi_{17}^8, \chi_4 \chi_7^3 \chi_{17}^8 \rangle(2, 1)$ $\langle \chi_5 \chi_{17}^8, \chi_{17}^8 \chi_{29}^{14} \rangle(4, 1)$, $\langle \chi_5^2 \chi_{13}^3, \chi_5^2 \chi_{17}^8 \rangle(2, 1)$ $\langle \chi_{13}^3 \psi_8, \psi_8 \chi_{17}^8 \rangle(4, 1)$, $\langle \chi_7^3 \chi_{17}^8, \chi_4 \chi_7^3 \rangle(1, 1)$ $\langle \chi_7^3 \chi_{17}^8, \chi_4 \psi_8 \chi_7^3 \rangle(1, 1)$, $\langle \chi_7^3 \chi_{17}^8, \chi_3 \chi_5^2 \chi_7^3 \rangle(8, 1)$
$(4^*, 2, 3)$	4	$\langle \chi_{13}^3, \chi_5^2, \chi_{13}^4 \rangle(1, 1)$
$(8^*, 2)$	4	$\langle \chi_3 \chi_{17}^2, \chi_3 \chi_4 \rangle(1, 1)$
$(4^*, 4^*)$	4	$\langle \chi_5, \chi_{29}^7 \rangle(2, 2)$, $\langle \chi_4 \psi_{16}, \chi_3 \chi_4 \chi_5 \rangle(2, 1)$
$(4^*, 2, 2)$	4	$\langle \chi_5, \psi_8, \chi_3 \chi_4 \rangle(1, 1)$

Corollary 1. *There are exactly 48 imaginary non-quadratic abelian number fields N with cyclic ideal class groups of 2-power orders such that the complex conjugation is the square of some automorphism of N : 20 out of them are non-cyclic number fields. Their class numbers are less than or equal to 8.*

This paper is organized as follows. Section 2 reviews some of the standard facts on imaginary abelian number fields. In Section 3 we briefly sketch our method of computations. Throughout this paper the following notations will be used. For a number field K , let O_K , C_K , d_K , h_K and ζ_K be the ring of integers, the ideal class group, the absolute value of discriminant, the class number and Dedekind zeta function of K , respectively. If K is abelian, let us denote by f_K the conductor of K . If K is a CM-field, we will denote by K^+ , h_K^- , ω_K , Q_K the maximal real subfield, the relative class number, the number of roots of unity in K and the Hasse unit index of K , respectively. For an odd prime p let χ_p be an odd Dirichlet character of conductor p , order $p-1$. For $\rho \geq 2$, let ψ_{p^ρ} be an even primitive Dirichlet character of conductor p^ρ , order $p^{\rho-1}$ with $\psi_{p^\rho}^p = \psi_{p^{\rho-1}}$. For the prime 2, let χ_4 be the odd

Dirichlet quadratic character of conductor 4. When $\rho \geq 3$, let ψ_{2^ρ} be the even primitive Dirichlet character of conductor 2^ρ , order $2^{\rho-2}$ with $\psi_{2^\rho}^2 = \psi_{2^{\rho-1}}$.

2. PRELIMINARIES

In this section we sum up some of the standard facts on imaginary abelian number fields which will be used in the sequel.

Proposition 1. (1) *Let F be an imaginary abelian number field, χ_F the group of primitive Dirichlet characters associated to F and χ_F^- the set of $\chi \in \chi_F$ such that $\chi(-1) = -1$. For a $\chi \in \chi_F$ we denote by f_χ the conductor of χ . We have*

$$h_F^- = Q_F \omega_F \prod_{\chi \in \chi_F^-} \left(-\frac{1}{2} B_{1,\chi} \right),$$

where $B_{1,\chi} = \sum_{a=1}^{f_\chi-1} \chi(a) a / f_\chi$.

(2) *Let $K \subset L$ be two CM-fields. Then h_K^- divides $4h_L^-$. In addition, if $[L : K]$ is odd, then h_K^- divides h_L^- .*

Proof. (1) is the content of [W, Theorem 4. 17]. (2) For the first statement see [Ok, Theorem 1] and [Ho, Theorem 5]. For the second statement see [Lem, Corollary 1] or [LOO, Theorem 5]. □

Proposition 2. *Let K be a CM-field of degree $2n$.*

(1) *We have*

$$h_K^- = \frac{Q_K \omega_K}{(2\pi)^n} \sqrt{\frac{d_K}{d_{K^+}}} \frac{\text{Res}_{s=1}(\zeta_K)}{\text{Res}_{s=1}(\zeta_{K^+})}.$$

(2) *The fact that $\beta \in [1 - 2/\log d_K, 1[$ and $\zeta_K(\beta) \leq 0$ implies*

$$\text{Res}_{s=1}(\zeta_K) \geq \frac{\varepsilon_K}{e} (1 - \beta),$$

where $\varepsilon_K = \max \left(1 - \frac{2\pi n e^{1/n}}{d_K^{1/2n}}, \frac{2}{5} \exp \left(-\frac{2\pi n}{d_K^{1/2n}} \right) \right)$.

(3) (a) *There exists a constant $\mu_k > 0$ such that for any abelian extension K/k of degree m unramified at all the infinite places we have*

$$\text{Res}_{s=1}(\zeta_K) \leq (\text{Res}_{s=1}(\zeta_k))^m \left(\frac{1}{2(m-1)} \log \left(\frac{d_K}{d_k^m} \right) + 2\mu_k \right)^{m-1}.$$

(b) *If k is a real abelian number field of degree $s \geq 2$, then*

$$\mu_k \text{Res}_{s=1}(\zeta_k) \leq \frac{s-1}{2^{s+1}} (\log f_k + 2\mu_{\mathbb{Q}})^s,$$

where $\mu_{\mathbb{Q}} = (2 + \gamma - \log(4\pi)) / 2$, γ is Euler's constant.

Proof. (1) is the content of [W, Chapter 4]. (2) See [Lou2, Proposition A] or [LO, Proposition 9]. (3) is the content of Theorem 1, Corollary 2 and Theorem 11 in [Lou3]. □

Proposition 3 ([Lou4, Lemma (b)]). *Let K be an imaginary cyclic number field of degree 2^n , $n \geq 1$. For a positive integer n we let ζ_n be a primitive n th root of unity. Then, $\omega_K = 2$, except when $K = \mathbb{Q}(\zeta_4)$ (in which case $\omega_K = 4$), or when $2^n + 1$ is prime and $K = \mathbb{Q}(\zeta_{2^n+1})$ (in which case $\omega_K = 2(2^n + 1)$).*

Proposition 4. *Let L be a CM-field. We denote by i_{L/L^+} the homomorphism $C_{L^+} \rightarrow C_L$ induced by mapping an ideal \mathfrak{a} to $\mathfrak{a}O_L$. We write $L = L^+(\sqrt{\alpha})$ for some $\alpha \in O_{L^+}$.*

- (1) ([Ha, Satz 24]) *If L is a cyclic number field, then $Q_L = 1$.*
- (2) ([Lem, Theorem 1]) *Assume that $\omega_L \equiv 2 \pmod 4$.*
 - (a) *If the principal ideal (α) is not a square of an ideal of O_{L^+} , then $Q_L = 1$ and i_{L/L^+} is injective.*
 - (b) *Assume that there is an ideal \mathfrak{b} in O_{L^+} such that $(\alpha) = \mathfrak{b}^2$.*
 - i) $Q_L = 2$, if \mathfrak{b} is principal.
 - ii) $Q_L = 1$ and $\ker(i_{L/L^+}) = \langle [\mathfrak{b}] \rangle$, otherwise.

Proposition 5. *Let N be an imaginary cyclic number field.*

- (1) *If $[N : \mathbb{Q}]$ is a power of 2 and if $h_N^- \leq 16$, then $[N : \mathbb{Q}] \leq 16$.*
- (2) *If $h_N^- \leq 4$, then $[N : \mathbb{Q}] \leq 22$.*

Proof. (1) See [PK2]. (2) See [CK]. □

3. MAIN RESULTS

Let s be a positive integer, r a non-negative integer. For $1 \leq i \leq s$ let m_i be a positive integer, for $1 \leq j \leq r$ let n_j be a power of a prime number. Assume that $m_1 \geq m_2 \geq \dots \geq m_s$ and $n_j < 2^{m_s}$ for each j such that n_j is a power of 2. The imaginary abelian field N is called of type $(2^{m_1^*}, \dots, 2^{m_s^*}, n_1, \dots, n_r)$ if N is a compositum of s imaginary cyclic number fields K_i of degree 2^{m_i} , $1 \leq i \leq s$, and r real cyclic number fields L_j of degree n_j , $1 \leq j \leq r$, such that $K_{i+1} \cap (K_1 \cdots K_i) = \mathbb{Q}$ for each $i = 1, \dots, s-1$ and $L_{j+1} \cap (K_1 \cdots K_s L_1 \cdots L_j) = \mathbb{Q}$ for each $j = 0, \dots, r-1$. Let G be the Galois group of N over \mathbb{Q} . Then G is isomorphic to the direct product $\prod_{i=1}^s \mathbb{Z}/2^{m_i}\mathbb{Z} \prod_{j=1}^r \mathbb{Z}/n_j\mathbb{Z}$.

This section is divided into subsections:

- 3.1. Proof of Theorem 1.
- 3.2. Proof of Theorem 2.
- 3.3. Proof of Corollary 1.

3.1. Proof of Theorem 1. In order to determine all imaginary abelian number fields with relative class number one we proceed as follows. Every imaginary abelian number field with relative class number one is a compositum of a finite number of distinct imaginary cyclic number fields with relative class number 1, 2, or 4. Since all imaginary cyclic number fields with relative class number ≤ 4 are known, there remain only finitely many computations to determine all imaginary abelian number fields with relative class number one ([S2], [S3], [A], [MW], [Lou5], [PK1], [PK2], and [CK]). Assume that N is an imaginary abelian number field of type $(2^{m_1^*}, \dots, 2^{m_s^*}, n_1, \dots, n_r)$ with relative class number one. A primitive Dirichlet character will be called a character in brief. Let τ_1, \dots, τ_s be odd characters of order 2^{m_i} , $1 \leq i \leq s$, and let $\varphi_1, \dots, \varphi_r$ be even characters of order n_j , $1 \leq j \leq r$, such that N is associated with the group $\langle \tau_1, \dots, \tau_s, \varphi_1, \dots, \varphi_r \rangle$. Since $h_N^- = 1$, the subfields M_i associated with $\langle \tau_i \rangle$, $1 \leq i \leq s$, satisfy that $h_{M_1 \cdots M_{i_a}}^- | 4$ for $1 \leq a \leq s$, $i_1, \dots, i_a \in \{1, \dots, s\}$. If n_j is odd, then the cyclic subfields L_i associated with $\langle \tau_i, \varphi_j \rangle$, $1 \leq i \leq s$, satisfy $h_{L_i}^- | 4$. If n_j is a power of 2, $n_j = 2^m$, then the relative class numbers of the cyclic subfields associated with $\langle \tau_i \varphi_j^k \rangle$, $1 \leq i \leq s$, $1 \leq k \leq 2^m$, divide 4. On the other hand, if there is no field

of type $(2^{m_1^*}, 2^{m_2^*}, \dots, 2^{m_s^*}, n_1, \dots, n_r)$ with relative class number dividing 4 such that all imaginary subfields have relative class number dividing 4, then we conclude that there is no field with relative class number one containing a subfield of type $(2^{m_1^*}, \dots, 2^{m_s^*}, n_1, \dots, n_r)$. For the fields N of type $(2^*, 2^*)$ or $(2^*, 2^*, 2^*)$ we determine Q_N using [HY1], [HY2] and [HY3]. For other fields we use Proposition 4. In order to determine whether an ideal \mathfrak{b} is principal or not we use the function `IdealsPrincipal` in KASH [KT]. By Proposition 1.(1) we can easily evaluate $h_{\bar{N}}$. Our computational results are summarized in the following. All fields mentioned below are given in Tables I and II.

Proposition 6. *Let l be an odd prime number.*

- (1) (a) *There is only one field of type $(2^*, 3, 3)$ with relative class number one : this field is associated with $\langle \chi_3, \chi_7^2, \psi_9 \rangle$ and has class number one.*
- (b) *There is no field M with relative class number one such that M contains a subfield of type $(2^*, 3, 3, 3)$ or a subfield of type $(2^*, 3, 9)$.*
- (2) *If $l \geq 5$, then there is no field with relative class number one in which a field of type $(2^*, l, l)$ is contained.*
- (3) *If $l \geq 3$, then there is no field with relative class number one in which a field of type $(4^*, l, l)$ is contained (cf. Proposition 15 below).*

Proposition 7. (1) (a) *There are exactly 147 fields of type $(2^*, 2^*)$ with relative class number one.*

- (b) *There are exactly 34 fields of type $(2^*, 2^*, 3)$ with relative class number one.*
- (c) *There is no field with relative class number one in which a field of type $(2^*, 2^*, 3, 3)$ or type $(2^*, 2^*, 9)$ is contained.*
- (2) (a) *There are 3 fields of type $(2^*, 2^*, 5)$ with relative class number one.*
- (b) *There is no field with relative class number one in which a field of type $(2^*, 2^*, 5, 5)$ is contained.*
- (3) *Let l be an odd prime number ≥ 7 . There is no field with relative class number one in which a field of type $(2^*, 2^*, l)$ is contained.*

Proposition 8. (1) *There are 18 fields of type $(4^*, 2^*)$ with relative class number one.*

- (2) *There are 3 fields of type $(4^*, 2^*, 3)$ with relative class number one.*
- (3) *There is no field with relative class number one containing a subfield of type $(4^*, 2^*, 5)$.*

Proposition 9. (1) *There are three fields of type $(8^*, 2^*)$ with relative class number one.*

- (2) *Let l be an odd prime number. There is no field with relative class number one containing a subfield of type $(8^*, 2^*, l)$.*

Proposition 10. *There is no field with relative class number one containing a field of type $(16^*, 2^*)$.*

Proposition 11. (1) *There are 6 fields of type $(4^*, 2)$ with relative class number one.*

- (2) *Let l be a prime. There is no field with relative class number one containing a subfield of type $(4^*, 2, l)$.*
- (3) *There is no field with relative class number one containing a subfield of type $(8^*, 2)$.*

- (4) *There is no field with relative class number one containing a subfield of type $(16^*, 2)$ (cf. Proposition 19).*
- (5) *There are 2 fields of type $(4^*, 4^*)$ with relative class number one.*

Proposition 12. (1) *There are 17 fields of type $(2^*, 2^*, 2^*)$ with relative class number one.*

- (2) *There are two fields of type $(2^*, 2^*, 2^*, 3)$ with relative class number one.*
- (3) *There is no field with relative class number one containing a subfield of type $(2^*, 2^*, 2^*, l)$ for every prime $l \geq 5$.*

Proposition 13. (1) *There are 7 fields of type $(4^*, 2^*, 2^*)$ with relative class number one.*

- (2) *There is no field with relative class number one containing a subfield of type $(4^*, 2^*, 2^*, l)$ for every odd prime l .*

Proposition 14. *There is no field with relative class number one containing a subfield of type $(2^*, 2^*, 2^*, 2^*)$.*

Theorem 1 follows from Propositions 6–14.

3.2. Proof of Theorem 2.

Lemma 1. *Let N be of type $(2^{m_1^*}, \dots, 2^{m_s^*}, n_1, \dots, n_s)$. The complex conjugation is the square of some automorphism of N if and only if $m_1 \geq \dots \geq m_s \geq 2$.*

Proof. Clear. □

According to [PK2, Proposition 3] it is clear that there is no field N of type $(2^{m_1^*}, \dots, 2^{m_s^*}, n_1, \dots, n_s)$ with relative class number 3 if $m_1 \geq \dots \geq m_s \geq 2$. The content of Theorem 2 is divided into Proposition 15–19 according to the Galois groups. Note that there are 8 fields of type $(4^*, 3)$ with relative class number ≤ 4 : 6 of them have relative class number 1 and 2 of them have relative class number 4. There is exactly one field of type $(4^*, 5)$ with relative class number ≤ 4 : $\mathbb{Q}(\zeta_{25})$. For an odd prime l with $l \geq 7$, there is no field of type $(4^*, l)$ with relative class number ≤ 4 (see [CK]).

Proposition 15. *Let l be an odd prime.*

- (1) *There is no field N with relative class number dividing 4 such that N contains a subfield of type $(4^*, l, l)$.*
- (2) *There is no field N with relative class number dividing 4 such that N contains a subfield of type $(8^*, l)$.*

Proof. (1) Let M be a number field of type $(4^*, l, l)$, φ an odd character of order 4, χ_1, χ_2 two characters of order l such that M is associated with the group $\langle \varphi, \chi_1, \chi_2 \rangle$. Suppose that there exists a field N containing M with $h_{\bar{N}}|4$. By Proposition 1.(2) all subfields of degree $4l$ associated with $\langle \varphi, \chi_1^i \chi_2^j \rangle$, $1 \leq i, j \leq l$ and $i + j \leq 2l$, have relative class number dividing 4. Using the results in [PK2] and [CK] we verify that there is no such characters φ, χ_1, χ_2 . From this we deduce (1).

(2) is proved similarly as (1). □

Proposition 16. (1) *There are 13 fields of type $(4^*, 2)$ with relative class number 2 and 32 fields of type $(4^*, 2)$ with relative class number 4.*

- (2) *There is no field of type $(4^*, 2, 3)$ with relative class number 2 and there is one field of type $(4^*, 2, 3)$ with relative class number 4.*

- (3) *There is no field with relative class number ≤ 4 containing a subfield of type $(4^*, 2, l)$, where l is an odd prime ≥ 5 .*
- (4) *There is no field with relative number ≤ 4 containing a subfield of type $(4^*, 2, 9)$.*
- (5) *There is no field of type $(4^*, 2, 2)$ with relative class number 2. There is one field of type $(4^*, 2, 2)$ with relative class number 4.*
- (6) *There is no field with relative class number ≤ 4 containing a subfield of type $(4^*, 2, 2, l)$, where l is a prime.*

Proof. (1) Let N be a number field of type $(4^*, 2)$, φ an odd character of order 4 and χ an even quadratic character such that N is associated with $\langle \varphi, \chi \rangle$. Let M_1 and M_2 be the quartic subfield associated with $\langle \varphi \rangle$ and $\langle \varphi\chi \rangle$, respectively. Assume that $h_N^-|4$. There are exactly 133 pairs of $\langle \varphi, \varphi\chi \rangle$ such that $h_{M_i}^-|8$, $i = 1$ and 2 . It is sufficient to consider M_i with $h_{M_i}^-|8$, instead of those with $h_{M_i}^-|16$, since $h_N^- = \frac{Q_N}{2} h_{M_1}^- h_{M_2}^-$. For these fields we determine Q_N and h_N^- . Note that if $h_N^-|4$, then $f_{\varphi^2} \in \{5, 8, 13, 17, 29\}$ where f_{φ^2} is the conductor of φ^2 . We will need this remark in 2.(a) below.

(2) The computation consists of two steps : (a) Determine all imaginary cyclic number fields of degree 12 with relative class numbers dividing 16 such that its quartic subfield can be embedded into a field N of type $(4^*, 2)$ with $h_N^-|4$. (b) We consider the composite of N and K , where N is one of the fields found in (1), K one of those obtained in (a).

(a) Let K be an imaginary cyclic number field of degree 12, L the quartic subfield and k the quadratic subfield of K . Assume that $h_K^-|16$ and L can be embedded into a field N of type $(4^*, 2)$ with $h_N^-|4$. Using Proposition 2 we find an upper bound for f_K . According to (1) $f_k \in \{5, 8, 13, 17, 29\}$. For those quadratic fields k we verify that $\zeta_k(s) \leq 0$ for $s \in]0, 1[$. Note that ζ_K/ζ_k is a product of L-functions which come in conjugate pairs. Hence $\zeta_K(s) \leq 0$ for $s \in]0, 1[$. By Proposition (2.2) we have

$$\text{Res}_{s=1}(\zeta_K) \geq \frac{\varepsilon_K}{e} \frac{2}{\log d_K}.$$

Since $f_K^6 \leq d_K \leq f_K^{11}$ ([Mu, Corollary 1], [HH, Lemma 10] or [T, Lemma 1]), we have

$$\begin{aligned} h_K^- &= \frac{Q_K \omega_K}{(2\pi)^6} \sqrt{\frac{d_K}{d_{K^+}}} \frac{\text{Res}_{s=1}(\zeta_K)}{\text{Res}_{s=1}(\zeta_{K^+})} \\ &\geq \frac{2\omega_K \eta_K}{(2\pi)^6} \frac{f_L f_K^2}{e \cdot 11 \log f_K} \frac{1}{\text{Res}_{s=1}(\zeta_{K^+})}, \end{aligned}$$

where $\eta_K = \max\left(1 - \frac{12\pi e^{1/6}}{\sqrt{f_K}}, \frac{2}{5} \exp\left(-\frac{12\pi}{\sqrt{f_K}}\right)\right)$.

According to Proposition 2.(3),

$$\text{Res}_{s=1}(\zeta_{K^+}) \leq \text{Res}_{s=1}(\zeta_k) \left(\frac{1}{4} \text{Res}_{s=1}(\zeta_k) \log \frac{d_{K^+}}{d_k^3} + 2\mu_k \text{Res}_{s=1}(\zeta_k) \right)^2$$

and

$$\mu_k \text{Res}_{s=1}(\zeta_k) \leq \frac{1}{8} (\log f_k + 2 \times 0.0231)^2.$$

Computing $\text{Res}_{s=1}(\zeta_k)$ for these 5 quadratic fields k we verify that $f_K \leq 18000$. For such fields K with $f_K \leq 18000$ we compute h_K^- and verify that there are 4

fields K with $h_K^- = 1$, 2 fields K with $h_K^- = 4$ and 2 fields K with $h_K^- = 8$. There is no field K with $h_K^- = 16$.

(b) Let E be a number field of type $(4^*, 2, 3)$, φ an odd character of order 4, χ an even quadratic character and τ a character of order 3 such that E is associated with the group $\langle \varphi, \chi, \tau \rangle$. Let N be the subfield associated with $\langle \varphi, \chi \rangle$, and K_1 and K_2 those associated with $\langle \varphi, \tau \rangle$ and $\langle \varphi\chi, \tau \rangle$, respectively. If $h_E^- \mid 4$, then N is one of the fields obtained in (1) and K_i , $i = 1$ and 2 , is one of those obtained in (a). There are only two fields E of which the subfields satisfy the above conditions: the fields associated with $\langle \chi_5, \chi_{13}^6, \chi_{13}^4 \rangle$ and $\langle \chi_{13}^3, \chi_5^2, \chi_{13}^4 \rangle$, respectively. The former has relative class number 16 and the latter has relative class number 4.

Similarly, we get (3), (4), (5) and (6). □

By the same reasoning as Proposition 16 we get the following.

- Proposition 17.** (1) *There is no field of type $(4^*, 4^*)$ with relative class number 2 and two fields with relative class number 4.*
 (2) *There is no field with relative class number ≤ 4 containing a subfield of type $(4^*, 4^*, l)$, l a prime.*

- Proposition 18.** (1) *There is no field of type $(8^*, 2)$ with relative class number 2 and one field with relative class number 4.*
 (2) *There is no field with relative class number ≤ 4 containing a subfield of type $(8^*, 2, l)$, l prime.*
 (3) *There is no field with relative class number ≤ 4 containing a subfield of type $(8^*, 4^*)$ or type $(8^*, 4)$.*

Proposition 19. *There is no field with relative class number ≤ 4 containing a subfield of type $(16^*, l)$, l prime.*

3.3. Proof of Corollary 1. According to [Lou1, Theorem 2], if an imaginary non-quadratic abelian number field N such that the complex conjugation is the square of some automorphism of N has cyclic ideal class group of 2-power order, then $h_N^- = 1$ or 2. The class numbers of the maximal real subfield of the fields in Theorems 1 and 2 are obtained from [K], [G], [Li], [Y1] and [KT]. (We note that [KT] is used only for the fields of type $(4, 2)$.) Therefore, we verify that among the fields in Theorem 2 there are 21 imaginary non-cyclic number fields N such that

- (i) $h_N^- = 1$ or 2,
- (ii) $h_{N^+} = 2^a$ for some non-negative integer a .

To determine which of those 21 fields has cyclic ideal class group of 2-power order we use Proposition 4 and Proposition 20 below.

Proposition 20. *Let M be a CM-field such that there exists a cyclic quartic M/K with $K \subset M^+ \subset M$. Let $C_M^{(2)}$ and $C_{M^+}^{(2)}$ be the 2-Sylow subgroups of C_M and C_{M^+} , respectively.*

- (1) *The three following conditions are equivalent :*
- (a) $C_M^{(2)}/i_{M/M^+} \left(C_{M^+}^{(2)} \right)$ *is non-cyclic.*
 - (b) $|C_M^{(2)}/i_{M/M^+} \left(C_{M^+}^{(2)} \right)| \geq 4$.
 - (c) $h_M^- \equiv 0 \pmod 4$ *if i_{M/M^+} is injective and $h_M^- \equiv 0 \pmod 2$ otherwise.*
- If these conditions are verified, then $C_M^{(2)}$ is not cyclic.*

- (2) If $C_M^{(2)}/i_{M/M^+}(C_{M^+}^{(2)})$ is cyclic, then $C_M^{(2)}$ and $C_{M^+}^{(2)}$ are cyclic. In addition, M satisfies one of the three following conditions :
- (a) $Q_M = 1$, i_{M/M^+} is injective and $t_M \leq 2$,
 - (b) $Q_M = 2$ and $t_M \leq 1$,
 - (c) i_{M/M^+} is not injective and $t_M \leq 1$.
- Here t_M is the number of prime ideals ramified in M/M^+ .

Proof. See Lemma II.1 and Proposition II.3 in [Gu]. □

ACKNOWLEDGEMENTS

We are greatly indebted to S. Louboutin for suggesting the problem and several helpful comments. We were notified after submission of this paper that Yamamura has recently completed the determination of the imaginary abelian number fields with relative class number one and class number > 1 ([Y2]).

REFERENCES

- [A] S. Arno, *The imaginary quadratic fields of class number 4*, Acta Arith. 60(1992), 321-334. MR **93b**:11144
- [CK] K. -Y. Chang and S. -H. Kwon, *Class number problem for imaginary cyclic number fields*, J. Number Theory 73(1998), 318-338. MR **99i**:11102
- [G] M. -N. Gras, *Classes et unités des extensions cycliques réelles de degré 4 de \mathbb{Q}* , Ann. Inst. Fourier(Grenoble) 29(1979), no 1, XIV, 107-124. MR **81f**:12003
- [Gu] G. Guerry, *Sur la 2-composante du group des classes de certaines extensions cycliques de degré 2^N* , J. Number Theory 53(1995), 159-172. MR **96j**:11151
- [Ha] H. Hasse, *Über die Klassenzahl abelscher Zahlkörper*, Akademie-Verlag, Berlin, 1952. Reprinted with an introduction by J. Martinet: Springer-verlag, Berlin, 1985. MR **14**:141a; MR **87j**:11122a
- [HH] K. Horie and M. Horie, *CM-fields and exponents of their ideal class groups*, Acta Arith. 55(1990), 157-170. MR **91k**:11098
- [Ho] K. Horie, *On a ratio between relative class numbers*, Math. Z. 211(1992), 505-521. MR **94a**:11171
- [HY1] M. Hirabayashi and K. Yoshino, *Remarks on unit indices of imaginary abelian number fields*, Manuscripta Math. 60(1988), 423-436. MR **89e**:11068
- [HY2] M. Hirabayashi and K. Yoshino, *Remarks on unit indices of imaginary abelian number fields II*, Manuscripta Math. 64(1989), 235-251. MR **90j**:11115
- [HY3] M. Hirabayashi and K. Yoshino, *Unit indices of imaginary abelian number fields of type $(2, 2, 2)$* , J. Number Theory, 34(1990), 346-361. MR **91e**:11125
- [K] T. Kubota, *Über den bizyklischen biquadratischen Zahlkörper*, Nagoya Math. J., 10(1956), 65-85. MR **18**:643e
- [KT] M. Daberkow, C. Fieker, J. Klüners, M. Pohst, K. Roegner and K. Wildanger, *Computational algebra and number theory*, KANT V4, J. Symbolic Comp. 24(1997), 267-283. MR **99g**:11150
- [Lem] F. Lemmermeyer, *Ideal class groups of cyclotomic number fields I*, Acta Arith. 72. 4(1995), 347-359. MR **96h**:11111
- [Li] F. J. van der Linden, *Class number computations of real abelian number fields*, Math. Comp. Vol. 39, No 160(1982), 693-707. MR **84e**:12005
- [LO] S. Louboutin and R. Okazaki, *The class number one problem for some non-abelian normal CM-fields of 2-power degrees*, Proc. London Math. Soc. Vol.76, part 3(1998), 523-548. MR **99c**:11138
- [LOO] S. Louboutin, R. Okazaki and M. Olivier, *The class number one problem for some non-abelian normal CM-fields*, Trans. Amer. Math. Soc. 349(1997), 3657-3678. MR **97k**:11149
- [Lou1] S. Louboutin, *CM-fields with cyclic ideal class groups of 2-power orders*, J. Number Theory 67(1997), 1-10. MR **98h**:11139

- [Lou2] S. Louboutin, *Lower bounds for relative class numbers of CM-fields*, Proc. Amer. Math. Soc. 120(1994), 425-434. MR **94d**:11089
- [Lou3] S. Louboutin, *Upper bounds on $|L(1, \chi)|$ and application*, Canad. J. Math., 50(4),1998, 794-815. MR **99f**:11149
- [Lou4] S. Louboutin, *Determination of all nonquadratic imaginary cyclic number fields of 2-power degrees with ideal class groups of exponents ≤ 2* , Math. Comp. Vol. 64, No. 209(1995), 323-340. MR **95c**:11124
- [Lou5] S. Louboutin, *Minoration au point 1 des fonctions L et détermination des corps sextiques abéliens totalement imaginaires principaux*, Acta. Arith. 62.2(1992),109-124. MR **93h**:11100
- [Mu] M. R. Murty, *An analogue of Artin's conjecture for abelian extensions*, J. Number Theory 18(1984), 241-248. MR **85j**:11161
- [MW] H. L. Montgomery and P.J. Weinberger, *Notes on small class numbers*, Acta Arith. **24** (1973/74) 529-542. MR **50**:9841
- [Ok] R. Okazaki, *Inclusion of CM-fields and divisibility of relative class numbers*, Preprint, 1996, Doshisha Univ.
- [PK1] Y.-H. Park and S.-H. Kwon, *Determination of all imaginary abelian sextic number fields with class number ≤ 11* , Acta Arith. 82. 1(1997), 27-43. MR **98i**:11094
- [PK2] Y.-H. Park and S.-H. Kwon, *Determination of all non-quadratic imaginary cyclic number fields of 2-power degree with class number ≤ 20* , Acta Arith. 83. 3(1998), 211-223. MR **99a**:11125
- [S1] H. M. Stark, *Some effective case of the Brauer-Siegel theorem*, Invent. Math. 23(1974), 135-152. MR **49**:7218
- [S2] H. M. Stark, *A complete determination of the complex quadratic fields of class number one*, Michigan Math. J. 14(1967), 1-27. MR **36**:5102
- [S3] H. M. Stark, *On complex quadratic fields with class number two*, Math. Comp. 29(1975), 289-302. MR **51**:5548
- [T] T. Tatuzawa, *On a theorem of Siegel*, Japanese J. Math.21(1951),163-178. MR **14**:452c
- [W] L. C. Washington, *Introduction to cyclotomic fields*, GTM 83. 2nd Ed., Springer-Verlag, 1996. MR **97h**:11130
- [Y1] K. Yamamura, *The determination of imaginary abelian number fields with class number one*, Math. Comp. 62(1994), 899-921. MR **94g**:11096
- [Y2] K. Yamamura, *Table of the imaginary abelian number fields with relative class number one and class number > 1* , Preprint(1999).

DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, 136-701, SEOUL, KOREA
E-mail address: jang@semi.korea.ac.kr

DEPARTMENT OF MATHEMATICS EDUCATION, KOREA UNIVERSITY, 136-701, SEOUL, KOREA
E-mail address: shkwon@semi.korea.ac.kr