

SUFFICIENT CONDITIONS FOR A LINEAR FUNCTIONAL TO BE MULTIPLICATIVE

K. SEDDIGHI AND M. H. SHIRDARREH HAGHIGHI

(Communicated by David R. Larson)

ABSTRACT. A commutative Banach algebra \mathcal{A} is said to have the $P(k, n)$ property if the following holds: Let M be a closed subspace of finite codimension n such that, for every $x \in M$, the Gelfand transform \hat{x} has at least k distinct zeros in $\Delta(\mathcal{A})$, the maximal ideal space of \mathcal{A} . Then there exists a subset Z of $\Delta(\mathcal{A})$ of cardinality k such that \hat{M} vanishes on Z , the set of common zeros of M . In this paper we show that if $X \subset \mathbf{C}$ is compact and nowhere dense, then $R(X)$, the uniform closure of the space of rational functions with poles off X , has the $P(k, n)$ property for all $k, n \in \mathbf{N}$. We also investigate the $P(k, n)$ property for the algebra of real continuous functions on a compact Hausdorff space.

1. INTRODUCTION

The well known Gleason-Kahane-Zelazko theorem ([6, 10]) states that if \mathcal{A} is any commutative unital Banach algebra and M is a closed subspace with $\text{codim}(M)=1$ such that M does not contain invertible elements, then M is a maximal ideal in \mathcal{A} . In other words if $\varphi \in \mathcal{A}^*$ is such that $\varphi(a) \in \sigma(a)$ for every $a \in \mathcal{A}$, then φ is multiplicative. This theorem has been generalized to higher codimensions (see [1, 12, 13, 7]). To state these generalizations we first need a definition.

Let \mathcal{A} be a commutative complex Banach algebra with identity. We say that \mathcal{A} satisfies the $P(k, n)$ property if the following holds: Let M be a closed subspace of \mathcal{A} of finite codimension n . Suppose that for every $x \in M$, the Gelfand transform \hat{x} has at least k distinct zeros in $\Delta(\mathcal{A})$, the maximal ideal space of \mathcal{A} . Then there exists a subset Z of $\Delta(\mathcal{A})$ of cardinality k such that \hat{x} vanishes on Z for every $x \in M$. We call Z a set of common zeros of M . In fact the hypothesis states that if every $x \in M$ is contained in k maximal ideals $I_1^x, I_2^x, \dots, I_k^x$ (depending on x), then there exist k maximal ideals I_1, \dots, I_k such that every $x \in M$ is contained in I_1, I_2, \dots, I_k .

Note that if for $k > 1$, the $P(k, n)$ holds for $C(X)$ the space of continuous complex valued functions on a compact Hausdorff space X , then every point of X is a G_δ set. We now discuss some background history of the subject.

Received by the editors January 31, 1999 and, in revised form, December 17, 1999.

2000 *Mathematics Subject Classification.* Primary 46J20; Secondary 46J10.

Key words and phrases. Multiplicative linear functional, the $P(k, n)$ property, Banach algebra, maximal ideal.

This research was partially supported by a grant from IPM, The Institute for Studies in Theoretical Physics and Mathematics.

In [12] Warner and Whitely conjectured that if X is a compact Hausdorff space such that each point of X is a G_δ set, then $C(X)$ satisfies the $P(k, n)$ property. This conjecture was settled by K. Jarosz [7]. Garimella and Rao [5] showed that $C^m[a, b]$ and $L^1(\mathbf{R})$ satisfy $P(k, n)$. Chen and Cohen [2] have proved that a selfadjoint regular n -point spectral Banach algebra satisfies $P(k, n)$. Rao [11] removed the spectrality condition in the Chen and Cohen theorem, thus making the result applicable to $C^m[a, b]$. He generalized the above result to any selfadjoint, regular commutative Banach algebra with identity. We would also like to investigate the $P(k, n)$ property. Extensions of the Gleason-Kahane-Zelazko theorem in other directions can be found in [9].

In §2 we generalize some results in [7] and [5] and show that if X is a compact and nowhere dense subset of the plane, then $R(X)$, the uniform closure of the algebra of rational functions with poles off X , has the $P(k, n)$ property for all $k, n \in \mathbf{N}$.

Note that the GKZ theorem does not hold for real Banach algebras. For example let $\mathcal{A} = \text{Re}C[0, 1]$ and $\varphi(f) = \frac{1}{2}(f(0) + f(1))$ for all f in \mathcal{A} . By the intermediate value theorem $\varphi(f) \in \sigma(f)$ for all $f \in \mathcal{A}$ but φ is not multiplicative. This example, in turn, shows that the proof of the GKZ theorem depends heavily on complex analysis techniques. But the GKZ theorem is true for $\text{Re}C(X)$, X compact Hausdorff, if and only if X is assumed to be totally disconnected (see Farnum and Whitely [4]). We may use the definition of the $P(k, n)$ property without any difficulty for real Banach algebras and in §3 we show that if X is a compact Hausdorff and totally disconnected space, then $\text{Re}C(X)$ has the $P(1, n)$ property for all $n > 1$ and also has the $P(k, n)$ property for all $k, n \in \mathbf{N}$, if we further assume that the points of X are G_δ . We also show that the $P(1, n)$ property holds for all unital commutative real Banach algebras with a totally disconnected maximal ideal space.

2. ALGEBRAS OF RATIONAL FUNCTIONS

To prove the $P(1, n)$ property for $C(X)$ where X is a compact Hausdorff space, Jarosz proves a lemma [7, Lemma 1] stating that if X is a compact subset of the real line and p_1, \dots, p_n are polynomials such that each linear combination of them has a zero in X , then there is a common zero for p_1, \dots, p_n in X . Garimella and Rao [5] generalized this result for those closed subsets X of \mathbf{C} that have an empty interior and called this result the *polynomial lemma*. The assumption that X has an empty interior is essential as the following example shows:

Let $p_1 = (iz + 1)^2$ and $p_2 = (z + i)^2$ on $\overline{\mathbf{D}}$ the closed unit disk. We show that every linear combination of p_1 and p_2 has a zero in the disk. To see this note that the Möbius transformation $\frac{iz+1}{z+i}$ sends $\overline{\mathbf{D}} \setminus \{-i\}$ to the lower closed half plane and so the function $h = \frac{(iz+1)^2}{(z+i)^2}$ sends it to the whole plane. Consequently, every linear combination of p_1 and p_2 has a zero in the disk while there is no common zero for p_1 and p_2 in the disk.

This example also shows that the assumption of being finite codimensional is essential in the generalizations of GKZ. See also Jarosz [7, §2].

In the proof of Jarosz and Garimella and Rao for the polynomial lemma, the method of several variables is used. Here we prove a generalization of this lemma that uses only elementary complex analysis and needs no assumption of closedness. We only assume that the closure of X in \mathbf{C} has an empty interior, i.e. X is nowhere dense in \mathbf{C} . But first we need a lemma.

Lemma 2.1. *Let X be a nowhere dense subset of \mathbf{C} and f be a holomorphic function on a neighborhood of X . Then $f(X)$ has an empty interior in \mathbf{C} . Consequently, there are infinitely many complex numbers outside $f(X)$.*

Proof. Denote by \overline{X} the closure of X in \mathbf{C} . By definition, for each $x \in X$ there is a neighborhood of x such that f is holomorphic there. In this proof we restrict ourselves to those neighborhoods V such that f is holomorphic on a neighborhood of \overline{V} . Note that f need not be defined on \overline{X} .

Let x be in X . Two cases are possible:

(i) $f'(x) \neq 0$. In this case there exists a bounded open set V containing x such that f is one-to-one on \overline{V} . Thus \overline{V} is compact and $f_1 = f|_{\overline{V}}$ is invertible and if $f_1(\overline{X} \cap \overline{V}) = U$ or equivalently $f_1^{-1}(U) = \overline{V} \cap \overline{X}$, then U has an empty interior in \mathbf{C} since otherwise f_1^{-1} , being holomorphic and nonconstant on U , sends open sets to open sets, and so $\overline{V} \cap \overline{X}$ and hence \overline{X} would contain an open set, which contradicts the hypothesis. Therefore $f(\overline{V} \cap \overline{X}) = f_1(\overline{V} \cap \overline{X})$ has an empty interior in \mathbf{C} .

(ii) $f'(x) = 0$. In this case again two cases may occur. First, there exists a neighborhood V of x such that $f' = 0$ on \overline{V} . Then f is constant on V and hence $f(\overline{V} \cap \overline{X})$ is a singleton and so is closed and has an empty interior. The second possibility is that there exists a neighborhood V such that f' has a single zero x in V . In this case we may suppose that V is an open ball with center x . Choose closed annuli V_n such that $\overline{V} \setminus \{x\} = \bigcup_{n=1}^{\infty} V_n$. By (i), for all $n = 1, 2, 3, \dots$, $f(V_n \cap \overline{X})$ has an empty interior in \mathbf{C} and therefore $f(\overline{V} \cap \overline{X}) = \bigcup_{n=1}^{\infty} f(V_n \cap \overline{X}) \cup \{f(x)\}$ has an empty interior in \mathbf{C} by the Baire category theorem.

Therefore for each $x \in X$, there exists a suitable neighborhood V of x such that $f(\overline{V} \cap \overline{X})$ has an empty interior in \mathbf{C} . But X can be covered with a countable union of such suitable neighborhoods V . Since for such a neighborhood V , $f(\overline{V} \cap \overline{X})$ is closed and has an empty interior in \mathbf{C} , their countable union also has an empty interior by the Baire category theorem. Hence, $f(X)$, being a subset of this countable union, has an empty interior in \mathbf{C} . □

We also need the following well-known facts from linear algebra.

Lemma 2.2. (i) *If a finite dimensional complex or real linear space is a countable union of subspaces, then one of these subspaces must contain the others.*

(ii) *If a complex or real linear space is a finite union of subspaces, then one of these subspaces must contain the others.* □

Theorem 2.3. *Suppose X is a nowhere dense subset of the plane and h_0, \dots, h_n are functions analytic on a neighborhood of X such that each linear combination of them has a zero in X . If one of these h_j , $0 \leq j \leq n$, has countably many zeros in X , then there exists a common zero for h_0, \dots, h_n in X .*

Proof. Suppose h_0 has countably many zeros in X . Set $Z = \{x \in X : h_0(x) = 0\}$. For a linear combination h of h_0, \dots, h_n consider the function $f = h/h_0$ on $X' = X \setminus Z$. Since X' is also nowhere dense in \mathbf{C} , by Lemma 2.1 there exists a complex number $\beta \neq 0$ such that $f(x) \neq \beta$ for all $x \in X'$. But the equation $h(x) = \beta h_0(x)$ has a solution in X and this solution is not in X' , so that it is in Z . Let M denote the complex linear space of linear combinations of h_0, \dots, h_n , i.e. $M = [h_0, \dots, h_n]$. We have shown that every element in M has a zero in Z . For a point $z \in Z$ let $M_z = \{f \in M : f(z) = 0\}$, so that $M = \bigcup_{z \in Z} M_z$. But Z is a

countable set and by Lemma 2.2 (i), one of these M_z must contain the others, i.e. h_0, \dots, h_n has a common zero in X . \square

Corollary 2.4. *Under the same hypothesis, if each linear combination of h_0, \dots, h_n has at least k zeros in X , then there exist k common zeros for h_0, \dots, h_n in X .*

Proof. There exists a common zero for h_0, \dots, h_n . Remove it from X and continue if necessary. \square

Corollary 2.5. *Suppose A is a linear space of functions on a nowhere dense subset X of the plane each of which is analytic on a neighborhood of X , and has at least k zeros in X . Furthermore assume that one element of A has finitely many zeros in X . Then there exist k common zeros for A in X .*

For example, A can be a space of polynomials or rational functions.

Proof. Let p in A have only finitely many zeros in X . The set $Z = \{x \in X : p(x) = 0\}$ is finite and has cardinality $\geq k$. For an arbitrary element $q \in A$, let $h = q/p$, so by Lemma 2.1, $h(X \setminus Z)$ has empty interior in \mathbf{C} . Thus there exists $\beta \neq 0$ such that $h(x) \neq \beta$ for all $x \in X \setminus Z$. But by the hypothesis, $q - \beta p$ is in A and has k zeros in X and none of the zeros are in $X \setminus Z$; hence all are in Z . Therefore q has at least k zeros in Z . For a subset K of Z of cardinality k , set

$$A_K = \{f \in A : f(x) = 0 \text{ for all } x \in K\}.$$

Thus $A = \bigcup_K A_K$ where K runs over all subsets of Z of cardinality k . But Z is a finite set and has only finitely many subsets of cardinality k . By Lemma 2.2 (ii) one of these A_K must contain the others, i.e. A has k common zeros in X . \square

Now the $P(k, n)$ property for the algebra of rational functions over a compact nowhere dense subset of the plane follows.

Theorem 2.6. *Let $R(X)$ be the uniform closure of the algebra of rational functions $\text{Rat}(X)$ with poles off a compact nowhere dense subset X of the plane. If M is a finite codimensional subspace of $R(X)$ such that every element of it has at least k zeros in X , then there are k common zeros for M in X . That is, the $P(k, n)$ property holds for $R(X)$.*

Proof. Note that the maximal ideal space of $R(X)$ is X . Let M be a finite codimensional subspace of $R(X)$ such that each element of M has at least k zeros in X . Let $M' = M \cap \text{Rat}(X)$. Since M is finite codimensional in $R(X)$, M' would be dense in M . Also each element of M' has at least k zeros in X and hence by Corollary 2.5, M' has k common zeros in X . Now M , being the uniform closure of M' , has k common zeros in X . \square

It follows that the $P(k, n)$ property holds for $C(X)$ whenever $X \subset \mathbf{C}$ is compact and nowhere dense with a connected complement (especially whenever $X \subset \mathbf{R}$) since rational functions are dense in $C(X)$ in this case.

The above proof is also applicable to $C^n[a, b]$ and to $C^\infty[a, b]$, although the latter has no Banach algebra norm, since polynomials are dense in these algebras.

3. WHAT HAPPENS FOR $\text{Re}C(X)$?

As stated in the introduction, the GKZ theorem does not hold in general for real Banach algebras and it is true for $\text{Re}C(X)$, X compact, if and only if X is totally disconnected [4]. In fact, if U is a nontrivial connected component of X , then we have an abundance of examples of continuous linear functionals φ with $\varphi(f) \in \text{Im}(f)$ for all f in $\text{Re}C(X)$ such that φ is not multiplicative.

Let x and y be two distinct points of U and $\varphi(f) = \frac{f(x)+f(y)}{2}$. Then for all f in $\text{Re}C(X)$, $\varphi(f) \in \text{Im}(f)$ by the intermediate value theorem and obviously φ is not multiplicative. If we consider U as a subset of the dual space of $\text{Re}C(X)$ (the points of U will then be the point mass measures at points of U), a deeper observation reveals that each element φ of $\overline{\text{co}}(U)$, the closure of the convex hull of U in the dual space of $\text{Re}C(X)$, has the property $\varphi(f) \in \text{Im}(f)$ for all f in $\text{Re}C(X)$.

Now we show that if X is a totally disconnected compact Hausdorff space, then $P(1, n)$ is true for $\text{Re}C(X)$ for all $n \in \mathbf{N}$.

Theorem 3.1. *If X is a totally disconnected compact Hausdorff space, then $\text{Re}C(X)$ has the $P(1, n)$ property for all $n \in \mathbf{N}$.*

Proof. Let M be a finite codimensional subspace of $\text{Re}C(X)$ consisting only of non-invertible elements. Since X is totally disconnected, X can be considered as a closed subspace of a discrete product $\{0, 1\}^J$ where J is some index set. For a finite subset J_0 of J denote

$$M(J_0) = \{f \in M : f \text{ depends only on the coordinates from } J_0\}$$

and

$$X(J_0) = \{x \in X : f(x) = 0 \text{ for all } f \in M(J_0)\}.$$

Clearly $M(J_0)$ can be considered as a subspace of $\mathbf{R}^{2^{J_0}}$ and since each element of $M(J_0)$ is non-invertible; then $X(J_0)$ would be closed (and hence compact) and nonvoid by Lemma 2.2 (i).

Now for a family J_1, \dots, J_m of finite subsets of J let $J_0 = \bigcup_{i=1}^m J_i$. Thus $X(J_0) \subseteq \bigcap_{i=1}^m X(J_i)$ and since $X(J_0)$ is not empty, then it follows that the class

$$\{X(J_0) : J_0 \text{ is a finite subset of } J\}$$

has the finite intersection property and therefore has a nonempty intersection Z which is compact and is a zero set for the subspace

$$M' = \{f \in M : f \text{ depends only on a finite number of coordinates from } J\}.$$

But the linear subspace

$$Y = \{f \in \text{Re}C(X) : f \text{ depends only on a finite number of coordinates from } J\}$$

is a dense subspace of $\text{Re}C(X)$ and M is finite codimensional in $\text{Re}C(X)$, so that $M' = Y \cap M$ is dense in M and therefore each element of M is zero on Z ; that is, M has at least a common zero in X . □

Now we are ready to prove the $P(k, n)$ property for $\text{Re}C(X)$ if we further assume that the points of X are G_δ . In general, all the points of the space $\{0, 1\}^J$ are not G_δ . For example, if $J = [0, 1]$, then the point $\mathbf{0}$ in $\{0, 1\}^J$, which is zero in all of its coordinates, is not a G_δ set.

Theorem 3.2. *If X is a totally disconnected compact Hausdorff space such that each point of X is a G_δ , then the $P(k, n)$ property holds for $\text{Re}C(X)$ for all $k, n \in \mathbf{N}$.*

Proof. Let M be a subspace of $\text{Re}C(X)$ of codimension n such that each element of M has at least k zeros in X . By Theorem 3.1, there is at least one common zero for M in X . Let $Z = \{x_1, \dots, x_m\}$ denote the set of common zeros for M in X . Since the multiplicative functionals on a commutative Banach algebra are linearly independent, it follows that $m \leq n$. If $k \leq m$ we are done. If $k > m$, put

$$M^\perp = [\delta_1, \dots, \delta_m, \mu_1, \dots, \mu_{m'}]$$

where $m + m' = n$ and δ_j denotes the point mass measure at x_j for $1 \leq j \leq m$. Let

$$M' = \{f \in \text{Re}C(X) : \mu_j(f) = \int f d\mu_j = 0, \text{ for all } j, 1 \leq j \leq m'\}.$$

We may suppose, by adding suitable linear combination of $\delta_1, \dots, \delta_m$ to each μ_j , that $|\mu_j|(Z) = 0$ for $1 \leq j \leq m'$.

Since there is no common zero for M' , by Theorem 3.1 there is a function $f \in M'$ such that $f \neq 0$ everywhere on X . By using the duality notions in the subalgebra

$$\{f \in \text{Re}C(X) : f(Z) = 0\},$$

of $\text{Re}C(X)$, we may choose f_j such that f_j is zero on Z and $\int f_i d\mu_j = \delta_{ij}$, $1 \leq i, j \leq m'$. Let $\epsilon > 0$ be such that $g = |f| - \epsilon \sum_j |f_j|$ is a strictly positive function. Set $s = \sup_{x \in X} g(x)$. Since each μ_j is regular and zero on Z , there exists an open neighborhood V of Z such that $|\mu_j|(V) < \epsilon/s$.

Since Z is a G_δ set, we may choose h in $\text{Re}C(X)$ such that $0 \leq h \leq 1$ on X , h is 1 only on Z and is zero outside V . Let $u = gh(\text{sgn}f)$ (since $f \neq 0$ everywhere on X , $\text{sgn}f$ is continuous). Then

$$|\int u d\mu_j| \leq |\mu_j|(V) \sup_{x \in X} g(x) < (\frac{\epsilon}{s})s = \epsilon.$$

If we consider the function $w = f - u + \sum_j \mu_j(u)f_j$, then a simple computation shows that w is zero on Z and $\int w d\mu_j = 0$, $1 \leq j \leq m'$. Thus w belongs to M . Furthermore, $|w| \geq |f| - |u| - \epsilon \sum_j |f_j| = (1 - h)g > 0$, outside Z . Hence $w \in M$ has exactly m zeros which is absurd. □

Corollary 3.3. *If X is a totally disconnected compact Hausdorff space with points G_δ and M is a subspace of codimension n in $\text{Re}C(X)$ such that each element of M has at least n zeros in X , then M is an ideal.* □

In [11, Theorem (0.4)], Rao proved that a complex unital commutative Banach algebra \mathcal{A} has the $P(k, n)$ property if and only if its corresponding semisimple algebra $\frac{\mathcal{A}}{\mathcal{B}}$ has this property where \mathcal{B} denotes the Jacobson radical of \mathcal{A} . The proof there is also applicable to real Banach algebras as well. Therefore we conclude the following:

Theorem 3.4. *Let \mathcal{A} be a unital real commutative Banach algebra with a totally disconnected maximal ideal space X . If M is a finite codimensional subspace of \mathcal{A} such that the Gelfand transform of each element of M has a zero in X , then M is contained in a maximal ideal.*

Proof. We can assume that \mathcal{A} is semisimple. Let Λ be the Gelfand map from \mathcal{A} into $\text{Re}C(X)$. Then $N = \overline{\Lambda(M)}$ is finite codimensional in $\text{Re}C(X)$ and each element of N is noninvertible in $\text{Re}C(X)$. So by Theorem 3.1 there exists a common zero for N in X and this is a common zero for M . Consequently M is contained in a maximal ideal. \square

4. SOME FURTHER RESULTS

In 1991, K. Jarosz [9] posed the following problems:

Problem 1. Does any commutative complex unital Banach algebra have the $P(1, n)$ property for $n \geq 2$?

Problem 2. In particular does the disk algebra have the $P(1, n)$ property for $n \geq 2$?

Problem 3. If some element f of a commutative complex Banach algebra \mathcal{A} does not belong to any regular maximal ideal of \mathcal{A} , does \mathcal{A} have the $P(1, n)$ property for $n \geq 1$?

Problem 4. If the points of the maximal ideal space of such an algebra \mathcal{A} are G_δ , does \mathcal{A} have the $P(k, n)$ property for all positive integers k and n ?

However, in 1989, Rao [11] showed that $P(2, 3)$ does not hold for $C^1(B)$, the algebra of all continuously differentiable functions on B where B denotes the closed unit ball of \mathbf{R}^3 . So the fourth problem of Jarosz is settled negatively. We give this example and modify it to give a negative answer to the third problem. However, the first two problems are still open.

Example 4.1 ([11]). Let $\mathcal{A} = C^1(B)$. It is clear that the maximal ideal space of \mathcal{A} is B . Let P and Q be arbitrary points of B with Q in the interior. Consider the three continuous linear functionals ν_1, ν_2 and ν_3 defined by $\nu_1(f) = f(P) + f_x(Q)$, $\nu_2(f) = f(Q)$, and $\nu_3(f) = f_x(Q) + if_y(Q)$ for all $f \in \mathcal{A}$, where x, y, z denote the coordinates of a point in \mathbf{R}^3 . We show that each element of the subspace

$$M = {}^\perp [\nu_1, \nu_2, \nu_3] = \{f \in \mathcal{A} : \nu_1(f) = \nu_2(f) = \nu_3(f) = 0\}$$

has at least two distinct zeros in B while M has only one common zero Q .

Let $f \in M$. If $f(P) = f(Q) = 0$, we are done. If $f(P) \neq 0$, then also by the definition of M , $f_x(Q)$ and $f_y(Q)$ are not equal to zero. Write $f = g + ih$ where g and h are real valued functions in \mathcal{A} . Consequently $g_x(Q) = h_y(Q)$ and $g_y(Q) = -h_x(Q)$. Since $f_x(Q) \neq 0$, the previous two quantities are not equal to zero, i.e. $g_x(Q) = h_y(Q) \neq 0$. Therefore, by the implicit function theorem, there are two 2-dimensional C^1 manifolds passing through Q such that $g = 0$ on one of them and $h = 0$ on the other. So on their intersection, which contains at least a C^1 curve passing through Q , f is identically zero.

To show that Problem 3 of Jarosz has also a negative answer, we modify the above example.

Example 4.2. We use the same notations as in Example 4.1. Put

$$\mathcal{A}_0 = \{f \in \mathcal{A} : f(Q) = 0\}.$$

It is not difficult to see that \mathcal{A}_0 is a non-unital commutative Banach algebra with the maximal ideal space $B' = B \setminus \{Q\}$ and there are functions in \mathcal{A}_0 which are nonzero on B' , i.e. do not belong to any regular maximal ideal of \mathcal{A}_0 . Now put

$$M_0 = {}^\perp [\nu_1, \nu_3] = \{f \in \mathcal{A}_0 : \nu_1(f) = \nu_3(f) = 0\}$$

in \mathcal{A}_0 . Then M_0 is of codimension 2 in \mathcal{A}_0 and as in the above we see that every element of M_0 has at least one zero in B' but there is no common zero for M_0 in B' . Therefore $P(1, 2)$ does not hold for \mathcal{A}_0 and the answer to the third problem would also be negative.

Finally we treat the most important type of non-unital commutative Banach algebras $C_0(X)$, the algebra of all continuous complex-valued functions on a locally compact Hausdorff space X vanishing at infinity.

If X is not σ -compact, then every element of $C_0(X)$ has infinitely many zeros in X . So, it does not make any sense to speak about the $P(k, n)$ property in this case. But if X is assumed to be σ -compact the $P(1, n)$ property follows for $C_0(X)$ from Theorem 2 of [8] for all $n \in \mathbf{N}$. If we further assume that all the points of X are G_δ , the $P(k, n)$ property follows for $C_0(X)$ from [7] for all $k, n \in \mathbf{N}$ by only considering the natural embedding of $C_0(X)$ into $C(X_\infty)$, where X_∞ denotes the one point compactification of X .

Here we present a simple proof of $P(1, 1)$ for $C_0(X)$ that is independent of [8]:

Theorem 4.3. *Let X be a locally compact Hausdorff space that is σ -compact. Then $C_0(X)$, the space of all complex valued functions on X vanishing at infinity, satisfies $P(1, 1)$.*

Proof. First we show that there exists a positive function f_0 in $C_0(X)$. Write $X = \bigcup_{n=1}^{\infty} K_n$, where K_n is compact for each $n \geq 1$. By the Urysohn lemma we find a sequence $g_n \in C_c(X)$ such that $0 \leq g_n \leq 1$ and $g_n = 1$ on K_n . Set $f_0 = \sum_{i=1}^{\infty} \frac{g_i}{2^i}$. Then $f_0 \in C_0(X)$ and $f_0 > 0$ on X .

Now, let M be a closed subspace of $C_0(X)$ of codimension 1 such that every $f \in M$ has a zero in X . Then $M^\perp = [\mu_0]$ for some regular Borel measure μ_0 on X . Hence

$$M = \{f \in C_0(X) : \int f d\mu_0 = 0\}.$$

Note that $f_0 \notin M$ because every element of M vanishes at some point of X . Let $\alpha = \int f_0 d\mu_0$. Clearly $\alpha \neq 0$. Replacing μ_0 by $\alpha^{-1}\mu_0$ we obtain a new $\mu_0 \perp M$ such that $\int f_0 d\mu_0 = 1$. Now given $f \in C_0(X)$ we write $g = f - (\int f d\mu_0)f_0$. Then $\int g d\mu_0 = \int f d\mu_0 - \int f d\mu_0 = 0$. Therefore $g \in M$. Hence there exists $x_0 \in X$ such that $g(x_0) = 0$, i.e. $\int f d\mu_0 = f(x_0)/f_0(x_0)$. Now if $f \geq 0$, then $\int f d\mu_0 \geq 0$; hence $\mu_0 \geq 0$.

We have shown that $\int f d\mu_0 \in \text{Im}(f/f_0)$ for every $f \in C_0(X)$. Replacing f by ff_0 we get $\int ff_0 d\mu_0 \in \text{Im}(f)$, $f \in C_0(X)$. From this we conclude that $f_0 d\mu_0 = d\delta_v$, the point mass at some $v \in X$ [3, Lemma 2.5]. Write $E = X \setminus \{v\}$. Then $\int_E f_0 d\mu_0 = 0$; hence $f_0 = 0$ a.e. μ_0 on E . Because f_0 is strictly positive, we conclude that $\mu_0(E) = 0$, that is, $\mu_0 = c\delta_v$ for some constant $c > 0$, so that,

$$M = \{f \in C_0(X) : f(v) = 0\}.$$

The proof is now complete. \square

REFERENCES

- [1] C. P. Chen, A generalization of the Gleason-Kahane theorem, *Pacific J. Math.* **107** (1983), 81-87. MR **85d**:46070
- [2] C. P. Chen and J. P. Cohen, Ideals of finite codimension in commutative Banach algebras, manuscript.
- [3] F. Ershad and K. Seddighi, Multiplicative linear functionals in a commutative Banach algebra, *Arch. Math.* **65** (1995), 71-79. MR **96h**:46079
- [4] N. Farnum and R. Whitely, Functionals on real $C(S)$, *Canad. J. Math.* **30** (1978), 490-498. MR **57**:13459
- [5] R. V. Garimella and N. V. Rao, Closed subspaces of finite codimension in some function algebras, *Proc. Amer. Math. Soc.* **101** (1987), 657-661. MR **88m**:46066
- [6] A. M. Gleason, A characterization of maximal ideals, *J. Analyse Math.* **19** (1967), 171-172. MR **35**:4732
- [7] K. Jarosz, Finite codimensional ideals in function algebras, *Trans. Amer. Math. Soc.* **287** (1985), 725-733. MR **86c**:46058
- [8] ———, Finite codimensional ideals in Banach algebras, *Proc. Amer. Math. Soc.* **101** (1987), 313-316. MR **88h**:46094
- [9] ———, Generalizations of the Gleason-Kahane-Zelazko theorem, *Rocky Mountain J. Math.* **21** (1991), 915-921. MR **92m**:46003
- [10] J. P. Kahane and W. Zelazko, A characterization of maximal ideals in commutative Banach algebras, *Studia Math.* **29** (1968), 339-343. MR **37**:1998
- [11] N. V. Rao, Closed ideals of finite codimension in regular selfadjoint Banach algebras, *J. Funct. Anal.* **82** (1989), 237-258. MR **90g**:46077
- [12] C. R. Warner and R. Whitely, A characterization of regular maximal ideals, *Pacific J. Math.* **30** (1969), 277-281. MR **54**:3420
- [13] ———, Ideals of finite codimension in $C[0, 1]$ and $L^1(\mathbf{R})$, *Proc. Amer. Math. Soc.* **76** (1979), 263-267. MR **81b**:46070

DEPARTMENT OF MATHEMATICS, SHIRAZ UNIVERSITY, SHIRAZ 71454, IRAN

DEPARTMENT OF MATHEMATICS, SHIRAZ UNIVERSITY, SHIRAZ 71454, IRAN

E-mail address: shir@sun01.susc.ac.ir